



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 405/06, A61K 31/35, C07D 407/00, 409/06, 417/12, 409/12, 409/14, 417/06, 405/14, 405/12, 417/14, 413/06	A2	(11) International Publication Number: WO 00/15634 (43) International Publication Date: 23 March 2000 (23.03.00)
(21) International Application Number: PCT/US99/18986 (22) International Filing Date: 18 August 1999 (18.08.99) (30) Priority Data: 60/099,946 11 September 1998 (11.09.98) US (71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BOYER, Frederick, Earl, Jr. [US/US]; 1624 Ironwood Drive, Canton Township, MI 48188 (US). DOMAGALA, John, Michael [US/US]; 47693 Red Run, Canton, MI 48187 (US). ELLSWORTH, Edmund, Lee [US/US]; 6899 Kestrel Court, Brighton, MI 48116 (US). GAJDA, Christopher, Andrew [CA/US]; 2684 Bellwood, Ann Arbor, MI 48104 (US). HAGEN, Susan, Elizabeth [US/US]; 1035 Paddington Road, Canton Township, MI 48187 (US). LOVDAHL, Michael, James [US/US]; 2948 Renfrew Street, Ann Arbor, MI 48105 (US). LUNNEY, Elizabeth, Ann [US/US]; 619 Ridgewood Court, Ann Arbor, MI 48103 (US). MARKOSKI, Larry, James [US/US]; 303 W. Green 202D, Champaign, IL 61820 (US).		JOSYULA, Vara, Prasad, Venkata, Nagendra [IN/US]; 3129 Fawnmeadow Court, Ann Arbor, MI 48105 (US). TAIT, Bradley, Dean [US/US]; 8431 Congress Drive, Canton, MI 48187 (US). (74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al. (81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: HIV PROTEASE INHIBITORS (57) Abstract <p>The present invention relates to novel dihydropyrones with tethered heterocycles having improved pharmacologic properties which potently inhibit the HIV aspartyl protease blocking HIV infectivity. The dihydropyrones are useful in the development of therapies for the treatment of viral infections and diseases, including AIDS. The present invention is also directed to methods of synthesis of the dihydropyrones and intermediates useful in the preparation of the final compounds.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-


HIV PROTEASE INHIBITORS

BACKGROUND OF THE INVENTION

The HIV-protease enzyme is absolutely essential for the replication and dissemination of HIV throughout the body (Navia M.A. and McKeever B.M.,
5 *Ann. New York Acad. Sci.*, 1990;616:73-85) and has become an extremely important target for the design and development of anti-HIV therapeutic agents (von der Helm K., *Biol. Chem.* 1996;377:756-774). Several peptidomimetic HIV protease inhibitors have been successfully developed (such as indinavir, saquinavir, ritonavir, and nelfinavir), which demonstrate significant clinical
10 success in lowering plasma viral load, reducing the onset to AIDS, and decreasing the frequency of opportunistic infections (Deeks S.G., Smith M., Holodniy M., and Kahn J.O., *JAMA.*, 1997;277:145-153).


Yet the current HIV protease inhibitors by their peptidomimetic nature have relatively poor solubility, high biliary excretion, limited bioavailabilities and
15 significant liver metabolism. These drawbacks in turn increase the need for high doses of drug which increases the frequency of various side effects and multiple drug interactions (Barry M., Gibbons S., Back D., and Mulcahy F., *Clin Pharmacokinet.*, 1997;32:194-209). More importantly, resistance to the current HIV protease inhibitors has emerged (Shock H.B., Garsky V.M., and Kuo L.,
20 *J. Biol. Chem.*, 1996;271:31957-31963) resulting in treatment failures (Fatkenheuer G., Theisen A., Rockstroh J., Grabow T., et al., *AIDS*, 1997;11:F113-F116). From this discussion, it is apparent that while HIV protease is an excellent antiviral target for the treatment of HIV infection and AIDS, there is a critical need to identify non-peptide inhibitors with improved pharmacological
25 properties and which are not cross resistant with the current drugs (Wallace R.W., *DDT*, 1997;2:83-84).

United States Patent Number 5,789,440 recites non-peptidic HIV protease inhibitors of formula A



A

The patent application is hereby incorporated by reference. Excellent HIV protease inhibition was achieved, but the antiviral activity at the cellular level was in some cases less than desired for an ideal therapeutic agent due to poor overall pharmacological properties (Tummino P.J., Vara Prasad J.V.N., Ferguson D., Nauhan C., et al., *BioOrganic and Med. Chem.*, 1996;4:1401-1410). These efforts however led to a core structure B where R₁ and R₂ were alkyl groups filling the




B

10

S₁' and S₂' pockets, respectively, and the phenyl of the phenethyl group at C₆ filled the S₂ pocket very efficiently. This core structure was recognized as a valuable platform for additional study (Tait B.D., Hagen S., Domagala J.M., Ellsworth E.L., et al., *J. Med. Chem.*, 1997;40:3781-3792).

15

Additional dihydropyrones C were reported when it was unexpectedly discovered that certain polar groups judiciously placed at R₁-R₅ led to greatly



C

-3-

improved antiviral cellular activity. See United States Patent Application Number 08/883,743. The patent application is hereby incorporated by reference. Among the preferred compounds were those where R₁ and R₅ were OH, NH₂, or CH₂OH. In such cases, the preferred R₄ included a small alkyl chain or ring and
5 R₆ was methyl. In addition to improved cellular antiviral activity, the compounds also showed good pharmacokinetics in animals relative to the non-polar substituted compounds. These compounds were also not cross resistant with current HIV Protease inhibitors (Hagen S.E., Vara Prasad J.V.N., Boyer F.E., Domagala J.M., et al., *J. Med.*, 1997;40:3707-3711; Vander Roest S., Wold S.,
10 and Saunders J., *37th Interscience Conference on Antimicrobial Agents and Chemotherapy*, September 28 - October 1, 1997, Toronto, Canada. Abstract I-84; Domagala J.M., Boyer F., Ellsworth E., Gajda C., et al., *5th Conference on Retroviruses and Opportunistic Infections*, February 1-5, 1998, Chicago, IL. Abstract 638).

15 While the compounds C were notable for their improved pharmacological properties relative to the non-polar substituted core molecules B, these highly favorable properties were conferred directly by the use of OH, NH₂ and NR₂ groups placed on the lipophilic rings. The rings themselves were important for binding to the enzyme "pockets" and for holding the t-butyl group and the
20 groups R₁-R₃ and R₅ in their proper places within the enzyme's active site.

It is well-known in the pharmacological sciences that OH and NH₂ groups, especially phenols and anilines, offer distinct metabolic sites resulting in deactivation of the drug and more rapid clearance. In particular, phenols may be glucuronidated and amines and anilines are substrates for rapid acetylation
25 (Goodman L.S. and Gilman A., *The Pharmacological Basis of Therapeutics*, Permagon Press, New York, NY 1985:13-16). Such modifications generally inactivate the drug by preventing its binding to the structurally stringent active site of the enzyme. The modifications also reduce the plasma level of the active agent (Caldwell J., in *Concepts in Drug Metabolism*, edited by Jenner P. and Testa B.,
30 Marcel Dekker, New York, NY, Part A, 1980:235-238). Another suggested problem with amines and anilines is their possible oxidation to electropositive

-4-

nitrogen species which have mutagenic potential (Sobels F.H., *Mut. Res.*, 1985;157:107-110; Bus J.S. and Popp J.A., *Fd Chem. Toxic*, 1987;25:619-626; Rodrigues-Arnaiz R. and Aranda J.H., *Env. Mol. Mutagenesis*, 1994;24:75-79). Thus, while the polar groups are vital for good antiviral efficacy, solubility, and oral absorption, they also present sites for metabolism and possible mutagenesis.

This hereby incorporates by reference 5888L1-01-TMC filed on even date herewith now United States Patent Application Serial Number 60/099,944 filed September 11, 1998, entitled "A Method of Making Dihydropyrone HIV Protease Inhibitors" by Victor Fedij, et al.

SUMMARY OF THE INVENTION

The present invention relates to the extraordinary discovery that the phenyl groups bearing the important polar groups (especially the OH and NH₂ groups) in formula C above, the very polar groups that improved cellular antiviral activity and improved pharmacokinetics in animals relative to compounds of formula B, can themselves be replaced by certain selected heterocycles. These heterocycles bind to the enzyme in the manner of the phenyls to preserve essential enzymatic activity, but additionally present the polar atoms of the heterocycle in such a way as to maintain cellular antiviral activity without requiring peripheral OH and NH₂ groups. Since these compounds have fewer peripheral OH and NH₂ groups, they have even greater pharmacokinetic improvements and less potential for amine based toxicity.

The dihydropyrones with selected heterocycles replacing the phenyls bearing polar substituents are useful in the development of treatments for infections caused by viruses, especially by retroviruses that rely on aspartyl protease activities for replication and infectivity. One such virus is HIV. These dihydropyrones provide higher plasma levels in animals and man due to reduced metabolism and/or clearance, and they provide less toxicity risk due to the removal of certain reactive anilino functionalities. For these reasons, the compounds of this invention are very useful for lowering viral load in individuals


-5-

infected with HIV. Furthermore, the compounds reduce diseases and syndromes associated with viral pathogenesis. One such syndrome is AIDS.

The compounds of the instant invention will be useful according to: Guidelines for the Use of Antiretroviral Agents in HIV-Infected Adults and Adolescents, supplement to the *J. of the International Association of Physicians in AIDS Care*, Supp. No. 1, Vol. 5, pp 4-26 and David A. Katzenstein, Antiretroviral Therapy for HIV: What to do in 1999. *The Journal of Critical Illness*, April 1999;14(4):196.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to compounds or pharmaceutically acceptable salts thereof, of Formula I



R_1 is H, a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R_2 is H, a straight or branched alkyl of 1-5 carbons;

R_3 is H, $(CR'_2)_nOR$, $(CR'_2)_nN(R)_2$, $(CR'_2)_nNR'COR$, $(CR'_2)_nCO_2R$, $(CR'_2)_nOCOR$, $(CR'_2)_nCON(R)_2$, $(CR'_2)_nOCON(R)_2$, $(CR'_2)_nR$, $(CR'_2)_nNR'CON(R)_2$, $(CR'_2)_nNR'CO_2R$, $(CR'_2)_nOSO_2N(R)_2$, $(CR'_2)_nNR'SO_2OR$, $(CR'_2)_nNR'SO_2N(R)_2$, $(CR'_2)_nOSO_2R$, $(CR'_2)_nNR'SO_2R$, $(CR'_2)_nSO_pR$, $(CR'_2)_nNR'CSN(R)_2$, $(CR'_2)_nNR'C(NR')N(R)_2$, $(CR'_2)_nSO_2N(R)_2$, $(CR'_2)_nC(NR')N(R)_2$, $(CR'_2)_nCOR$, $O(CR'_2)_mOR$, $NR(CR'_2)_mOR$, F, Cl, Br, CF_3 , CN, or =O;

R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of 1-6 carbons, a cycloalkyl of 3-6 carbons, $(CR'_2)_nOR$, $(CR'_2)_nN(R)_2$, F, Cl, Br, CN, CF_3 ,

-6-

- $=O$, $(CR'_2)_pNR'COR$, $(CR'_2)_pSO_pR$, $(CR'_2)_pR$, $(CR'_2)_pOCOR$,
 $O(CR'_2)_mOR$, $NR(CR'_2)_mOR$, $(CR'_2)_pNR'CON(R)_2$,
 $(CR'_2)_pOCON(R)_2$, $(CR'_2)_pNR'CO_2R$, $(CR'_2)_pCOR$, $(CR'_2)_pCO_2R$,
 $(CR'_2)_pCON(R)_2$, $(CR'_2)_pNR'SO_2R$, $(CR'_2)_pSO_2N(R)_2$,
5 $(CR'_2)_pNR'SO_2OR$, $(CR'_2)_pOSO_2N(R)_2$, $(CR'_2)_pNR'SO_2N(R)_2$,
 $(CR'_2)_pC(NR')N(R)_2$, $(CR'_2)_pNR'C(NR')N(R)_2$, $(CR'_2)_pHet$;
any two of R_1 - R_3 or R_4 - R_6 may together form a ring of 5-6 total atoms which
may contain 0-3 heteroatoms;
 n is an integer of from 0 to 3;
10 m is an integer of from 2 to 4;
 p is an integer from 0 to 2;
 R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;
 R is independently H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a
 $(CH_2)_n$ heterocycle of 5-6 atoms containing 1-2 heteroatoms and wherein
15 the $(R)_2$ in $N(R)_2$ may form a heterocycle containing the nitrogen, all
optionally substituted by F, Cl, Br, OR' , CN, CO_2R' , $N(R')_2$, $NR'COR'$,
 CF_3 , or $=O$;
 R' is independently H, a straight or branched alkyl of 1-4 carbons, or phenyl;
 R'' is independently H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' ,
20 or $N(R')_2$;
 Ar_1 and Ar_2 are independently phenyl or Het with the proviso that at least one Ar
is Het;
Het is a heterocycle of from 5-6 atoms having from 1-4 heteroatoms or a fused
heterocycle of from 9-10 atoms having 1-3 heteroatoms.
25 A compound which upon administering to a human being converts into a
compound according to Formula I is within the scope of this invention.
Preferred compounds of Formula I are those wherein:
 R_1 is isopropyl or t-butyl;
 R_2 is H, methyl, or ethyl;

-7-

- R_3 is H, $(CR'_2)_nOR$, $(CR'_2)_nN(R)_2$, $(CR'_2)_nNR'COR$, $(CR'_2)_nCO_2R$,
 $(CR'_2)_nOCOR$, $(CR'_2)_nCON(R)_2$, $(CR'_2)_nOCON(R)_2$,
 $(CR'_2)_nNR'CON(R)_2$, $(CR'_2)_nNR'CO_2R$, $(CR'_2)_nOSO_2N(R)_2$,
 $(CR'_2)_nNR'SO_2OR$, $(CR'_2)_nNR'SO_2N(R)_2$, $(CR'_2)_nOSO_2R$,
5 $(CR'_2)_nNR'SO_2R$, $(CR'_2)_nSO_pR$, $(CR'_2)_nNR'CSN(R)_2$, $(CR'_2)_nCOR$,
 $O(CR'_2)_mOR$, $NR(CR'_2)_mOR$, F, Cl, Br, CF_3 , CN, or =O;
- R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of 1-6 carbons, a
 cycloalkyl of 3-6 carbons, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, F, Cl, Br, CN, CF_3 ,
 =O, $(CR'_2)_pNR'COR$, $(CR'_2)_pSO_pR$, $(CR'_2)_pNR'CON(R)_2$,
 10 $(CR'_2)_pOCON(R)_2$, $(CR'_2)_pNR'CO_2R$, $(CR'_2)_pCOR$, $(CR'_2)_pCO_2R$,
 $(CR'_2)_pCON(R)_2$, $(CR'_2)_pNR'SO_2R$, $(CR'_2)_pSO_2N(R)_2$,
 $(CR'_2)_pNR'SO_2OR$, $(CR'_2)_pOSO_2N(R)_2$, $(CR'_2)_pHet$ and any two of
 R_1 - R_3 or R_4 - R_6 may together form a ring of 5-6 total atoms which may
 contain 0-2 heteroatoms;
- 15 n is an integer of 0 to 3;
 m is an integer of 2 to 4;
 p is 0 to 2;
 R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;
 R is H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a
 20 $(CH_2)_n$ heterocycle of 5-6 atoms containing 1-2 heteroatoms and wherein
 the $(R)_2$ in $N(R)_2$ may form a heterocycle containing the nitrogen, all
 optionally substituted by F, Cl, Br, OR' , $N(R')_2$, $NR'COR'$, CF_3 , or =O;
 R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;
 R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or $N(R')_2$;
 25 Ar_1 is phenyl or Het; and
 Ar_2 is Het wherein Het is a heterocycle of from 5-6 atoms having 1-4 heteroatoms
 selected from: furan, pyrrole, thiophene, oxazole, isoxazole, thiazole,
 pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine,
 pyrimidine, pyrazine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine,

-8-

piperazine, piperidine, morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of from 9-10 atoms having from 1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

Other preferred compounds of Formula I are those wherein:

R_1 is isopropyl or t-butyl;

R_2 is H, methyl or ethyl;

R_3 is H, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, $(CH_2)_nNR'COR$, $(CH_2)_nCON(R)_2$,

$(CH_2)_nOCON(R)_2$, $(CH_2)_nNR'CON(R)_2$, $(CH_2)_nNR'CO_2R$,
 $(CH_2)_nOSO_2N(R)_2$, $(CH_2)_nNR'SO_2OR$, $(CH_2)_nNR'SO_2N(R)_2$,
 $(CH_2)_nOSO_2R$, $(CH_2)_nNR'SO_2R$, $(CH_2)_nSO_2R$, $(CH_2)_nNR'CSN(R)_2$,
 $(CH_2)_nCOR$, $O(CH_2)_mOR'$, $NR(CH_2)_mOR'$, or $C(CH_3)_2OR'$;

R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of 1-6 carbons, a

cycloalkyl of 3-6 carbons, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, F, Cl, Br, CN, CF_3 ,
 $=O$, $(CH_2)_pNR'COR$, $(CH_2)_pNR'CON(R)_2$, $(CH_2)_pOCON(R)_2$,
 $(CH_2)_pNR'CO_2R$, $(CH_2)_pCOR$, $(CH_2)_pCON(R)_2$, $(CH_2)_pNR'SO_2R$,
 $(CH_2)_pNR'SO_2OR$, $(CH_2)_pOSO_2N(R)_2$, wherein p is 0 to 2, or R_4 and
 R_5 may together form a ring of 5-6 total atoms which may contain

0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a

$(CH_2)_n$ heterocycle of 5-6 atoms containing 1-2 heteroatoms and wherein
the $(R)_2$ in $N(R)_2$ may be a heterocycle containing the nitrogen, all
optionally substituted by F, Cl, Br, OR' , $N(R')_2$, $NR'COR'$, or $=O$;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or $N(R')_2$;

-9-

Ar₁ is phenyl; and

Ar₂ is a heterocycle of 5-6 atoms having from 1-4 heteroatoms selected from:

- 5 furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine, piperidine, morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of 9-10 atoms having from 1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

Still other preferred compounds of Formula I are those wherein:

R₁ is H, a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R₂ is H, a straight or branched alkyl of 1-5 carbons;

- 15 R₃ is H, (CH₂)_nOR, (CH₂)_nN(R)₂, (CH₂)_nNR'COR, (CH₂)_nCON(R)₂, (CH₂)_nOCON(R)₂, (CH₂)_nNR'CON(R)₂, (CH₂)_nNR'CO₂R, (CH₂)_nOSO₂N(R)₂, (CH₂)_nNR'SO₂OR, (CH₂)_nNR'SO₂N(R)₂, (CH₂)_nOSO₂R, (CH₂)_nNR'SO₂R, (CH₂)_nSO₂R, (CH₂)_nCOR, O(CH₂)_mOR, NR(CH₂)_mOR, C(CH₃)₂OR', F, Cl, Br, CF₃, or =O;

- 20 R₄, R₅, and R₆ are independently H, a straight or branched alkyl of 1-6 carbons, a cycloalkyl of 3-6 carbons, (CH₂)_nOR, (CH₂)_nN(R)₂, F, Cl, Br, =O, (CH₂)_pNR'COR, (CH₂)_pNR'CON(R)₂, (CH₂)_pOCON(R)₂, (CH₂)_pNR'CO₂R, (CH₂)_pCOR, (CH₂)_pCON(R)₂, (CH₂)_pNR'SO₂R, (CH₂)_pNR'SO₂OR, (CH₂)_pOSO₂N(R)₂, wherein p is 0 to 2;

- 25 R₄ and R₅ may together form a ring of 5-6 total atoms which may contain 0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

R₇ is methyl, ethyl, isopropyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl;

-10-

R is H, a straight or branched alkyl of 1-4 carbons, $(\text{CH}_2)_n\text{Ph}$, or a $(\text{CH}_2)_n$ heterocycle of 5-6 atoms containing 1-2 heteroatoms and wherein the $(\text{R})_2$ in $\text{N}(\text{R})_2$ may be a heterocycle containing the nitrogen, all optionally substituted by F, Cl, Br, OR' , $\text{N}(\text{R}')_2$, $\text{NR}'\text{COR}'$, or $=\text{O}$;

5 R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or $\text{N}(\text{R}')_2$;

Ar_1 is phenyl or Het; and

Ar_2 is Het wherein Het is a heterocycle of 5-6 atoms having from 1-4 heteroatoms

selected from: furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine, piperidine, morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of from 9-10 atoms having 1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

Still other preferred compounds of Formula I are those wherein:

R_1 is H, methyl, ethyl, isopropyl, or t-butyl;

R_2 is H or methyl;

20 R_3 is H, $(\text{CH}_2)_n\text{OR}$, $(\text{CH}_2)_n\text{N}(\text{R})_2$, $(\text{CH}_2)_n\text{NR}'\text{COR}$, $(\text{CH}_2)_n\text{CON}(\text{R})_2$, $(\text{CH}_2)_n\text{OCON}(\text{R})_2$, $(\text{CH}_2)_n\text{NR}'\text{CON}(\text{R})_2$, $(\text{CH}_2)_n\text{NR}'\text{CO}_2\text{R}$, $(\text{CH}_2)_n\text{OSO}_2\text{N}(\text{R})_2$, $(\text{CH}_2)_n\text{NR}'\text{SO}_2\text{OR}$, $(\text{CH}_2)_n\text{NR}'\text{SO}_2\text{N}(\text{R})_2$, $(\text{CH}_2)_n\text{OSO}_2\text{R}$, $(\text{CH}_2)_n\text{NR}'\text{SO}_2\text{R}$, $(\text{CH}_2)_n\text{SO}_2\text{R}$, $(\text{CH}_2)_n\text{COR}$, $\text{O}(\text{CH}_2)_m\text{OR}'$, $\text{NR}(\text{CH}_2)_m\text{OR}'$, $\text{C}(\text{CH}_3)_2\text{OR}'$, F, Cl, Br, CF_3 , or $=\text{O}$;

25 R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of 1-6 carbons, a cycloalkyl of 3-6 carbons, $(\text{CH}_2)_n\text{OR}$, $(\text{CH}_2)_n\text{N}(\text{R})_2$, F, Cl, Br, CF_3 , $(\text{CH}_2)_p\text{NR}'\text{COR}$, $(\text{CH}_2)_p\text{NR}'\text{CON}(\text{R})_2$, $(\text{CH}_2)_p\text{OCON}(\text{R})_2$, $(\text{CH}_2)_p\text{NR}'\text{CO}_2\text{R}$, $(\text{CH}_2)_p\text{COR}$, $(\text{CH}_2)_p\text{CON}(\text{R})_2$, $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{R}$, $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{OR}$, $(\text{CH}_2)_p\text{OSO}_2\text{N}(\text{R})_2$, $(\text{CH}_2)_p\text{Het}$ wherein p is 0 to 2;

-11-

any two of R_4 - R_6 may together form a ring of 5-6 total atoms which may contain

0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

5 R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a

$(CH_2)_n$ heterocycle of 5-6 atoms having from 1-2 heteroatoms and wherein the $(R)_2$ in $N(R)_2$ may form a heterocycle having nitrogen, all optionally substituted by F, Cl, Br, OR' , $N(R')_2$, $NR'COR'$, or $=O$;

10 R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or $N(R')_2$;

Ar_1 is a heterocycle of 5-6 atoms having from 1-4 heteroatoms selected from:

15 furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine, piperidine, morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of 9-10 atoms having 1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline; and

20 Ar_2 is phenyl.

More preferred compounds of Formula I are those wherein:

R_1 is isopropyl or t-butyl;

R_2 is H or methyl;

25 R_3 is H, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, $(CH_2)_nOCON(R)_2$, $(CH_2)_nNR'CON(R)_2$, $(CH_2)_nNR'CO_2R$, $(CH_2)_nOSO_2N(R)_2$, $(CH_2)_nNR'SO_2OR$, $(CH_2)_nNR'SO_2N(R)_2$, $(CH_2)_nNR'SO_2R$, $(CH_2)_nSO_2R$, $O(CH_2)_mOR'$, $NR(CH_2)_mOR'$, or $C(CH_3)_2OR'$;

R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of 1-6 carbons, a

30 cycloalkyl of 3-6 carbons, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, F, Cl, Br, $=O$,

-12-

$(\text{CH}_2)_p\text{NR}'\text{COR}$, $(\text{CH}_2)_p\text{NR}'\text{CON}(\text{R})_2$, $(\text{CH}_2)_p\text{OCON}(\text{R})_2$,
 $(\text{CH}_2)_p\text{NR}'\text{CO}_2\text{R}$, $(\text{CH}_2)_p\text{COR}$, $(\text{CH}_2)_p\text{CON}(\text{R})_2$, $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{R}$,
 $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{OR}$, $(\text{CH}_2)_p\text{OSO}_2\text{N}(\text{R})_2$, wherein p is 0 to 2;

R_4 and R_5 may together form a ring of 5-6 total atoms which may contain

5 0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, $(\text{CH}_2)_n\text{Ph}$, or a

10 $(\text{CH}_2)_n$ heterocycle of 5-6 atoms having from 1-2 heteroatoms and wherein
 the $(\text{R})_2$ in $\text{N}(\text{R})_2$ may be a heterocycle containing the nitrogen, all
 optionally substituted by F, Cl, Br, OR' , $\text{N}(\text{R}')_2$, $\text{NR}'\text{COR}'$, or $=\text{O}$;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or $\text{N}(\text{R}')_2$;

15 Ar_1 is phenyl; and

Ar_2 is furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole, tetrazole,
 or pyridine.

Other more preferred compounds of Formula I are those wherein:

R_1 is H, a straight or branched alkyl of 1-6 carbons or a carbocycle of

20 3-6 carbons;

R_2 is H, methyl or ethyl or isopropyl;

R_3 is H, $(\text{CH}_2)_n\text{OR}$, $(\text{CH}_2)_n\text{N}(\text{R})_2$, $(\text{CH}_2)_n\text{NRCOR}$, $(\text{CH}_2)_n\text{CON}(\text{R})_2$,

$(\text{CH}_2)_n\text{OCON}(\text{R})_2$, $(\text{CH}_2)_n\text{NR}'\text{CON}(\text{R})_2$, $(\text{CH}_2)_n\text{NR}'\text{CO}_2\text{R}$,

$(\text{CH}_2)_n\text{OSO}_2\text{N}(\text{R})_2$, $(\text{CH}_2)_n\text{NR}'\text{SO}_2\text{OR}$, $(\text{CH}_2)_n\text{NR}'\text{SO}_2\text{N}(\text{R})_2$,

25 $(\text{CH}_2)_n\text{OSO}_2\text{R}$, $(\text{CH}_2)_n\text{NR}'\text{SO}_2\text{R}$, $(\text{CH}_2)_n\text{SO}_2\text{R}$, $\text{O}(\text{CH}_2)_m\text{OR}$,

$\text{NR}(\text{CH}_2)_m\text{OR}'$, $\text{C}(\text{CH}_3)_2\text{OR}'$, or $=\text{O}$;

R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of 1-6 carbons, a
 cycloalkyl of 1-6 carbons, $(\text{CH}_2)_n\text{OR}$, $(\text{CH}_2)_n\text{N}(\text{R})_2$, F, Cl, Br, $=\text{O}$,

$(\text{CH}_2)_p\text{NR}'\text{COR}$, $(\text{CH}_2)_p\text{NR}'\text{CON}(\text{R})_2$, $(\text{CH}_2)_p\text{OCON}(\text{R})_2$,

-13-

$(\text{CH}_2)_p\text{NR}'\text{CO}_2\text{R}$, $(\text{CH}_2)_p\text{COR}$, $(\text{CH}_2)_p\text{CON}(\text{R})_2$, $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{R}$,
 $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{OR}$, $(\text{CH}_2)_p\text{OSO}_2\text{N}(\text{R})_2$, wherein p is 0 to 2;

R_4 and R_5 may together form a ring of 5-6 total atoms which may contain

0-2 heteroatoms;

5 n is an integer of 0 to 3;

m is an integer of 2 to 4;

R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, $(\text{CH}_2)_n\text{Ph}$, or a

10 $(\text{CH}_2)_n$ heterocycle of 5-6 atoms containing 1-2 heteroatoms and wherein
the $(\text{R})_2$ in $\text{N}(\text{R})_2$ may be a heterocycle having nitrogen, all optionally
substituted by F, Cl, Br, OR' , $\text{N}(\text{R}')_2$, $\text{NR}'\text{COR}'$, or $=\text{O}$;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or $\text{N}(\text{R}')_2$;

Ar_1 is thiophene, thiazole, pyridine, benzothiophene, benzthiazole, benzoxazole,

15 quinoline or isoquinoline; and

Ar_2 is phenyl or Het wherein Het is a heterocycle of 5-6 atoms having from

1-4 heteroatoms selected from: furan, pyrrole, thiophene, oxazole,
isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole,
pyridine, pyridazine, pyrimidine, pyrazine, tetrahydrofuran,
20 tetrahydrothiophene, pyrrolidine, piperazine, piperidine, morpholine,
thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of
9-10 atoms having 1-3 heteroatoms selected from: benzofuran, indole,
indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole,
quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

25 Still other more preferred compounds of formula IV are those wherein:

R_1 is isopropyl or t-butyl;

R_2 is H, methyl, or ethyl;

R_3 is CH_2OH , NH_2 , $\text{OCH}_2\text{CH}_2\text{OH}$, NHCOR , $\text{OSO}_2\text{N}(\text{R})_2$, $\text{NR}'\text{SO}_2\text{OR}$,
 $\text{NR}'\text{SO}_2\text{R}$, or OSO_2R ;

-14-

- R₄, R₅ and R₆ are independently H, a straight or branched alkyl of 1-6 carbons, a cycloalkyl of 1-6 carbons, (CH₂)_nOR, (CH₂)_nN(R)₂, F, Cl, Br, =O, (CH₂)_pNR'COR, (CH₂)_pNR'CON(R)₂, (CH₂)_pOCON(R)₂, (CH₂)_pNR'CO₂R, (CH₂)_pCOR, (CH₂)_pCON(R)₂, (CH₂)_pNR'SO₂R, (CH₂)_pNR'SO₂OR, (CH₂)_pOSO₂N(R)₂, wherein p is 0 to 2;
- R₄ and R₅ may together form a ring of 5-6 total atoms which may contain 0-2 heteroatoms;
- R₇ is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons,
- R is H, a straight or branched alkyl of 1-4 carbons, (CH₂)_nPh, or a (CH₂)_nheterocycle of 5-6 atoms having from 1-2 heteroatoms and wherein the (R)₂ in N(R)₂ may be a heterocycle having nitrogen, all optionally substituted by F, Cl, Br, OR', N(R')₂, NR'COR', or =O;
- R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;
- R'' is H, a straight or branched alkyl of from 1-4 carbons, F, Cl, Br, OR', or N(R')₂;
- Ar₁ is phenyl; and
- Ar₂ is a heterocycle of 5-6 atoms having from 1-4 heteroatoms selected from: furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine, piperidine, morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of 9-10 atoms having from 1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.
- Still other more preferred compounds of Formula I are those wherein;
- R₁ is H, methyl, ethyl, isopropyl, or t-butyl;
- R₂ is H or methyl;

-15-

- R_3 is H, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, $(CH_2)_nNR'COR$, $(CH_2)_nCON(R)_2$,
 $(CH_2)_nOCON(R)_2$, $(CH_2)_nNR'CON(R)$, $(CH_2)_nNR'CO_2R$,
 $(CH_2)_nOSO_2N(R)_2$, $(CH_2)_nNR'SO_2OR$, $(CH_2)_nNR'SO_2N(R)_2$,
 $(CH_2)_nOSO_2R$, $(CH_2)_nNR'SO_2R$, $(CH_2)_nSO_2R$, $(CH_2)_nNR'CSN(R)_2$,
 5 $(CH_2)_nCOR$, $O(CH_2)_mOR$, $NR(CH_2)_mOR'$, $C(CH_3)_2OR'$, F, Cl, Br, or
 $=O$;
 R_4 , R_5 and R_6 are independently H, methyl, ethyl, OH, CH_2OH , CH_2CH_2OH , F,
 Cl, NH_2 ;
 any two R_4 - R_6 may form a ring of 5-6 atoms having from 1-2 heteroatoms;
 10 n is an integer of 0 to 3;
 m is an integer of 2 to 4;
 R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;
 R is H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a
 $(CH_2)_nheterocycle$ of 5-6 atoms having from 1-2 heteroatoms or a
 15 heterocycle having a nitrogen, all optionally substituted by F, Cl, Br, OR' ,
 $N(R')_2$, $NR'COR'$, or $=O$;
 R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;
 R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or $N(R')_2$;
 Ar_1 is furan, thiophene, thiazole, pyridine, imidazole, benzofuran,
 20 benzothiophene, benzimidazole, benzthiazole, quinoline, or isoquinoline;
 and
 Ar_2 is phenyl.

Still other more preferred compounds of formula IV are those wherein:

- R_1 is H, methyl, ethyl, isopropyl, or t-butyl;
 25 R_2 is H or methyl;
 R_3 is H, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, or $=O$;
 R_4 , R_5 , and R_6 are independently H, methyl, OH, CH_2OH , CH_2CH_2OH , NH_2 , or
 F;
 R_7 is H, isopropyl, butyl, sec-butyl, cyclobutyl, cyclopentyl, or cyclohexyl;

-16-

R is H, methyl, ethyl, phenyl, or CH₂Ph and wherein the (R)₂ of N(R)₂ may be a heterocycle having a nitrogen;

R'' is H, F, or CH₃;

Ar₁ is furan, thiophene, thiazole, pyridine, imidazole, benzofuran,

5 benzothiophene, benzimidazole, benzthiazole, quinoline, or isoquinoline;
and

Ar₂ is phenyl.

Most preferred compounds of Formula I are those wherein:

R₁ is H, methyl, ethyl, isopropyl, or t-butyl;

10 R₂ is H, methyl, ethyl, or isopropyl;

R₃ is H, NH₂, OH, CH₂OR, CH₂N(R)₂, CH₂CON(R)₂, CH₂OSO₂N(R)₂,
CH₂NHSO₂OR, CH₂NHSO₂R, CH₂OSO₂R, Cl, Br, or OCH₂CH₂OH;

R₄, R₅, and R₆ are independently H, methyl, ethyl, isopropyl, OH, NH₂, CH₂OR,
CH₂N(R)₂, =O, F, Cl, Br, or CH₂NRCOR;

15 R₇ is methyl, ethyl, isopropyl, cyclopropyl, cyclobutyl, or cyclopentyl;

R is H, methyl, ethyl, Ph, CH₂Ph, and wherein the (R)₂ in N(R)₂ may be a heterocycle having a nitrogen;

R'' is H, F, or CH₃;

Ar₁ is thiophene, thiazole, furan, pyridine, benzothiophene, benzofuran,

20 benzthiazole, benzoxazole, quinoline, or isoquinoline; and

Ar₂ is furan, thiophene, oxazole, isoxazole, imidazole, thiazole, pyrazole,
pyridine, benzofuran, benzothiophene, benzimidazole, benzthiazole,
quinoline, or isoquinoline.

Other most preferred compounds of Formula I are those wherein:

25 R₁ is isopropyl or t-butyl;

R₂ is H, methyl, or ethyl;

R₃ is H, NH₂, OH, CH₂OR, CH₂N(R)₂, CH₂CON(R)₂, OSO₂N(R)₂,
NHSO₂OR, NHSO₂R, OSO₂R, or OCH₂CH₂OH;

-17-

R₄, R₅, and R₆ are independently H, methyl, ethyl, isopropyl, OH, NH₂, CH₂OR, CH₂N(R)₂, =O, F, Cl, Br, or CH₂NRCOR;

R₇ is methyl, ethyl, isopropyl, cyclopropyl, cyclobutyl, or cyclopentyl;

R is H, methyl, ethyl, Ph, CH₂Ph, and wherein the (R)₂ in N(R)₂ may be a

5 heterocycle containing the nitrogen;

R'' is H, F, or CH₃;

Ar₁ is phenyl; and

Ar₂ is furan, thiophene, oxazole, isoxazole, imidazole, thiazole, pyrazole, pyridine, benzofuran, benzothiophene, benzimidazole, benzthiazole, quinoline, or isoquinoline.

10

Still other most preferred compounds of Formula I are those wherein:

R₁ is isopropyl or t-butyl;

R₂ is H, methyl, or ethyl;

R₃ is NH₂, CH₂OH, OCH₂CH₂OH, or CH₂CH₂OH;

15

R₄, R₅, and R₆ are independently H, NH₂, CH₂OH, =O, methyl, ethyl, or isopropyl;

R₇ is isopropyl;

R'' is H, F, or CH₃;

Ar₁ is phenyl; and

20 Ar₂ is furan, thiophene, imidazole, thiazole, pyrazole, or pyridine.

Still other most preferred compounds of Formula I are those wherein:

R₁ is H, methyl, ethyl, isopropyl, or t-butyl;

R₂ is H or methyl;

R₃ is H, CH₂OH, NH₂, or =O;

25

R₄, R₅, and R₆ are independently H, OH, CH₂OH, NH₂, or F;

R₇ is isopropyl, sec-butyl, isobutyl, or cyclopentyl;

R is H, methyl, ethyl, Ph, CH₂Ph, and wherein the (R)₂ in N(R)₂ may be a heterocycle having a nitrogen;

R'' is H, F, or CH₃;

-18-

Ar₁ is furan, thiophene, imidazole, thiazole, pyrazole, or pyridine; and

Ar₂ is phenyl.

Especially preferred compounds of the invention are:

- 5 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-4-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-2-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 10 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-2-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 15 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 20 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 25 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 30 (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one;

-19-

(S)-3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one;

6-[2-(4-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

5 3-(2-Amino-5-isopropyl-benzothiazol-6-yl-sulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-phenethyl-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-yl-sulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one;

10 (S)-3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

(S)-(6-{4-Hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-isopropyl-7-methyl-benzothiazol-2-yl)-carbamic acid methyl ester;

(S)-3-(2-Amino-6-tert-butyl-benzothiazol-4-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

(S)-3-(3-tert-Butyl-benzo[b]thiophen-2-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(3-tert-Butyl-benzo[b]thiophen-2-yl-sulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

4-Hydroxy-3-(5-hydroxymethyl-2-isopropyl-thiophen-3-ylsulfanyl)-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

25 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

30 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

-20-

N-(4-{2-[5-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl]-ethyl}-thiazol-2-yl)-acetamide;

5 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

10 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

15 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-isopropyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

20 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxy-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

25 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(1H-pyrazol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

30 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-[2-(1H-pyrazol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyrimidin-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

-21-

- 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyrimidin-5-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 6-[2-(2-Amino-pyrimidin-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 5 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(2-amino-pyrimidin-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-4-yl-ethyl)-5,6-dihydro-pyran-2-one;
- (S)-4-Hydroxy-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-6-phenethyl-5,6-dihydro-pyran-2-one;
- 10 (S)-4-Hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;
- 3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 15 (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- (S)-3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- (R)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 20 3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;
- 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 25 3-(2-tert-Butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 30 3-[2-tert-Butyl-4-(2-hydroxy-ethyl)-5-methyl-phenylsulfanyl]-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

-22-

5-Trifluoromethyl-pyridine-2-sulfonic acid {5-tert-butyl-4-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-2-methyl-phenyl}-amide; and

5 5-Trifluoromethyl-pyridine-2-sulfonic acid {5-tert-butyl-4-[4-hydroxy-6-isopropyl-2-oxo-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-2-methyl-phenyl}-amide.

Other preferred compounds are:

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(3-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

10 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

15 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-Cyclopentyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

20 3-(4-Amino-2-cyclopentyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-cyclopentyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

25 3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-furan-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(3-hydroxymethyl-furan-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-23-

- 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-furan-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-furan-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 5 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-furan-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 10 3-(2-Cyclopentyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 15 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 20 6-[2-(2-Amino-thiazol-4-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 6-[2-(2-Amino-thiazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 25 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-4-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-5-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 30 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-oxazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

-24-

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

5 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

10 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(1H-benzoimidazol-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(1H-Benzoimidazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(3-amino-1H-indazol-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

15 6-[2-(3-Amino-1H-indazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

25 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(3-amino-1H-indazol-4-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(3-Amino-1H-indazol-4-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

30 3-(2-tert-Butyl-6-hydroxymethyl-5-methyl-pyridin-3-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-5-methyl-pyridin-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-25-

- 3-(3-tert-Butyl-2-hydroxymethyl-6-methyl-pyridin-4-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(5-tert-Butyl-2-methyl-pyridin-4-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 5 3-(4-tert-Butyl-5-hydroxymethyl-thiophen-3-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(4-tert-Butyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 6-[2-(4-Amino-phenyl)-ethyl]-3-(2-tert-butyl-5-methyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 10 3-(2-tert-Butyl-5-methyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 6-[2-(4-Amino-phenyl)-ethyl]-3-(4-tert-butyl-5-hydroxymethyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 15 6-[2-(4-Amino-phenyl)-ethyl]-3-(3-tert-butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(3-tert-Butyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(3-tert-Butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 20 3-(3-tert-Butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 25 3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 30 3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

-26-

3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

N-(7-tert-Butyl-6-{6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-acetamide;

5 N-(7-tert-Butyl-6-{4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-acetamide;

10 N-(7-tert-Butyl-6-{6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-methanesulfonamide;

N-(7-tert-Butyl-6-{4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-methanesulfonamide;

15 3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-6-hydroxymethyl-5-methyl-pyridin-3-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

25 3-(3-tert-Butyl-2-hydroxymethyl-6-methyl-pyridin-4-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-27-

N-{7-tert-Butyl-6-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-4-methyl-benzothiazol-2-yl}-acetamide;


5 N-{7-tert-Butyl-6-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-4-methyl-benzothiazol-2-yl}-methanesulfonamide;

3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

10 3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

(S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one; and
compounds useful as intermediates in the preparation of the final products are:

a compound of Formula 1A



wherein

R₄, R₅, and R₆ are independently H, a straight or branched alkyl of 1-6 carbons, a cycloalkyl of 3-6 carbons, (CR'₂)_nOR, (CR'₂)_nN(R)₂, F, Cl, Br, CN, CF₃, =O, (CR'₂)_pNR'COR, (CR'₂)_pSO_pR, (CR'₂)_pR, (CR'₂)_pOCOR, O(CR'₂)_mOR, NR(CR'₂)_mOR, (CR'₂)_pNR'CON(R)₂, (CR'₂)_pOCON(R)₂, (CR'₂)_pNR'CO₂R, (CR'₂)_pCOR, (CR'₂)_pCO₂R, (CR'₂)_pCON(R)₂, (CR'₂)_pNR'SO₂R, (CR'₂)_pSO₂N(R)₂, (CR'₂)_pNR'SO₂OR, (CR'₂)_pOSO₂N(R)₂, (CR'₂)_pNR'SO₂N(R)₂, (CR'₂)_pC(NR')N(R)₂, (CR'₂)_pNR'C(NR')N(R)₂, (CR'₂)_pHet;
25 any two of R₄-R₆ may together form a ring of 5-6 total atoms which may contain 0-3 heteroatoms;

-28-

n is an integer of from 0 to 3;

p is an integer from 0 to 2;

R₇ is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R is independently H, a straight or branched alkyl of 1-4 carbons, (CH₂)_nPh, or a

5 (CH₂)_nheterocycle of 5-6 atoms containing 1-2 heteroatoms and wherein the (R)₂ in N(R)₂ may form a heterocycle containing the nitrogen, all optionally substituted by F, Cl, Br, OR', CN, CO₂R', N(R')₂, NR'COR', CF₃, or =O;

R' is independently H, a straight or branched alkyl of 1-4 carbons, or phenyl;

10 R'' is independently H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR', or N(R')₂;

Ar₂ is phenyl or Het wherein Het is a heterocycle of from 5-6 atoms having from 1-4 heteroatoms.

Terms useful in describing the instant invention are as follows:

15 The term "alkyl" means a straight or branched hydrocarbon radical having from 1 to 12 carbon atoms unless otherwise specified and includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, undecyl, and dodecyl. The alkyl groups may contain one or more sites of unsaturation such as double or triple
20 carbon-carbon bonds. The alkyl group is unsubstituted or substituted by from 1 to 3 substituents selected from F, Cl, Br, OH, NH₂, CN, NO₂, OCH₃, OCH₂CH₂OH, NHCH₃, or N(CH₃)₂.

The term "cycloalkyl" means a hydrocarbon ring which contains from 3 to 12 carbon atoms unless otherwise specified, for example, cyclopropyl, cyclobutyl, 25 cyclopentyl, cyclohexyl, and adamantyl. Where possible, the cycloalkyl group may contain double bonds. The cycloalkyl ring may be unsubstituted or substituted by from 1 to 3 substituents selected from alkyl, alkoxy, thioalkoxy all as defined herein, hydroxy, thiol, nitro, halogen, amino, formyl, carboxyl, nitrile, -NH-CO-R, -CO-NHR-, -CO₂R, -COR, aryl, or heteroaryl wherein alkyl (R), aryl, 30 and heteroaryl are defined as herein.

The term "carbocycle" means cycloalkyl as defined above.

-29-

The term "heteroatoms" means a nitrogen, sulfur, or oxygen.

The term "heterocycle" means a heterocyclic radical which are 5-6 atoms having 1-4 heteroatoms and selected from 1-4 heteroatoms selected from: furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine, piperidine, morpholine, thiomorpholine, oxolane, dioxane, sulfolane, unsubstituted or substituted by 1 to 2 substituents selected from alkyl as defined above. For heterocycles containing sulfur, the oxidized sulfur heterocycles containing SO or SO₂ groups are also included.

The term "fused heterocycle" refers to a heterocycle that is adjoined as two consecutive positions with a phenyl ring or another heterocycle. such rings may include 2-, 3-, 4-, 5-, 6-, 7-, or 8-quinoliny, 1-, 3-, 4-, 5-, 6-, 7-, or 8-isoquinoliny, 2-, 3-, 4-, 5-, 6-, or 7-indolyl, 2-, 3-, 4-, 5-, 6-, or 7-indoliny, 2-, 3-, 4-, 5-, 6-, or 7-benzo[b]thienyl, 2-, 4-, 5-, 6-, or 7-benzoxazolyl, 2-, 4-, 5-, 6-, or 7-benzimidazolyl, 2-, 4-, 5-, 6-, or 7-benzothiazolyl.

For purposes of the syntheses of the compounds of the present invention, reactive functional groups present in starting materials, reaction intermediates, or reaction products may be protected during chemical reactions using protecting groups which render the reactive functional groups substantially inert to the reaction conditions (see for example, *Protective Groups in Organic Synthesis*, 2nd ed., T. W. Green and P. G. Wuts, John Wiley & Sons, New York, NY 1991). Thus, for example, protecting groups such as the following may be utilized to protect suitable amino, hydroxyl, and other groups of related reactivity: carboxylic acyl groups, such as formyl, acetyl, trifluoroacetyl; alkoxycarbonyl groups, such as ethoxycarbonyl, *t*-butoxycarbonyl (BOC), β,β,β -trichloroethoxycarbonyl (TCEC), β -iodoethoxycarbonyl; aryloxycarbonyl groups, such as benzyloxycarbonyl, *p*-methoxybenzyloxycarbonyl, phenoxycarbonyl; trialkyl silyl groups, such as trimethylsilyl and *t*-butyldimethylsilyl (TBDMS); and groups such as trityl, tetrahydropyranyl, vinyloxycarbonyl, *o*-nitrophenylsulfenyl, diphenylphosphinyl, *p*-toluenesulfonyl, and benzyl may all be utilized. The protecting group may be removed, after completion of the synthetic reaction of

-30-

interest, by procedures known to those skilled in the art. For example, a BOC group may be removed by acidolysis, a trityl group by hydrogenolysis, TBDMS by treatment with fluoride ions, and TCEC by treatment with zinc.

5 Some of the compounds of Formula I are capable of further forming pharmaceutically acceptable acid-addition and/or base salts. All of these forms are within the scope of the present invention.

Pharmaceutically acceptable acid addition salts of the compounds of Formula I include salts derived from nontoxic inorganic acids such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydriodic, hydrofluoric, 10 phosphorous, and the like, as well as the salts derived from nontoxic organic acids, such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, alkanedioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, etc. Such salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, monohydrogenphosphate, 15 dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, trifluoroacetate, propionate, caprylate, isobutyrate, oxalate, malonate, succinates suberate, sebacate, fumarate, maleate, mandelate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, phthalate, benzenesulfonate, toluenesulfonate, phenylacetate, citrate, lactate, maleate, tartrate, 20 methanesulfonate, and the like. Also contemplated are salts of amino acids such as arginate and the like and gluconate, galacturonate (see, for example, Berge S.M., et al., "Pharmaceutical Salts," *Journal of Pharmaceutical Science*, 1977;66:1-19.

The acid addition salt of said basic compounds are prepared by contacting the free base form with a sufficient amount of the desired acid to produce the salt 25 in the conventional manner.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chloroprocaine, 30 choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge, supra., 1977).

-31-

The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner.

5 Certain of the compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms, are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention.

10 Certain of the compounds of the present invention possess one or more chiral centers and each center may exist in the R(D) or S(L) configuration. The present invention includes all enantiomeric and epimeric forms as well as the appropriate mixtures thereof. Configuration drawn is most preferred.

15 The compounds of the present invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds of the present invention can be administered by inhalation, for example, intranasally. Additionally, the compounds of the present invention can be administered transdermally. It will be obvious to those skilled in the art that the following dosage forms may comprise
20 as the active component, either a compound of Formula I or a corresponding pharmaceutically acceptable salt of a compound of Formula I.

25 For preparing pharmaceutical compositions from the compounds of the present invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided active component.

30 In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

-32-

The powders and tablets preferably contain from five or ten to about seventy percent of the active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component with or without other carriers, is surrounded by a carrier, which is thus in association with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted, and the active component is dispersed homogeneously therein, as by stirring. The molten homogenous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water propylene glycol solutions. For parenteral injection, liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing and thickening agents as desired.

Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-known suspending agents.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

-33-

The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is divided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

The quantity of active component in a unit dose preparation may be varied or adjusted from 0.1 mg to 100 mg preferably 0.5 mg to 100 mg according to the particular application and the potency of the active component. The composition can, if desired, also contain other compatible therapeutic agents.

In therapeutic use as antagonists of a retroviral protease, as agents for the treatment of infections caused by a retrovirus including HIV, or as agents for the treatment of diseases due to AIDS, the compounds utilized in the pharmaceutical method of this invention are administered at the initial dosage of about 0.01 mg to about 100 mg/kg daily. A daily dose range of about 0.01 mg to about 10 mg/kg is preferred. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day, if desired.

The compounds of the instant invention, protease inhibitors, can be used as a component of initial patient antiretroviral therapy for HIV as, for example, in the following regimen:

Choose one combination of nucleoside reverse transcriptase inhibitors

Zidovudine + lamivudine

30 Zidovudine + didanosine

Zidovudine + zalcitabine

Didanosine + stavudine

-34-

Didanosine + lamivudine

Stavudine + lamivudine

Abacavir + zidovudine + lamivudine

AND one of the compounds of the instant invention.

- 5 In addition, one may use one of the nucleoside reverse transcriptase inhibitor combinations above with one of the agents below plus a compound of the instant invention.

Nelfinavir

Indinavir

- 10 Saquinavir (soft gel-cap now preferred over hard gel-cap)

Ritonavir

Nevirapine

Delavirdine

Efavirenz

- 15 The compounds of the instant invention can be used in multiple combinations with available antiretroviral drugs for combination therapy. The following are illustrative:

Drug class/agents	Dosage
Nucleoside reverse transcriptase inhibitors	
Abacavir	300 mg bid
Didanosine	200 mg bid (for patients <60 kg, 125 mg bid)
Lamivudine	150 mg bid (for patients <50 kg, 2 mg/kg bid)
Stavudine	40 mg bid (for patients <60 kg, 30 mg/kg bid)
Zalcitabine	0.75 mg tid
Zidovudine	300 mg bid
Non-nucleoside reverse transcriptase inhibitors	
Delavirdine	400 mg tid
Efavirenz	600 mg qd
Nevirapine	200 mg bid
Protease inhibitors	
Indinavir	800 mg q8h
Nelfinavir	750 mg tid
Ritonavir	600 mg bid
Saquinavir	600 mg tid (hard gel-cap formulation), 1200 mg tid (soft gel-cap formulation)

-35-

The compounds of the instant invention can be used with or instead of the following for the treatment of established HIV infection.

<i>Preferred</i>	Strong evidence of clinical benefit and/or sustained suppression of plasma viral load. One choice each from column A and column B. Drugs are listed in random, not priority, order:	
	Column A	Column B
	Indinavir (AI)	ZDV + ddI (AI)
	Nelfinavir (AII)	d4T + ddI (AII)
	Ritonavir (AI)	ZDV + ddC (AI)
	Saquinavir – SGC (AII)	ZDV + 3TC (AI)
	Ritonavir + Saquinavir SGC or HGC (BII)	d4T + 3TC (AII)
	Efavirenz (AII)	
<i>Alternative</i>	Less likely to provide sustained virus suppression, or data inadequate. Nevirapine or delavirdine + two NRTIs (Column B, above) (BII)	

The compounds of the instant invention can be used with the following drugs available through treatment investigational new drug protocols.

Drug	Adefovir (Preveon)	Abacavir (1592-U89)	Amprenavir (Agenerase; 141W94)
Source	Gilead 800-GILEAD-5	Glaxo-Wellcome 800-501-4672	Vertex; Glaxo-Wellcome 800-248-9757
Class	Nucleotide RT Inhibitor	Nucleoside RT Inhibitor	Protease Inhibitor
Usual dose	60 mg po qd or 120 mg po qd + L-carnitine 500 mg po qd	300 mg po bid	1200 mg po bid
Side effects (major)	Proximal renal tubular dysfunction, nausea, elevated LFTs	Hypersensitivity: 2-5% usually in first 4 weeks (fever, nausea, vomiting, morbilliform rash)	Nausea, diarrhea, rash, headache

5



The compounds of the present invention can be prepared according to the various synthetic schemes that follow. Protecting groups may be used when

-36-

appropriate throughout many of the schemes. Although specifically noted in certain schemes, the appropriate use and choice of protecting groups is well-known by one skilled in the art, and is not limited to the specific examples below. It is also understood that such groups not only serve to protect chemically reactive sites, but also to enhance solubility or otherwise change physical properties. A good general reference for protecting group preparation and deprotection is "Protecting Groups in Organic Synthesis" by Theodora Green. A number of general reactions such as oxidations and reductions are not shown in detail but can be done by methods understood by one skilled in the art. General transformations are well-reviewed in "Comprehensive Organic Transformation" by Richard Larock, and the series "Compendium of Organic Synthetic Methods" published by Wiley-Interscience. In general, the starting materials were obtained from commercial sources unless otherwise indicated.

There are three major components of the synthesis of the desired products: the formation of a ketone precursor; the cyclization of that ketone into the dihydropyrone ring system; and the addition of the thiol component to complete the preparation of the compounds of the invention. Schemes 1 through 6 address the various methods for preparation of ketones indicated as C, D, G, and K.

Scheme 1



20

Aldehyde synthesis


The desired aldehydes which can be used in Schemes 2 and 3 are prepared by a number of methods as indicated in Scheme 1. The appropriate ester can be

-37-

reduced to the corresponding alcohols and then oxidized to the aldehyde. Another method involves preparation of an anion with an appropriate base and trapping with a reagent, such as DMF, to prepare the aldehyde.

-38-


Scheme 2



- 5 The desired ketones can be prepared as shown in Schemes 2 and 3. The appropriate α halo ketone can be displaced with triphenylarsine to prepare an arsenate reagent. In a similar manner, triphenylphosphine can be used to prepare a Wittig reagent which will also carry out the reaction shown in Scheme 2. Upon treating the arsenate reagent and the aldehyde with base such as potassium carbonate, the chalcone derivative C is formed. Compound C can then be reduced using hydrogen with an appropriate catalyst to prepare ketone D.

-39-

Scheme 3



Alternatively, the desired enones C can be prepared by reaction of the appropriate methylketones and aldehydes using $Ba(OH)_2$ in EtOH as described in *Synthesis*, 1983:502-504, *An Quim, Ser. C*, 1981;77(2):222-224, and *Pol. J. Chem.*, 1982;56(10-112):1435. In a similar fashion, the enones C can also be prepared by reaction of the appropriate methylketones and aldehydes using NaOH in EtOH as described by Kohler and Chadwell, *Org Synth. Coll.*, 1941;1:78. The relative ratio of aldehyde, ketone, base, and temperature (RT to reflux) varied according to the substitution pattern of the aldehyde and ketone. There have also been reports of conversion of A and B to chalcones C under acidic conditions using H_2SO_4 and acetic anhydride. This may be accomplished by direct treatment with acid or the acid may be used to dehydrate the aldol intermediate if it does not dehydrate during the reaction with base.


The resulting chalcones C were converted into ketones D by reduction as noted previously in Scheme 2. The reduction was generally accomplished by hydrogenation using palladium on barium sulfate at room temperature in tetrahydrofuran. On occasion, overreduction to the alcohol was observed. The isolated alcohol or the crude mixture can be oxidized to the ketone with Jones reagent.

The reduction of the enone to the corresponding ketone can be carried out via a number of alternative methods including but not limited to: metals in ammonia as described in *Org. Reactions*, 1976;23:1-253; ethylene glycol and

-40-

$\text{RuCl}_2[\text{P}(\text{Ph})_3]_3$ as described in *Synthesis*, 1973:359; and copper hydride reagents as described in *J. Amer. Chem. Soc.*, 1974;96:3686.


Scheme 4



- 5 Preparation of the ketone G can be accomplished from an appropriate acid(E) and appropriate nucleophile using the Weinreb methodology described in *Tetrahedron Letters*, 1987:1857. The appropriate acid can be converted to the acid chloride by treatment with neat thionyl chloride or by treatment with oxalyl chloride in methylene chloride with a catalytic amount of DMF. Treatment of the
- 10 acid chloride with N,O-dimethylhydroxylamine with bases such as pyridine or triethylamine in inert solvents such as methylene chloride from 0°C to reflux will give intermediate F. Intermediate F can then be converted to G by treatment with the appropriate nucleophile, such as Grignard reagents, at 0°C to reflux in solvents such as tetrahydrofuran or ether. An alternative method of converting the acid E to
- 15 ketone G is by treatment with alkyl lithiums at -78°C to reflux as described in *J. Med. Chem.*, 1996;39:2659 and *J. Amer. Chem. Soc.* 1970;92:2590. Another route is treatment of the acid chloride with nucleophiles (organomagnesium, copper, cadmium, or zinc reagents) as reviewed in *Org. React.*, 1954;8:28 and *Tetrahedron Lett.*, 1970:4647.


-41-

Scheme 5



An alternative preparation of ketone D can be accomplished by reaction of
 an allylic alcohol I with an appropriate halide H using palladium acetate as
 described in *Tetrahedron*, 1979;35:329 and *Tetrahedron Lett.*, 1991;32:2121.
 There are a number of catalysts and conditions which will effect this
 transformation such as: $\text{Pd}(\text{OAc})_2$, NaHCO_3 , DMF, tetrabutylammonium chloride
 at room temperature to reflux; $\text{Pd}(\text{OAc})_2$, triethylamine, acetonitrile at room
 temperature to reflux. The desired ketone D can also be prepared by acid
 catalyzed reaction of allylic alcohol I, or an enone, with an appropriate
 heterocycles as described in *Heterocycles*, 1987;25:399.


Scheme 6



An alternate preparation of ketone G can be accomplished by reaction of
 aldehyde A with an appropriate nucleophile such as an organo lithium or Grignard
 reagents from -78°C to room temperature in inert solvents to afford the alcohol.
 The desired ketone G can be formed by oxidation of the resulting alcohol with
 Jones reagent or other oxidizing conditions such as the Swern oxidation which are
 well-known to one skilled in the art.

-42-

Scheme 7




Acid E can be converted to desired ketone K utilizing a four step route indicated in Scheme 7. The appropriate acid E can be converted into the
 5 corresponding β -ketoester J utilizing conditions described in *Tetrahedron Letters*, 1992:1945 and *Angew. Chem. Int. Ed.*, 1979:72. The resulting β -ketoester L can be treated with a base, such as NaH or lithium diisopropylamide (LDA), then with an appropriate alkylating agent, R'X at 0°C to room temperature to give the alkylated β -ketoester. The β -ketoester can be decarboxylated with acid in
 10 tetrahydrofuran at reflux. Alternatively the β -ketoester can be hydrolyzed with base to the acid, then heated with or without acid to give the desired ketone K.

The ketones C, D, G, and K and those prepared from Scheme 7 are cyclized to the dihydropyrone ring following the routes shown in Schemes 8 and 9. Scheme 9 also describes methods for obtaining optically pure forms of the
 15 dihydropyrone ring which is a recognized and important aspect of the current invention. The same methods can be used to produce racemic compounds if the resolution step is removed.

-43-

Scheme 8




The dihydropyrone ring system can be prepared as described in *Can. J. Chem.*, 1974;52:2157-2164 and *Synthetic Communications*, 1988;18(9):949-963.

- 5 Reaction of the dianion of acetoacetate (L) with the appropriate ketone (C, D, G, or K) in inert solvents such as tetrahydrofuran at 0°C to room temperature gave aldol intermediate M. H₂O can be added directly to the reaction to effect closure. Alternatively the reaction can be worked up by addition of acetic acid or ammonium chloride and the aldol product isolated and characterized or taken on
- 10 crude. The aldol product M can be closed by treating with dilute sodium hydroxide with or without tetrahydrofuran present. The tetrahydrofuran may be necessary to assist in solubilizing the aldol intermediate M. Protecting groups on intermediate M can be removed before closure to assist in solubilizing the intermediate in base.

- 15 Other routes are available to form the dihydropyrone ring such as those described in: *J. C. S. Chem. Comm.*, 1979:578; *J. Am. Chem. Soc.*, 1984;106:4294; and *J. Org. Chem.*, 1975;40:1610.


-44-

Scheme 9



2) Separate (Chiralcel OD or Enzymatic Resolution)

3) Base



2) Separate by recrystallization or chromatography

3) Base

The dihydropyrone ring system can also be prepared optically enriched as shown in Scheme 9. The appropriate ketone (C, D, G, or K) can be reacted with the appropriate Reformatsky reagent or the equivalent lanthanide species (5 *Tetrahedron*, 1981;37(Supp. 1):175; *J. Org. Chem.*, 1984;49:3904) in inert solvents such as tetrahydrofuran at -78°C to reflux to afford the aldol intermediate. The two antipodes can be separated by a chiral HPLC column such as Chiralcel OD (90% hexane:0.1% TFA:10% isopropyl alcohol) or selective enzymatic hydrolysis with enzymes such as *Candida antarctica* "B" lipase in (10 phosphate buffer with co-solvents such as isopropyl alcohol at room temperature. The resolved ester can be hydrolyzed to the acid O* using base under standard conditions.

Intermediate R* can also be prepared in a chiral form by reaction of (15 ketone C, D, G or K with a chiral ester Q to give an intermediate which is a mixture of diastereomers (*J. Org. Chem.*, 1982;47:91; *Tetrahedron*, 1980;36:227).

-45-


The aldol mixture can be separated by recrystallization or by chromatography to give each enantiomer. Hydrolysis of the ester using base affords acid R*.

Intermediate O can be prepared in racemic form and then resolved by classical means such as co-crystallization with a chiral amine such as 1-(1-naphthyl)ethylamine in solvents such as H₂O. Analysis of the chiral salt by x-ray may allow the determination of the absolute stereochemistry of intermediate O*.

The intermediates O* or R* can be converted to P* by activation of the acid and treatment with the magnesium salt of a half acid ester. Then P*, like M, may be converted to N or N* by base.

Schemes 10 to 14 describe the synthesis of a number of the tosyl reagents (X, BB, HH) which are used to introduce the 3-thiol moiety to the dihydropyrones of the invention.

Scheme 10



Phenol S can be reacted with electrophiles, such as HC(OEt)₃ in inert solvents with a Lewis acid, to give the para-substituted phenol. The phenol can be protected with groups such as methoxyethoxymethyl or tertbutyldimethylsilyl which are known to one skilled in the art. The derivatized phenol can be converted to the corresponding thiol using a variety of methods (*Tetrahedron Lett.*,

-46-


1996:4523; *Chem. Lett.*, 1985:1307, and *Tetrahedron Lett.*, 1993:393). The Newman-Kwart rearrangement is also useful for the conversion of phenol to thiophenol as described in *J. Org. Chem.*, 1966:3980; *Synth.*, 1975:43; and *J. Chem. Eng. Data*, 1975;20:443. Phenol U can be treated with bases such as

5 sodium hydride and dimethylthiocarbamoyl chloride in solvents such as DMF or tetrahydrofuran at 0°C to reflux to give V. Vigorous heating of V at temperatures in the 200°C to 330°C range affords intermediate X. The free thiol can be prepared by reduction of derivatives such as W with diisobutylaluminum hydride (DIBAL-H) or sodium borohydride in inert solvents such as toluene or

10 tetrahydrofuran from -78°C to room temperature or by hydrolysis in base. The desired thiosylate X can be prepared by reaction of the thiol with tosyl bromide and base such as triethylamine or pyridine in inert solvents such as carbon tetrachloride at 0°C to room temperature.

-47-

Scheme 11




Another route into desired thiotosyl reagents is indicated in Scheme 11.

Aniline Y can be converted to the corresponding phenol using a variety of
 5 conditions (*J. Org. Chem.*, 1951;16:586; *Org. Synth.*, 1955; Coll. Vol. 3:130). The
 phenol can be treated with sodium thiocyanate, sodium bromide, and bromine in
 MeOH at 0°C to 50°C to incorporate the thiocyanate to give Z (*Synth.*, 1992:656).
 The phenol can be modified or protected using conditions which are understood
 by one skilled in the art. The thiocyanate can be converted to the thiol (AA) by
 10 treatment with dithiothreitol (DTT) in phosphate buffer in ethanol at room
 temperature to reflux or by treatment with lithium aluminum hydride (LAH) in

-48-

inert solvents such as tetrahydrofuran at 0°C to room temperature. The desired thiotosylate BB can be prepared by reaction of the thiol with tosyl bromide and base as described in Scheme 10.

Scheme 12




5

10

Thiol AA can be converted to the disulfide (CC) by treatment with iodine and triethylamine in EtOAc. The disulfide can act as a protecting group for the sulfur. Various reactions can then be carried out on CC and the disulfide converted back to the thiol by treatment with dithiothreitol (DTT) using similar conditions as described in Scheme 11. Reaction of the free thiophenol with tosyl bromide as previously described affords the desired tosyl reagent BB.

-49-

Scheme 13




Thiocyanate Z can be converted to the thiol and then to thiotosylate DD using conditions which have been described in previous schemes.

- 5 Thiotosylate DD can be derivatized or a protecting group attached using conditions known to one skilled in the art to give BB.

-50-

Scheme 14



The nitro aromatic EE (*J. Org. Chem.*, 1951:586) can be reduced to the corresponding aniline by hydrogenation over Raney Nickel at room temperature.


5 The thiocyanate can be introduced para to the amine in a similar manner as described in Scheme 11. Aniline FF can be converted to GG by modification of the amine, conversion to thiol, and tosylation as previously described.

Alternatively aniline II can be protected, converted to the thiol, and reacted with tosyl bromide to give JJ. Aniline JJ can be deprotected and then modified to give HH. The reactions in this scheme have been previously described or are
 10 known to one skilled in the art.

Scheme 15 shows the convergent preparation of the desired compounds of the invention. The dihydropyrones (N) described in Schemes 8 and 9 are reacted with the tosyl reagents (X, BB, HH) described in Schemes 10 to 14 to produce the
 15 target compounds.

-51-


Scheme 15



The desired dihydropyrone KK can be prepared by reaction with the appropriate dihydropyrone N or N*, thiosylate (X, BB, HH), and potassium carbonate in dimethylformamide (DMF) at room temperature. Other bases and solvents will also effect this reaction such as triethylamine in ethanol or sodium hydride in tetrahydrofuran at 0°C to reflux. Alternatively, the intermediate N can be brominated with NBS in tert-butanol and then displaced with the thiols such as AA and GG.

Scheme 16 describes alternative methods for effecting the separation of enantiomers into optically pure forms after formation of the dihydropyrone ring.

Scheme 16




It has been shown that tetrahydropyrone can be resolved as described in the ICI patent WO 93/06235. The 4-hydroxydihydropyrone can be converted to an ester and the ester hydrolyzed with an enzyme to afford chiral material. This or other similar enzymatic processes may be applicable to the dihydropyrone as indicated in Scheme 16. The 4-hydroxydihydropyrone may also be resolved by

-52-

classical means by conversion into a salt with chiral amines such as 1-(1-naphthyl)ethylamine. The salt may be recrystallized and then freed to afford the desired chiral material MM*.

Scheme 17



5

Scheme 17 shows the general preparation of a number of substituted heterocycles. When X is a halogen, a metal halogen exchange can be preformed to prepare the appropriate anion which can be trapped with an appropriate electrophile (E). This can be used to prepare heterocyclic aldehydes by trapping with DMF, thiols by trapping with sulfur, and alkyl groups by trapping with alkyl ketones or alkyl halides. In certain cases an appropriate base will deprotonate a heterocyclic hydrogen to prepare an anion directly. The following references exemplify Scheme 17: *J. Chem. Soc., Perkin Trans. 1*, 1997:3465; *J. Chem. Soc., Perkin Trans. 1*, 1995:2913; *J. Org. Chem.*, 1971;36(18):2690.

10

Preparation of Aldehydes

Example A-1

5-Methyl-thiazole-4-carbaldehyde

A solution of 2.48 g (14.4 mmol) of 5-methyl-thiazole-4-carboxylic acid ethyl ester (*J. Chem. Soc., Perkin Trans. 1*, 1982:159-164) in dichloromethane (CH_2Cl_2) (50 mL) was cooled to -78°C under nitrogen. DIBAL (Diisobutylaluminum hydride) (1.0 M in CH_2Cl_2 , 15 mmol) was added dropwise, and the solution was stirred at low temperature for 45 minutes. Another 10 mmol of DIBAL was then added dropwise, and the mixture was stirred for another 45 minutes. A solution of methanol (MeOH):acetic acid (10 mL:5 mL) was added slowly, followed by H_2O . The organic layer was separated, washed with brine, and dried (MgSO_4). Concentration gave a residue which was chromatographed on silica gel, eluting with 2:1 hexane:ethyl acetate (EtOAc), to give the title compound.

^1H NMR (CDCl_3): δ 2.76 (s, 3 H), 8.58 (s, 1 H), 10.14 (s, 1 H).

Example A-2

2-Methyl-thiazole-4-carbaldehyde

A mixture of 5.52 g (30 mmol) of 4-chloromethyl-2-methylthiazole hydrochloride and 0.5 M NaOH (180 mL) was refluxed for 8 hours and then stirred for 18 hours at room temperature. EtOAc and H_2O were added. The organic phase was separated, washed with brine, and dried (MgSO_4). Concentration gave 4-(hydroxymethyl)-2-methylthiazole.

^1H NMR (CDCl_3): δ 2.63 (s, 3 H), 4.66 (s, 2 H), 6.96 (s, 1 H).

The crude product (2.2 g, 17 mmol) was oxidized with MnO_2 (20 g) in 75 mL of CHCl_3 for 16 hours. The suspension was filtered, and the filtrate was concentrated. The residue was chromatographed on silica gel, eluting with 2:1 hexane:EtOAc, to give the title compound.

^1H NMR (CDCl_3): δ 2.71 (s, 3 H), 7.98 (s, 1 H), 9.89 (s, 1 H).

-54-

Example A-3

2-iso-Propyl-thiazole-4-carbaldehyde

A solution of 2.7 g (13 mmol) of 2-isopropyl-thiazole-4-carboxylic acid ethyl ester (*J. Med. Chem.*, 1998:602-617) in CH₂Cl₂ (50 mL) was cooled to
5 -78°C under nitrogen and treated dropwise with DIBAL (15 mL of 1.0 M in CH₂Cl₂; 15 mmol). The mixture was stirred at low temperature for 45 minutes and then treated with another 15 mL of DIBAL. The solution was stirred for 1 hour at -78°C. Five percent citric acid was added, and the mixture was extracted with EtOAc (3 × 100 mL). The combined extracts were washed with brine, dried
10 (MgSO₄), and concentrated. Chromatography of the residue over silica gel, eluting with 3:1 hexane:EtOAc, gave the title compound.
¹H NMR (CDCl₃): δ 1.37 (d, 6 H), 3.30-3.35 (m, 1 H), 8.01 (s, 1 H), 9.93 (s, 1 H).

Example A-4

4-iso-Propyl-thiazole-5-carbaldehyde

A solution of 5.9 g (30 mmol) of 4-isopropyl-thiazole-5-carboxylic acid ethyl ester (*J. Chem. Soc. Perkin Trans 1*, 1982:159-164) in toluene (100 mL) was cooled in an ice bath under nitrogen and treated dropwise with DIBAL (150 mL of 1.0 M; 150 mmol). The mixture was stirred at low temperature for
20 45 minutes and then allowed to warm to room temperature overnight. H₂O was added cautiously, and the mixture was extracted with EtOAc (3 × 200 mL). The combined extracts were washed with brine, dried (MgSO₄), and concentrated. The crude 5-(hydroxymethyl)-4-isopropyl-thiazole thus obtained was used as is in the next step.
25 ¹H NMR (CDCl₃): δ 1.30 (d, 6 H), 3.13-3.20 (m, 1 H), 4.85 (s, 2 H), 8.67 (s, 1 H).

A mixture of 4.65 g (30 mmol) of crude 5-(hydroxymethyl)-4-isopropyl-thiazole in 200 mL of chloroform (CHCl₃) was treated with 45 g of MnO₂ and stirred at room temperature for 2.5 hours. The suspension was filtered, and the

-55-

filtrate was concentrated. The residue was chromatographed on silica gel, eluting with 3:1 hexane:EtOAc, to give the title compound.

^1H NMR (CDCl_3): δ 1.40 (d, 6 H), 3.62-3.69 (m, 1 H), 8.97 (s, 1 H), 10.17 (s, 1 H).

5

Example A-5

5-iso-Propyl-thiazole-4-carbaldehyde

A solution of 6.8 g (37 mmol) of 5-isopropyl-thiazole-4-carboxylic acid methyl ester (*J. Chem. Soc., Perkin Trans. 1*, 1982:159-164) in CH_2Cl_2 (120 mL) was cooled to -78°C under nitrogen and treated dropwise with DIBAL (37 mL of 1.0 M in CH_2Cl_2 ; 37 mmol). The mixture was stirred at low temperature for 45 minutes and then treated with another 20 mL of DIBAL. The solution was stirred for 1 hour at -78°C . H_2O was added, and the mixture was extracted with EtOAc (3×150 mL). The combined extracts were washed with brine, dried (MgSO_4), and concentrated. Chromatography of the residue over silica gel, eluting with 2:1 hexane:EtOAc, gave the title compound.

^1H NMR (CDCl_3): δ 1.34 (d, 6 H), 4.09-4.16 (m, 1 H), 8.63 (s, 1 H), 10.20 (s, 1 H).

10
15

Example A-6

4-iso-Propyl-thiazole-2-carbaldehyde

A solution of 2.1 g (11 mmol) of 4-isopropyl-thiazole-2-carboxylic acid ethyl ester (*J. Med. Chem.*, 1998:602-617) in CH_2Cl_2 (50 mL) was cooled to -78°C under nitrogen and treated dropwise with DIBAL (12 mL of 1.0 M in CH_2Cl_2 ; 12 mmol). The mixture was stirred at low temperature for 1 hour and then treated with another 7 mL of DIBAL. The solution was stirred for 30 minutes at -78°C . H_2O was added, and the mixture was extracted with EtOAc (3×100 mL). The combined extracts were washed with brine, dried (MgSO_4), and concentrated. Chromatography of the residue over silica gel, eluting with 3:1 hexane:EtOAc, gave the title compound.

20
25

-56-

^1H NMR (CDCl_3): δ 1.35 (d, 6 H), 3.16-3.23 (m, 1 H), 7.33 (s, 1 H), 9.96 (s, 1 H).

The remaining aldehydes were either commercially available or known in the chemical literature, as summarized in Table A below.

TABLE A. Preparation of Aldehydes








Example	Aryl	Method of Preparation	^1H NMR Data
A-7		Commercially available	
A-8		<i>Synthesis</i> , 1987:998-1001	(CDCl_3): δ 8.22 (s, 1 H), 8.88 (s, 1 H), 10.10 (s, 1 H).
A-9		<i>Synthesis</i> , 1987:998-1001	(CDCl_3): δ 8.51 (s, 1 H), 9.10 (s, 1 H), 10.10 (s, 1 H).
A-10		<i>Acta. Chemica Scandinavica</i> , 1966;20:2649-2657	(CDCl_3): δ 2.49 (s, 3 H), 7.29 (s, 1 H), 9.86 (s, 1 H).
A-11		<i>Acta. Chemica Scandinavica</i> , 1966;20:2649-2657	(CDCl_3): δ 2.51 (s, 3 H), 7.71 (s, 1 H), 9.80 (s, 1 H).
A-12		<i>J. Amer. Chem. Soc.</i> , 1982:4934-4943	(CDCl_3): δ 2.76 (s, 3 H), 8.94 (s, 1 H), 10.11 (s, 1 H).
A-13		<i>Bull. Chim. France</i> , 1967:2235-2238	(CDCl_3): δ 2.19 (s, 3 H), 7.74 (s, 1 H), 9.77 (s, 1 H), 11.69 (br s, 1 H).

TABLE A. Preparation of Aldehydes (cont'd)










Example	Aryl	Method of Preparation	¹ H NMR Data
A-14		Commercially available	
A-15		Commercially available	
A-16		Commercially available	
A-17		Commercially available	
A-18		<i>J Org. Chem.</i> , 1990;55(26):6317-6328	(CDCl ₃): δ 6.69 (m, 1 H), 6.85 (m, 1 H), 7.46 (m, 1 H), 9.2 (br s, 1 H), 9.82 (s, 1 H).
A-19		Commercially available	
A-20		Commercially available	
A-21		Commercially available	
A-22		Commercially available	

TABLE A. Preparation of Aldehydes (cont'd)














Example	Aryl	Method of Preparation	¹ H NMR Data
A-23		Commercially available	
A-24		Commercially available	
A-25		Commercially available	
A-26		Commercially available	
A-27		Commercially available	
A-28		Commercially available	
A-29		Commercially available	
A-30		<i>Chem Pharm. Bull.</i> , 1986;34(10):4116-4125	(CDCl ₃): δ 6.67 (m, 1 H), 7.25 (m, 1 H), 7.45 (d, 1 H), 7.74 (d, 1 H), 8.15 (s, 1 H), 8.70 (br s, 1 H), 10.00 (s, 1 H).
A-31		<i>J.O.C.</i> , 1987;53:107-109	

TABLE A. Preparation of Aldehydes (cont'd)

Example	Aryl	Method of Preparation	¹ H NMR Data
A-32		Commercially available	
A-33		Commercially available	
A-34		Commercially available	
A-35		Commercially available	

Methods for the Preparation of Enones: General Method 1. Preparation of Enones Via Triphenylarsonium Salts

A mixture of the appropriate aldehyde from Examples A-1 to A-29 (1.0 equiv.), the crude arsonate salt (usually (3-methyl-2-oxobutyl)triphenylarsonium bromide, prepared in *Synthesis*, 1988:975-977; 1.1 equiv.), potassium carbonate (1.1 equiv.), and 1% H₂O in CH₃CN was stirred for 1 to 18 hours at room temperature. The solids were filtered, and the filtrate was chromatographed on silica gel to remove the arsenate by-products.

Example B-1

10 4-Methyl-1-(4-methyl-thiazol-5-yl)-pent-1-en-3-one

The title compound was prepared as described in General Method 1 using 1.4 g (11.0 mmol) of 4-methyl-5-thiazolecarbaldehyde (Example A-12), 6.2 g (13.1 mmol) of (3-methyl-2-oxobutyl)triphenylarsonium bromide, 1.8 g (13.1 mmol) of potassium carbonate, 75 mL of acetonitrile, and 0.75 mL of H₂O.

15 Chromatography of the residue, eluting with 3:1 hexane:EtOAc gave the title compound.

-60-

¹H NMR (CDCl₃): δ 1.13 (d, 6 H), 2.54 (s, 3 H), 2.75-2.82 (m, 1 H), 6.50 (d, 1 H), 7.73 (d, 1 H), 8.66 (s, 1 H).

General Method 2. Preparation of Enones Via Condensation with Ba(OH)₂

The desired enones were prepared by reaction of the appropriate methylketones and aldehydes by the methods described in *Synthesis*, 1983:502-504; *An. Quim, Ser. C*, 1981;77(2):222-224; *Org. Synth Coll.*, 1941;1:78; and *Pol. J. Chem.*, 1982;56(10-112):1435. To a reaction flask was added aldehyde (1-2 equiv.), ketone (1-2 equiv.), 95% EtOH, and anhydrous or hydrated Ba(OH)₂ (23 mg/mmol). The reaction was stirred at room temperature or heated at reflux for up to 2 days. The reaction was cooled to room temperature. The EtOH was evaporated, and the crude reaction was partitioned between EtOAc and 1N HCl. The aqueous layer was extracted with EtOAc. The combined organic extracts were dried (MgSO₄) and concentrated. The crude product was purified by flash chromatography or recrystallization.

Example B-2

1-Furan-2-yl-4-methyl-pent-1-en-3-one


The titled compound was prepared according to General Method 2 by reacting 2-furaldehyde (117 mmol), 2-methyl-butan-3-one (117 mmol), anhydrous Ba(OH)₂ (2.2 g) and EtOH (200 mL) to afford the desire compound.

¹H NMR (CDCl₃): δ 1.18 (d, 6 H), 2.83 (m, 1 H), 6.46 (m, 1 H), 6.65 (m, 1 H), 6.73 (d, 1 H), 7.38 (d, 1 H), 7.50 (m, 1 H).

The following enones were prepared using either General Method 1 (the arsonate salt) or General Method 2 (the Ba(OH)₂ method) from the corresponding aldehydes in Examples A-1 to A-35:

-61-











TABLE B. Preparation of Enones



Example	R ₇	General Method	NMR Data (δ for vinyl protons. Solvent: CDCl ₃) or Mass Spec
B-3		Gen. Method 1	7.05, 7.64
B-4		Gen. Method 1	7.14, 7.53
B-5		Gen. Method 1	6.58, 7.72
B-6		Gen. Method 1	6.90, 7.52
B-7		Gen. Method 1	7.01, 7.57
B-8		Gen. Method 1	7.06, 7.43
B-9		Gen. Method 1	7.12, 7.54
B-10		Gen. Method 1	7.11, 7.49
B-11		Gen. Method 1	6.52, 7.78
B-12		Gen. Method 1	7.24, 7.64


-62-

TABLE B. Preparation of Enones (cont'd)

			
Example	R ₇	General Method	NMR Data (δ for vinyl protons. Solvent: CDCl ₃) or Mass Spec
B-13		Gen. Method 1	6.95, 7.60
B-14		Gen. Method 1	6.93, 7.41
B-15		Gen. Method 1	7.33, 7.60
B-16		Gen. Method 1	6.96, 7.51
B-17		Gen. Method 1	MS (APCI): 243 (M+H)
B-18		Gen. Method 1	MS (APCI): 243 (M+H)
B-19		Gen. Method 1	MS (APCI): 243 (M+H)
B-20		Gen. Method 1	MS (APCI): 261 (M+H)
B-21		Gen. Method 1	MS (APCI): 261 (M+H)

-63-

TABLE B. Preparation of Enones (cont'd)



Example	R ₇	General Method	NMR Data (δ for vinyl protons. Solvent: CDCl ₃) or Mass Spec
B-22		Gen. Method 1	MS (APCI): 200 (M+H)
B-23		Gen. Method 2	6.74, 7.86
B-24		Gen. Method 2	6.90, 7.73
B-25		Gen. Method 2	6.74, 7.57
B-26		Gen. Method 2	6.80, 7.56
B-27		Gen. Method 2	
B-28		Gen. Method 1	MS (APCI): 164 (M+H)
B-29		Gen. Method 1	MS (APCI): 165 (M+H)
B-30		Gen. Method 2	6.59, 7.56
B-31		Gen. Method 1	6.82, 7.78

-64-

TABLE B. Preparation of Enones (cont'd)

Example	R ₇	General Method	NMR Data (δ for vinyl protons. Solvent: CDCl ₃) or Mass Spec
B-32		Gen. Method 1	6.78, 6.99
B-33		Gen. Method 2	7.11, 7.70
B-34		Gen. Method 2	6.79, 7.49
B-35		Gen. Method 2	6.84, 7.66
B-36		Gen. Method 2	6.73, 7.43

Names corresponding to the enones in Table B above are:

B-3: 4-Methyl-1-thiazol-2-yl-pent-1-en-3-one;

B-4: 4-Methyl-1-thiazol-4-yl-pent-1-en-3-one;

B-5: 4-Methyl-1-thiazol-5-yl-pent-1-en-3-one;

5 B-6: 4-Methyl-1-(5-methyl-thiazol-2-yl)-pent-1-en-3-one;

B-7: 4-Methyl-1-(4-methyl-thiazol-2-yl)-pent-1-en-3-one;

B-8: 4-Methyl-1-(2-methyl-thiazol-4-yl)-pent-1-en-3-one;

B-9: 4-Methyl-1-(5-methyl-thiazol-4-yl)-pent-1-en-3-one;

B-10: 1-(2-Isopropyl-thiazol-4-yl)-4-methyl-pent-1-en-3-one;

10 B-11: 1-(4-Isopropyl-thiazol-5-yl)-4-methyl-pent-1-en-3-one;

B-12: 1-(5-Isopropyl-thiazol-4-yl)-4-methyl-pent-1-en-3-one;

-65-

- B-13: 1-(4-Isopropyl-thiazol-2-yl)-4-methyl-pent-1-en-3-one;
 B-14: N-[4-(4-Methyl-3-oxo-pent-1-enyl)-thiazol-2-yl]-acetamide;
 B-15: 4-Methyl-1-pyridin-2-yl-pent-1-en-3-one;
 B-16: 4-Methyl-1-pyridin-4-yl-pent-1-en-3-one;
 5 B-17: 4-Methyl-1-(2-trifluoromethyl-phenyl)-pent-1-en-3-one;
 B-18: 4-Methyl-1-(3-trifluoromethyl-phenyl)-pent-1-en-3-one;
 B-19: 4-Methyl-1-(4-trifluoromethyl-phenyl)-pent-1-en-3-one;
 B-20: 4-Methyl-1-(3-fluoro-5-trifluoromethyl-phenyl)-pent-1-en-3-one;
 B-21: 4-Methyl-1-(4-fluoro-3-trifluoromethyl-phenyl)-pent-1-en-3-one;
 10 B-22: 4-(4-Methyl-3-oxo-pent-1-enyl)-benzonitrile;
 B-23: 4-Methyl-1-(3-fluoro-2-methyl-phenyl)-pent-1-en-3-one;
 B-24: 4-Methyl-1-(2-fluoro-phenyl)-pent-1-en-3-one;
 B-25: 4-Methyl-1-(4-fluoro-phenyl)-pent-1-en-3-one;
 B-26: 4-Methyl-1-(3-fluoro-phenyl)-pent-1-en-3-one;
 15 B-27: 4-Methyl-1-(3,4-difluoro-phenyl)-pent-1-en-3-one;
 B-28: 4-Methyl-1-(1H-pyrrol-3-yl)-pent-1-en-3-one;
 B-29: 1-(1H-Imidazol-4-yl)-4-methyl-pent-1-en-3-one;
 B-30: 4-Methyl-1-thiophen-3-yl-pent-1-en-3-one;
 B-31: 1-(1H-Indol-5-yl)-4-methyl-pent-1-en-3-one;
 20 B-32: 4-Methyl-1-(2-methyl-thiophen-3-yl)-pent-1-en-3-one;
 B-33: 4-Methyl-1-(2,6-difluoro-phenyl)-pent-1-en-3-one;
 B-34: 4-Methyl-1-(3,5-difluoro-phenyl)-pent-1-en-3-one;
 B-35: 4-Methyl-1-(2,4-difluoro-phenyl)-pent-1-en-3-one; and
 B-36: 4-Methyl-1-(3,4,5-trifluoro-phenyl)-pent-1-en-3-one.

25 **Methods for the Preparation of Requisite Ketones: General Method 3.**

Preparation of Ketones by Hydrogenation of Enones

The substituted arylpropiophenones were prepared by hydrogenation of the corresponding enones at room temperature at 50 psi in tetrahydrofuran (THF) utilizing a number of catalysts such as: 5% palladium (Pd) on barium sulfate
 30 (BaSO₄) or (Ph₃P)₃RhCl or 10% Pd on carbon or 5% Pd on CaCO₃. The catalyst

-66-

was filtered off and the resulting ketone recrystallized or purified by flash chromatography.

Example C-1

4-Methyl-1-pyridin-4-yl-pentan-3-one

5 The title compound was prepared according to General Method 3 using 4-methyl-1-pyridin-4-yl-pent-1-en-3-one (Example B-16; 1.81 g), (Ph₃P)₃RhCl (0.5 g), and THF (100 mL) at 50 psi. The title compound was flash chromatographed using 98:2 CH₂Cl₂:MeOH as eluent. ¹H NMR (CDCl₃): δ 1.15 (d, 6 H), 2.5-2.7 (m, 1 H), 2.7-2.8 (m, 2 H), 2.85-2.95 (m, 2 H), 7.1 (d, 2 H), 8.5 (d, 2 H).

10

Example C-2

4-Methyl-1-(4-methyl-thiazol-5-yl)-pentan-3-one

 The title compound was prepared as described in General Method 3 using 1.73 g (8.86 mmol) of 4-methyl-1-(4-methyl-thiazol-5-yl)-pent-1-en-3-one (prepared in Example B-1) and 0.4 g of 5% Pd/BaSO₄ in THF:MeOH (25 mL:25 mL). The crude product was chromatographed on silica gel, eluting with 2:1 hexane:EtOAc, to give the title compound. ¹H NMR (CDCl₃): δ 1.03 (d, 6 H), 2.35 (s, 3 H), 2.49-2.56 (m, 1 H), 2.72 (t, 2 H), 2.98 (t, 2 H), 8.49 (s, 1 H).

15

Synthesis of Appropriate Halogenated Heterocycles Necessary for Ketone Synthesis (General Methods 4a and 4b)

20

Example Starting Material - 1 (SM)

(5-Bromo-thiophen-2-yl)-methanol

 The title compound was prepared by reduction of 5-bromo-2-thiophenecarbaldehyde (137 mmol) with sodium borohydride (137 mmol) in MeOH (500 mL). The reaction was stirred for 2 hours at 0°C and 2 hours at room temperature. The MeOH was evaporated, and saturated ammonium chloride was added followed by 2N HCl. The aqueous layers were extracted with EtOAc, dried (MgSO₄), and concentrated. Flash chromatography over silica gel using 100%

25

-67-

CH₂Cl₂ as eluent afforded the title compound. ¹H NMR (CDCl₃): δ 4.74 (d, 2 H), 6.75 (m, 1 H), 6.91 (d, 1 H).

Example Starting Material - 2

(4-Bromo-thiophen-2-yl)-methanol

5 The title compound was prepared in a similar manner to Example SM-1 by reduction of 4-bromo-2-thiophenecarboxaldehyde (122 mmol) with sodium borohydride (12 mmol) in MeOH (500 mL).

¹H NMR (CDCl₃): δ 4.80 (d, 2 H), 6.93 (d, 1 H), 7.18 (d, 1 H).

Example Starting Material - 3

10 **(3-Bromo-thiophen-2-yl)-methanol**

 The title compound was prepared by reduction of 3-bromothiophene-2-carboxylic acid methyl ester (45 mmol) with lithium aluminum hydride (45 mmol) in THF (150 mL) at 0°C for 1 hour and then overnight at room temperature. The reaction was worked up by addition of 1 mL of H₂O, 1 mL of 15
15 15% NaOH, and 3 mL of H₂O followed by filtration through celite. Concentration of the filtrate gave the title compound. ¹H NMR (CDCl₃): δ 4.80 (s, 2 H), 6.96 (d, 1 H), 7.26 (d, 1 H).

Example Starting Material - 4

1-Trityl-4-iodopyrazole

20 The title compound was prepared by combining 4-iodopyrazole (10 g, 52 mmol), triphenylmethyl chloride (14.4 g, 51.6 mmol), triethylamine (NEt₃) (7.2 mL, 52 mmol), and DMF (80 mL). After stirring overnight, the mixture was poured onto ice water. The precipitated solid was collected and recrystallized to give the title compound as a solid, mp 193-194°C.

25 ¹H NMR (CDCl₃): δ 7.11 (m, 6 H), 7.32 (m, 9 H), 7.41 (s, 1 H), 7.67 (s, 1 H).

-68-

Example Starting Material - 5 (SM)

(4-Bromo-thiophen-3-yl)-methanol

The title compound was prepared in a similar manner to Example SM-1 by reduction of 4-bromo-3-thiophenecarbaldehyde (*Bull. Soc. Chim.*

5 *France*, 1967;11:4115) with sodium borohydride in MeOH. ¹H NMR (CDCl₃): δ 2.05 (s, 2 H), 7.27-7.34 (m, 2 H).

Example Starting Material - 6 (SM)

(2-Bromo-thiophen-3-yl)-methanol

10 A solution of 10.0 g (39.0 mmol) of 2-bromo-3-bromomethyl-thiophene (*J Chem. Soc Perkin Trans. II*, 1983:813) in 130 mL of acetone was treated with a solution of 11.4 g (67.1 mmol) of AgNO₃ in 110 mL of H₂O. The mixture was stirred at room temperature for 1 hour and filtered. The solid was washed with Et₂O and acetone; the combined filtrate and washings were concentrated. The residue was extracted with CH₂Cl₂, and the organic layer was dried (MgSO₄) and
15 concentrated. The crude product was chromatographed over silica gel, eluting with hexane:EtOAc:CH₂Cl₂ (80:5:15 to 60:30:10) to give the title compound.
¹H NMR (CDCl₃) δ 1.67 (t, 1 H), 4.63 (d, 2 H), 7.03 (d, 1 H), 7.26 (d, 1 H).

Example Starting Material - 7 (SM)

3-Bromo-4-ethyl-fluorobenzene

20 2-Bromo-4-fluoroacetophenone (10.0 g, 46.1 mmol) in THF (120 mL) was treated with BF₃·OEt₂ (22.9 g, 20.4 mL, 161 mmol) and followed by NaBH₃CN (7.24 g, 155 mmol) added portionwise. The resulting mixture was heated to reflux under N₂ overnight and then cooled to room temperature. Diethyl ether (Et₂O) was added; the organic layer was washed with saturated NaHCO₃/H₂O and brine,
25 dried (MgSO₄), filtered, and concentrated. The resulting residue was subjected to flash silica gel chromatography, eluting with 4:1 (hexanes:EtOAc) to afford the intermediate alcohol. ¹H NMR (CDCl₃): δ 1.46 (d, 3 H), 1.98 (bs, 1 H), 5.21 (q, 1 H), 7.03-7.10 (m, 1 H), 7.26 (dd, 1 H), 7.58 (dd, 1 H).

-69-

The alcohol isolated above (4.0 g, 18 mmol) was dissolved in hexanes (20 mL) and treated with trimethylsilyl chloride (11.9 g, 13.9 mL, 109 mmol), NaI (16.4 g, 109 mmol), and CH₃CN (5.7 mL, 109 mmol). The resulting slurry was stirred at room temperature under N₂ overnight. Et₂O was added followed by

5 H₂O. The phases were separated, and the aqueous phase was extracted again with Et₂O; the combined organic phases were washed with sodium bisulfite (NaHSO₃)/H₂O and brine, dried (MgSO₄), filtered, and concentrated. The resulting residue was subjected to flash silica gel chromatography, eluting with hexanes to afford the title compound. ¹H NMR (CDCl₃): δ 1.20 (t, 3 H), 2.72 (q,

10 2 H), 6.93-6.99 (m, 1 H), 7.18 (dd, 1 H), 7.27 (dd, 1 H).

Example Starting Material - 8 (SM)

(2-Bromo-5-fluoro-phenyl)-methanol

2-Bromo-5-fluorobenzylbromide (2.0 g, 7.46 mmol) was dissolved in dioxane (25 mL) and H₂O (25 mL), and CaCO₃ (3.84 g, 38.39 mmol) was added.












15 The mixture was heated to reflux overnight. The solution was concentrated, and the residue was partitioned between H₂O and CH₂Cl₂. The organic phase was washed with 1N HCl and brine, dried (MgSO₄), filtered, and concentrated to afford the title compound. ¹H NMR (CDCl₃): δ 2.00 (t, 1 H), 4.72 (d, 2 H), 6.85-

20 6.92 (m, 1 H), 7.26 (dd, 1 H), 7.48 (dd, 1 H).

The remaining halides were commercially available, as summarized in Table C below.

-70-

TABLE C. Aryl Halides

Example	Aryl
SM-9	
SM-10	
SM-11	
SM-12	
SM-13	
SM-14	
SM-15	
SM-16	
SM-17	
SM-18	
SM-19	

-71-

Methods for the Preparation of Requisite Ketones: General Method 4a and 4b. Preparation of Ketones via Pd-Catalyzed Coupling

General Method 4a is the method described in *Tetrahedron*, 1979;35:329 and *Tetrahedron Letters*, 1991;32:2121. The appropriate aryl halide (Br or I; 1 equiv.), allylic alcohol (1-2 equiv.), tetrabutyl ammonium chloride (1 equiv.), sodium bicarbonate (2-3 equiv.), DMF (0.1-1 mL per mmol of halide), and palladium acetate (0.01-0.1 equiv.) were added to a reaction vessel. The solution was heated to 40°C to 100°C for 1 to 24 hours. On occasion, pyrrolidine (0.2-1 equiv.) was also added. The reaction was cooled to room temperature and partitioned between H₂O and CH₂Cl₂. The solution was filtered through celite and the aqueous layer extracted two times with CH₂Cl₂. The organic extracts were washed with brine and then dried (MgSO₄). Purification was usually carried out by flash chromatography.

General Method 4b is the method described in *Tetrahedron*, 1979;35:329-340 with slight modifications. The appropriate aryl halide (Br or I; 1 equiv.), allylic alcohol (1-2 equiv.), sodium iodide (0.01-0.050 equiv.), sodium bicarbonate (1-3 equiv.), triphenyl phosphine (0.01-0.050 equiv.), DMF (0.1-1 mL per mmol of halide), and palladium acetate (0.01-0.1 equiv.) were added to a reaction vessel. The solution was heated to 40°C to 100°C for 1 to 24 hours. Pyrrolidine (0.2-1 equiv.) was also added at times. The reaction was cooled to room temperature and partitioned between H₂O and CH₂Cl₂. The solution was filtered through celite and the aqueous layer extracted two times with CH₂Cl₂. The organic extracts were washed with brine and then dried (MgSO₄). Purification was usually carried out by flash chromatography.

Example C-3

4-Methyl-1-pyridin-3-yl-pentan-3-one

The title compound was prepared according to General Method 4a using 3-iodo-pyridine (36.6 mmol), 4-methyl-1-penten-3-ol (54.9 mmol), tetrabutyl ammonium chloride (36.6 mmol), sodium bicarbonate (91.5 mmol), pyrrolidine (~1.5 mL), DMF (15 mL), and palladium acetate (0.51 g). The title compound was flash chromatographed eluting with EtOAc:CH₂Cl₂:hexane 50:25:25.

-72-

¹NMR (CDCl₃): δ 1.15 (d, 6 H), 2.5-2.7 (m, 1 H), 2.7-2.9 (m, 2 H), 2.85-2.95 (m, 2 H), 7.2-7.3 (m, 1 H), 7.5-7.6 (m, 1 H), 8.4-8.5 (m, 2 H).

Example C-4

1-Furan-3-yl-4-methyl-pentan-3-one

5 The title compound was prepared according to General Method 4a using 3-bromofuran (17 mmol), 4-methyl-1-penten-3-ol (25.5 mmol), tetrabutyl ammonium chloride (17 mmol), sodium bicarbonate (42.5 mmol), DMF (15 mL), and palladium acetate (0.9 mmol). The title compound was flash chromatographed eluting with EtOAc:hexane (5:95 to 10:90).

10 ¹NMR (CDCl₃): δ 1.05 (d, 6 H), 2.5-2.7 (m, 1 H), 2.65 (s, 4 H), 6.2 (s, 1 H), 7.25 (m, 1 H), 7.3 (m, 1 H).

Example C-5

4-Methyl-1-thiophen-3-yl-pentan-3-one

15 The title compound was prepared according to General Method 4b using 3-bromothiophene (20 mmol), 4-methyl-1-penten-3-ol (30 mmol), sodium iodide (0.7 mmol), sodium bicarbonate (24 mmol), triphenylphosphine (0.6 mmol), DMF (15 mL), and palladium acetate (0.2 mmol). The title compound was flash chromatographed using EtOAc:hexane (5:95 to 10:90).

20 ¹NMR (CDCl₃): δ 1.05 (d, 6 H), 2.5-2.7 (m, 1 H), 2.7-2.8 (t, 2 H), 2.85-2.95 (t, 2 H), 6.90-9.95 (m, 2 H), 7.2-7.3 (m, 1 H).

 Alternatively, the title compound could be prepared by hydrogenation (General Method 3) of the enone prepared in Example B-30.

General Method 5. Preparation of Silylated Intermediates.

25 The appropriate alcohol (1 equiv.), and imidazole (1.2 equiv.) were added to a reaction vessel followed by CH₂Cl₂ or THF (7-10 mL per mmol of alcohol). t-Butyldimethylsilyl chloride (1.1 equiv.) was added and the reaction stirred at room temperature (3 hours to 4 days). The reaction was filtered, washed with H₂O and brine, dried (MgSO₄), and concentrated. The product was either flashed chromatographed or carried on crude.





-73-

Example C-6

1-[5-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-2-yl]-4-methyl-pentan-3-one








- 5 The title compound was prepared according to General Method 5 using 1-(5-hydroxymethyl-thiophen-2-yl)-4-methyl-pentan-3-one (see Example C-24 below; 73.0 mmol), t-butyldimethylsilyl chloride (80.3 mmol), imidazole (80.3 mmol) and CH₂Cl₂ (300 mL). ¹H NMR (CDCl₃): δ 0.09 (s, 6 H), 0.92 (s, 9 H), 1.09 (d, 6 H), 2.59 (m, 1 H), 2.81 (t, 2 H), 3.06 (t, 2 H), 4.78 (s, 2 H), 6.61 (d, 1 H), 6.69 (d, 1 H).
- 10 The following ketones were prepared as indicated [either from reduction of the corresponding enones from Examples B-1 to B-36 (General Method 3) OR from the corresponding halides from Examples SM-1 to SM-19 (General Method 4a or 4b)]. In some cases, the following ketones were also prepared from silylation of an existing ketone using General Method 5:

TABLE D. Preparation of Ketones

			
Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-7		Gen. Method 3	(CDCl ₃): δ 1.06 (d, 6 H), 2.56-2.63 (m, 1 H), 2.97 (t, 2 H), 3.24 (t, 2 H), 7.13 (d, 1 H), 7.60 (d, 1 H)
C-8		Gen. Method 3	(CDCl ₃): δ 1.04 (d, 6 H), 2.53-2.60 (m, 1 H), 2.87 (m, 2 H), 3.08 (m, 2 H), 6.96 (m, 1 H), 8.70 (m, 1 H)
C-9		Gen. Method 3	(CDCl ₃): δ 1.08 (d, 6 H), 2.55-2.61 (m, 1 H), 2.78 (t, 2 H), 3.15 (t, 2 H), 7.60 (s, 1 H), 8.63 (s, 1 H)


-74-






TABLE D. Preparation of Ketones (cont'd)

			
Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-10		Gen. Method 3	(CDCl ₃): δ 1.03 (d, 6 H), 2.33 (s, 3 H), 2.52-2.59 (m, 1 H), 2.88-2.92 (m, 2 H), 3.09-3.13 (m, 2 H), 7.18 (s, 1 H)
C-11		Gen. Method 3	(CDCl ₃): δ 1.05 (d, 6 H), 2.34 (s, 3 H), 2.55-2.62 (m, 1 H), 2.93 (t, 2 H), 3.18 (t, 2 H), 6.65 (s, 1 H)
C-12		Gen. Method 3	(CDCl ₃): δ 1.02 (d, 6 H), 2.50-2.56 (m, 1 H), 2.62 (s, 3 H), 2.80-2.84 (m, 2 H), 2.91-2.94 (m, 2 H), 6.68 (s, 1 H)
C-13		Gen. Method 3	(CDCl ₃): δ 1.00 (d, 6 H), 2.35 (s, 3 H), 2.50-2.57 (m, 1 H), 2.86 (s, 4 H), 8.45 (s, 1 H)
C-14		Gen. Method 3	(CDCl ₃): δ 1.02 (d, 6 H), 1.32 (d, 6 H), 2.52-2.58 (m, 1 H), 2.83 (t, 2 H), 2.95 (t, 2 H), 3.20-3.27 (m, 1 H), 6.70 (s, 1 H)
C-15		Gen. Method 3	(CDCl ₃): δ 1.07 (d, 6 H), 1.26 (d, 6 H), 2.53-2.60 (m, 1 H), 2.75 (t, 2 H), 3.04 (t, 2 H), 3.07-3.14 (m, 1 H), 8.56 (s, 1 H)

-75-







TABLE D. Preparation of Ketones (cont'd)



Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-16		Gen. Method 3	(CDCl ₃): δ 1.07 (d, 6 H), 1.27 (d, 6 H), 2.56-2.63 (m, 1 H), 2.89-2.97 (m, 4 H), 3.28-3.35 (m, 1 H), 8.53 (s, 1 H)
C-17		Gen. Method 3	(CDCl ₃): δ 1.10 (d, 6 H), 1.26 (d, 6 H), 2.60-2.65 (m, 1 H), 2.98 (t, 2 H), 2.99-3.05 (m, 1 H), 3.24 (t, 2 H), 6.68 (s, 1 H)
C-18		Gen. Method 3	(CDCl ₃): δ 1.04 (d, 6 H), 2.20 (s, 3 H), 2.52-2.59 (m, 1 H), 2.76-2.80 (m, 2 H), 2.86-2.90 (m, 2 H), 6.51 (s, 1 H)
C-19		Gen. Method 3	(CDCl ₃): δ 1.15 (d, 6 H), 2.5-2.7 (m, 1 H), 2.9-3.0 (m, 2 H), 3.0-3.1 (m, 2 H), 7.05-7.15 (m, 1 H), 7.2 (d, 1 H), 7.5-7.6 (dt, 1 H), 8.5 (d, 1 H)
C-20		Gen. Method 3	(CDCl ₃): δ 1.15 (d, 6 H), 2.5-2.7 (m, 1 H), 2.7-2.9 (m, 2 H), 2.85-2.95 (m, 2 H), 5.95 (m, 1 H), 6.25 (m, 1 H), 7.3 (m, 1 H)








-76-

TABLE D. Preparation of Ketones (cont'd)

			
Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-21		Gen. Method 3 (complete reduction of C-20)	MS (APCI): 171 (M+H)
C-22		Gen. Method 4b	(CDCl ₃): δ 1.1 (d, 6 H), 2.5-2.7 (m, 1 H), 2.8-2.9 (t, 2 H), 3.05-3.15 (t, 2 H), 6.75-6.80 (m, 1 H), 6.85-6.95 (m, 1 H), 7.05-7.15 (m, 1 H)
C-23		Gen. Method 4b	(CDCl ₃): δ 1.1 (d, 6 H), 2.5-2.7 (m, 1 H), 2.7-2.8 (m, 2 H), 2.8-2.9 (m, 2 H), 4.75 (s, 2 H), 6.85 (s, 1 H), 6.9 (s, 1 H)
C-24		Gen. Method 4b	(CDCl ₃): δ 1.1 (d, 6 H), 2.5-2.7 (m, 1 H), 2.8-2.9 (t, 2 H), 3.05-3.15 (t, 2 H) 4.75 (s, 2 H), 6.65 (d, 1 H), 6.8 (d, 1 H)
C-25		Gen. Method 4b	(CDCl ₃): δ 1.0 (d, 6 H), 2.4-2.5 (m, 1 H), 2.7-2.8 (m, 2 H), 2.8-2.9 (m, 2 H), 4.7 (s, 2 H), 6.75 (d, 1 H), 7.1 (d, 1 H)










-77-

TABLE D. Preparation of Ketones (cont'd)

			
Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-26		Gen. Method 5 (from C-25)	(CDCl ₃): δ 0.00 (s, 6 H), 0.82 (s, 9 H), 0.95 (d, 6 H), 2.44 (m, 1 H), 2.62 (t, 2 H), 2.72 (t, 2 H), 4.70 (s, 2 H), 6.70 (d, 1 H), 7.01 (d, 1 H)
C-27		Gen. Method 5 (from C-23)	(CDCl ₃): δ 0.09 (s, 6 H), 0.85 (s, 9 H), 0.92 (d, 6 H), 2.58 (m, 1 H), 2.73 (m, 2 H), 2.82 (m, 2 H), 4.81 (s, 2 H), 6.74 (s, 1 H), 6.82 (s, 1 H)
C-28		Gen. Method 3	(CDCl ₃): δ 1.08 (d, 6 H), 2.21 (d, 3 H), 2.53-2.62 (m, 1 H), 2.70 (t, 2 H), 2.89 (t, 2 H), 6.84-6.92 (m, 2 H), 7.03-7.10 (m, 1 H)
C-29		Gen. Method 3	
C-30		Gen. Method 3	(CDCl ₃): δ 1.05 (d, 6 H), 2.50-2.60 (m, 1 H), 2.71-2.76 (m, 2 H), 2.83-2.89 (m, 2 H), 6.95 (m, 2 H), 7.13 (m, 2 H)
C-31		Gen. Method 4a	¹ HNMR (CDCl ₃): δ 1.04 (d, 6H), 2.55 (sp, 1H), 2.64- 2.74 (m, 4H), 7.11-7.17 (m, 7H), 7.26-7.35 (m, 9H), 7.48 (s, 1H)







-78-

TABLE D. Preparation of Ketones (cont'd)

			
Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-32		Gen. Method 3	MS (APCI): 166 (M+H)
C-33		Gen. Method 3	MS (APCI): 167 (M+H)
C-34		Gen. Method 4a	MS (APCI): 179 (M+H)
C-35		Gen. Method 4a	MS (APCI): 194 (M+H)
C-36		Gen. Method 3	MS (APCI): 200 (M-H)
C-37	Ph		Prepared as reported in <i>Bull. Soc. Chim. Fr</i> , 1956:1653.
C-38		Gen. Method 4	MS (APCI): 206 (M)
C-39		Gen. Method 5 (from C-38)	MS (APCI): 319 (M-H)
C-40		Gen. Method 3	¹ HNMR (CDCl ₃): δ 1.09 (d, 6 H), 2.56-2.63 (m, 1 H), 2.84 (t, 2 H), 3.01 (t, 2 H), 6.49 (m, 1 H), 7.03- 7.06 (m, 1 H), 7.17 (m, 1 H), 7.29-7.33 (m, 1 H), 7.46 (s, 1 H), 8.23 (s, 1H)







-79-

TABLE D. Preparation of Ketones (cont'd)

			
Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-41		Gen. Method 4b	(CDCl ₃): δ 1.03 (d, 6 H), 2.51-2.58 (m, 1 H), 2.81-2.87 (m, 4 H), 4.60 (s, 2 H), 6.90 (d, 1 H), 7.17 (d, 1 H)
C-42		Gen. Method 5 (from ABOVE)	(CDCl ₃): δ 0.0 (s, 6 H), 0.83 (s, 9 H), 0.99 (d, 6 H), 2.50 (m, 1 H), 2.71 (m, 4 H), 4.56 (s, 2 H), 6.81 (d, 1 H), 7.06 (s, 1 H)
C-43		Gen. Method 4b	(CDCl ₃): δ 1.09 (d, 6 H), 2.19 (s, 3 H), 2.60 (m, 1 H), 2.75- 2.80 (m, 4 H), 6.85-6.90 (m, 2 H)
C-44		Gen. Method 4b	(CDCl ₃): δ 1.04-1.06 (d, 6 H), 2.13 (s, 3 H), 2.51- 2.58 (m, 1 H), 2.72-2.75 (t, 2 H), 2.94-2.98 (t, 2 H), 6.72-6.75 (d, 1 H), 6.97-6.98 (d, 1 H)
C-45		Gen. Method 3	(CDCl ₃): δ 1.06 (d, 6 H), 2.37 (s, 3 H), 2.52- 2.59 (m, 1 H), 2.67- 2.71 (t, 2 H), 2.77-2.81 (t, 2 H), 6.78-6.79 (d, 1 H), 6.98-7.00 (d, 1 H)








-80-

TABLE D. Preparation of Ketones (cont'd)

			
Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-46		Gen. Method 4a	(CDCl ₃): δ 1.05 (d, 6 H), 2.50-2.61 (m, 1 H), 2.88 (t, 2 H), 3.12 (t, 2 H), 4.62 (d, 2 H), 6.96 (d, 1 H), 7.08 (d, 1 H)
C-47		Gen. Method 5 (from C-46)	(CDCl ₃): δ 0.08 (s, 6 H), 0.91 (s, 9 H), 1.08 (d, 6 H), 2.50-2.70 (m, 1 H), 2.78-2.81 (t, 2 H), 3.03- 3.08 (t, 2 H), 4.63 (s, 2 H), 6.94 (d, 1 H), 7.03 (d, 1 H)
C-48		Gen. Method 3	(CDCl ₃): δ 1.09 (d, 6 H), 2.52-2.66 (m, 1 H), 2.71- 2.76 (m, 2 H), 2.90-2.95 (m, 2 H), 6.79-6.89 (m, 2 H), 7.09-7.19 (m, 1 H)
C-49		Gen. Method 3	(CDCl ₃): δ 1.07 (d, 6 H), 2.50-2.64 (m, 1 H), 2.73- 2.78 (m, 2 H), 2.84-2.90 (m, 2 H), 6.59-6.67 (m, 2 H), 7.69-7.72 (m, 1 H)
C-50		Gen. Method 3	(CDCl ₃): δ 1.06 (d, 6 H), 2.51-2.60 (m, 1 H), 2.71- 2.77 (m, 2 H), 2.84-2.89 (m, 2 H), 6.72-6.81 (m, 2 H), 7.12-7.20 (m, 1 H)

-81-

TABLE D. Preparation of Ketones (cont'd)

			
Example	R ₈	General Method	Analytical Data (¹ H NMR or MS)
C-51		Gen. Method 3	(CDCl ₃): δ 1.07 (d, 6 H), 2.49-2.61 (m, 1 H), 2.71- 2.76 (m, 2 H), 2.80-2.85 (m, 2 H), 6.74-6.84 (m, 2 H)
C-52		Gen. Method 4a	
C-53		Gen. Method 4a	(CDCl ₃): δ 1.10 (d, 6 H), 1.19 (t, 3 H), 2.54-2.64 (m, 3 H), 2.69-2.77 (m, 2 H), 2.82-2.90 (m, 2 H), 6.80-6.87 (m, 2 H), 7.08- 7.13 (m, 1 H)
C-54		Gen. Method 4a	(CDCl ₃): δ 1.04 (d, 6 H), 2.37 (bs, 1 H), 2.50-2.66 (m, 1 H), 2.77-2.82 (m, 2 H), 2.84-2.94 (m, 2 H), 4.70 (s, 2 H), 6.88-6.91 (m, 1 H), 7.08-13 (m, 2 H)
C-55		Gen. Method 5 (from C-54)	
C-56		Gen. Method 4a	

-82-

Example C-57

2,2,2-Trifluoro-N-[4-(4-methyl-3-oxo-pentyl)-thiazol-2-yl]-acetamide

A solution of C-18 (from Table D above; 1.21 g, 5.03 mmol) in 6N HCl (50 mL) and THF (5 mL) was refluxed for 4 hours and then cooled to room temperature. Solid NaHCO₃ was added portionwise with caution until pH 7.2 was achieved. The suspension was extracted with EtOAc. The combined extracts were washed with brine, dried (MgSO₄), and concentrated. The residue was chromatographed over silica gel, eluting with EtOAc, to give the deprotected compound. ¹H NMR (CDCl₃): δ 1.04 (d, 6 H), 2.52-2.59 (m, 1 H), 2.75 (m, 4 H), 6.06 (s, 1 H).

A solution of the ketone prepared above (0.75 g, 3.8 mmol) in CH₂Cl₂ (50 mL) was cooled in an ice bath, treated with NEt₃ (0.6 mL, 4.3 mmol) and trifluoromethyl acetic anhydride (0.6 mL, 4.3 mmol), and allowed to warm to room temperature. H₂O was added. The organic layer was separated, washed with brine, and dried (MgSO₄). Concentration gave an oil which was chromatographed over silica gel, eluting with EtOAc, to give the title compound. ¹H NMR (CDCl₃): δ 1.10 (d, 6 H), 2.58-2.65 (m, 1 H), 2.83-2.86 (m, 2 H), 2.91-2.95 (m, 2 H), 6.61 (s, 1 H).

Names corresponding to the ketones from Table D above are:

- C-7: 4-Methyl-1-thiazol-2-yl-pentan-3-one;
- C-8: 4-Methyl-1-thiazol-4-yl-pentan-3-one;
- C-9: 4-Methyl-1-thiazol-5-yl-pentan-3-one;
- C-10: 4-Methyl-1-(5-methyl-thiazol-2-yl)-pentan-3-one;
- C-11: 4-Methyl-1-(4-methyl-thiazol-2-yl)-pentan-3-one;
- C-12: 4-Methyl-1-(2-methyl-thiazol-4-yl)-pentan-3-one;
- C-13: 4-Methyl-1-(5-methyl-thiazol-4-yl)-pentan-3-one;
- C-14: 1-(2-Isopropyl-thiazol-4-yl)-4-methyl-pentan-3-one;
- C-15: 1-(4-Isopropyl-thiazol-5-yl)-4-methyl-pentan-3-one;
- C-16: 1-(5-Isopropyl-thiazol-4-yl)-4-methyl-pentan-3-one;
- C-17: 1-(4-Isopropyl-thiazol-2-yl)-4-methyl-pentan-3-one;

-83-

- C-18: N-[4-(4-Methyl-3-oxo-pentyl)-thiazol-2-yl]-acetamide;
C-19: 4-Methyl-1-pyridin-2-yl-pentan-3-one;
C-20: 1-Furan-2-yl-4-methyl-pentan-3-one;
C-21: 4-Methyl-1-(tetrahydro-furan-2-yl)-pentan-3-one;
5 C-22: 4-Methyl-1-thiophen-2-yl-pentan-3-one;
C-23: 1-(5-Hydroxymethyl-thiophen-3-yl)-4-methyl-pentan-3-one;
C-24: 1-(5-Hydroxymethyl-thiophen-2-yl)-4-methyl-pentan-3-one;
C-25: 1-(2-Hydroxymethyl-thiophen-3-yl)-4-methyl-pentan-3-one;
C-26: 1-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-4-
10 methyl-pentan-3-one;
C-27: 1-[5-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-4-
methyl-pentan-3-one;
C-28: 1-(3-Fluoro-2-methyl-phenyl)-4-methyl-pentan-3-one;
C-29: 1-(2-Fluoro-phenyl)-4-methyl-pentan-3-one;
15 C-30: 1-(4-Fluoro-phenyl)-4-methyl-pentan-3-one;
C-31: 4-Methyl-1-(1-trityl-1H-pyrazol-3-yl)-pentan-3-one;
C-32: 4-Methyl-1-(1H-pyrrol-3-yl)-pentan-3-one;
C-33: 1-(1H-Imidazol-4-yl)-4-methyl-pentan-3-one;
C-34: 4-Methyl-1-(pyrimidin-5-yl)-pentan-3-one;
20 C-35: 1-(2-Amino-pyrimidin-5-yl)-4-methyl-pentan-3-one;
C-36: 4-(4-Methyl-3-oxo-pentyl)-benzonitrile;
C-37: 4-Methyl-1-phenyl-pentan-3-one;
C-38: 1-(2-Hydroxymethyl-phenyl)-4-methyl-pentan-3-one;
C-39: 1-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-phenyl]-4-methyl-
25 pentan-3-one.
C-40: 1-(1H-Indol-5-yl)-4-methyl-pentan-3-one;
C-41: 1-(4-Hydroxymethyl-thiophen-3-yl)-4-methyl-pentan-3-one;
C-42: 1-[4-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-4-
methyl-pentan-3-one;
30 C-43: 4-Methyl-1-(4-methyl-thiophen-3-yl)-pentan-3-one;
C-44: 4-Methyl-1-(3-methyl-thiophen-2-yl)-pentan-3-one;
C-45: 4-Methyl-1-(2-methyl-thiophen-3-yl)-pentan-3-one;
C-46: 1-(3-Hydroxymethyl-thiophen-2-yl)-4-methyl-pentan-3-one;

-84-

C-47: 1-[3-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-2-yl]-4-methyl-pentan-3-one;

C-48: 1-(2,6-Difluoro-phenyl)-4-methyl-pentan-3-one;

C-49: 1-(3,5-Difluoro-phenyl)-4-methyl-pentan-3-one;

5 C-50: 1-(2,4-Difluoro-phenyl)-4-methyl-pentan-3-one;

C-51: 1-(3,4,5-Trifluoro-phenyl)-4-methyl-pentan-3-one;

C-52: 1-(5-Fluoro-2-methyl-phenyl)-4-methyl-pentan-3-one;

C-53: 1-(2-Ethyl-5-fluoro-phenyl)-4-methyl-pentan-3-one;

C-54: 1-(4-Fluoro-2-hydroxymethyl-phenyl)-4-methyl-pentan-3-one;

10 C-55: 1-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-4-fluoro-phenyl]-4-methyl-pentan-3-one; and

C-56: 1-(4-Fluoro-2-methyl-phenyl)-4-methyl-pentan-3-one.

General Method 6. Preparation of β -Ketoesters

15 Methyl acetoacetate was added dropwise to a slurry of hexane washed sodium hydride in anhydrous THF at 0°C and the reaction stirred at 0°C (15 minutes to 3 hours). N-Butyl lithium (nBuLi) was then added at 0°C and the reaction stirred at 0°C (15 minutes to 24 hours). A solution of the requisite ketone in THF was added, and the reaction mixture was stirred at 0°C to room
20 temperature for 15 minutes to 24 hours. To the reaction mixture was added acetic acid [or dilute HCl or saturated ammonium chloride (NH₄Cl)] with stirring, and the THF was removed on a rotoevaporator. The viscous reaction mixture was partitioned between H₂O and EtOAc. After separation of the layers, the aqueous layer was again extracted with EtOAc. The combined organic extracts were dried (MgSO₄) and concentrated. The aldol intermediates were either purified by flash
25 chromatography or taken on crude.

General Method 7. Desilylation of Silyl Ether Protecting Groups

The appropriate silanyloxy compound (1 equiv.) was added to a reaction vessel followed by THF (3-5 mL per mmol of silanyloxy compound). This solution was treated with tetrabutylammonium fluoride (1.2-2.0 equiv.) and stirred
30 at room temperature (1 hour to 1 day). The product was partitioned between

-85-

EtOAc and 1N HCl. The organic layer was dried (MgSO₄) and concentrated. The product was either flashed chromatographed or carried on crude.

Example D-1

5-Hydroxy-6-methyl-5-[2-(4-methyl-thiazol-5-yl)-ethyl]-3-oxo-heptanoic acid methyl ester

5


The title compound was prepared as described in General Method 6 from 1.00 g (8.61 mmol) of methyl acetoacetate, 0.38 g (9.50 mmol) of sodium hydride, 4.5 mL of 2.1 M nBuLi (9.45 mmol), and 1.53 g (7.75 mmol) of 4-methyl-1-(4-methyl-thiazol-5-yl)-pentan-3-one (prepared in Example C-2.) The crude

10

compound was used without purification in the next step.

The following compounds were prepared in similar fashion from the appropriate ketone (from Examples C-1 to C-57) and were used without purification.













-86-

TABLE E. Preparation of β -Ketoesters

Example	R ₉
D-2	
D-3	
D-4	
D-5	
D-6	
D-7	
D-8	
D-9	
D-10	
D-11	


-87-











TABLE E. Preparation of β -Ketoesters (cont'd)

	
Example	R ₉
D-12	
D-13	
D-14	
D-15	
D-16	
D-17	
D-18	
D-19	
D-20	
D-21	
D-22	

-88-













TABLE E. Preparation of β -Ketoesters (cont'd)




Example	R ₉
D-23	
D-24	
D-25	
D-26	
D-27	
D-28	
D-29	
D-30	
D-31	
D-32	

-89-

TABLE E. Preparation of β -Ketoesters (cont'd)

<div></div>	
Example	R ₉
D-33	
D-34	
D-35	
D-36	
D-37	
D-38*	
D-39	
D-40	
D-41	
D-42	
D-43	

-90-

TABLE E. Preparation of β -Ketoesters (cont'd)

Example	R ₉
D-44	
D-45	
D-46	
D-47	
D-48	
D-49	
D-50	
D-51	
D-52**	

* Can be prepared from D-37 as outlined in General Method 7

** Can be prepared from D-51 as outlined in General Method 7

-91-

The compounds from Table E above are named:

D-2: 5-Hydroxy-6-methyl-3-oxo-5-(2-thiazol-2-yl-ethyl)-heptanoic acid
methyl ester;

5 D-3: 5-Hydroxy-6-methyl-3-oxo-5-(2-thiazol-4-yl-ethyl)-heptanoic acid
methyl ester;

D-4: 5-Hydroxy-6-methyl-3-oxo-5-(2-thiazol-5-yl-ethyl)-heptanoic acid
methyl ester;

D-5: 5-Hydroxy-6-methyl-5-[2-(5-methyl-thiazol-2-yl)-ethyl]-3-oxo-
heptanoic acid methyl ester;

10 D-6: 5-Hydroxy-6-methyl-5-[2-(4-methyl-thiazol-2-yl)-ethyl]-3-oxo-
heptanoic acid methyl ester;

D-7: 5-Hydroxy-6-methyl-5-[2-(2-methyl-thiazol-4-yl)-ethyl]-3-oxo-
heptanoic acid methyl ester;

15 D-8: 5-Hydroxy-6-methyl-5-[2-(5-methyl-thiazol-4-yl)-ethyl]-3-oxo-
heptanoic acid methyl ester;

D-9: 5-Hydroxy-5-[2-(2-isopropyl-thiazol-4-yl)-ethyl]-6-methyl-3-oxo-
heptanoic acid methyl ester;

D-10: 5-Hydroxy-5-[2-(4-isopropyl-thiazol-5-yl)-ethyl]-6-methyl-3-oxo-
heptanoic acid methyl ester;

20 D-11: 5-Hydroxy-5-[2-(5-isopropyl-thiazol-4-yl)-ethyl]-6-methyl-3-oxo-
heptanoic acid methyl ester;

D-12: 5-Hydroxy-5-[2-(4-isopropyl-thiazol-2-yl)-ethyl]-6-methyl-3-oxo-
heptanoic acid methyl ester;

25 D-13: 5-[2-(2-Acetylamino-thiazol-4-yl)-ethyl]-5-hydroxy-6-methyl-3-
oxo-heptanoic acid methyl ester;

D-14: 5-Hydroxy-6-methyl-3-oxo-5-(2-pyridin-2-yl-ethyl)-heptanoic acid
methyl ester;

D-15: 5-Hydroxy-6-methyl-3-oxo-5-(2-pyridin-4-yl-ethyl)-heptanoic acid
methyl ester;

30 D-16: 5-Hydroxy-6-methyl-3-oxo-5-(2-pyridin-3-yl-ethyl)-heptanoic acid
methyl ester;

D-17: 5-(2-Furan-2-yl-ethyl)-5-hydroxy-6-methyl-3-oxo-heptanoic acid
methyl ester;

-92-

D-18: 5-(2-Furan-3-yl-ethyl)-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-19: 5-Hydroxy-6-methyl-3-oxo-5-[2-(tetrahydro-furan-2-yl)-ethyl]-heptanoic acid methyl ester;

5 D-20: 5-Hydroxy-6-methyl-3-oxo-5-(2-thiophen-3-yl-ethyl)-heptanoic acid methyl ester;

D-21: 5-Hydroxy-6-methyl-3-oxo-5-(2-thiophen-2-yl-ethyl)-heptanoic acid methyl ester;

10 D-22: 5-{2-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-23: 5-{2-[5-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-24: 5-{2-[5-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-2-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

15 D-25: 5-Hydroxy-5-[2-(5-hydroxymethyl-thiophen-3-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

D-26: 5-Hydroxy-5-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

20 D-27: 5-Hydroxy-5-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

D-28: 5-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-29: 5-[2-(2-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

25 D-30: 5-[2-(4-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-31: 5-Hydroxy-6-methyl-3-oxo-5-[2-(1-trityl-1H-pyrazol-3-yl)-ethyl]-heptanoic acid methyl ester;

30 D-32: 5-Hydroxy-6-methyl-3-oxo-5-[2-(1H-pyrrol-3-yl)-ethyl]-heptanoic acid methyl ester;

D-33: 5-Hydroxy-5-[2-(1H-imidazol-4-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

-93-

D-34: 5-Hydroxy-6-methyl-3-oxo-5-(2-pyrimidin-5-yl-ethyl)-heptanoic acid methyl ester;

D-35: 5-[2-(2-Amino-pyrimidin-5-yl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

5 D-36: 5-[2-(4-Cyano-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-37: 5-{2-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-phenyl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

10 D-38: 5-Hydroxy-5-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

D-39: 5-Hydroxy-5-[2-(1H-indol-5-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

D-40: 5-Hydroxy-6-methyl-3-oxo-5-{2-[2-(2,2,2-trifluoro-acetylamino)-thiazol-4-yl]-ethyl}-heptanoic acid methyl ester;

15 D-41: 5-Hydroxy-6-methyl-5-[2-(4-methyl-thiophen-3-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;

D-42: 5-Hydroxy-6-methyl-5-[2-(3-methyl-thiophen-2-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;

20 D-43: 5-{2-[3-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-2-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-44: 5-Hydroxy-5-[2-(3-hydroxymethyl-thiophen-2-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

D-45: 5-[2-(2,6-Difluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

25 D-46: 5-[2-(3,5-Difluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-47: 5-[2-(2,4-Difluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

30 D-48: 5-[2-(3,4,5-Trifluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-49: 5-[2-(5-Fluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

-94-

D-50: 5-[2-(2-Ethyl-5-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

D-51: 5-{2-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-4-fluoro-phenyl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester; and

5 D-52: 5-Hydroxy-5-[2-(4-fluoro-2-hydroxymethyl-phenyl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester.

General Method 8. Preparation of the Intermediate 4-Hydroxy-5,6-dihydro-pyran-2-ones

10 The aldol intermediate was dissolved in THF (1 volume) and treated with 9 to 10 volumes of NaOH (0.1N-1.0N). The reaction was stirred from 1 hour to 24 hours at room temperature. The base solution was extracted with Et₂O and then cooled to 0°C. The mixture was acidified to pH 4 to 5 using HCl (0.1N to 6N) or acetic acid. On occasion, the product could be isolated by filtration. Alternatively the acidified extracts were extracted with EtOAc. The organic
15 extracts were combined, dried (MgSO₄) and concentrated. Purification was accomplished by trituration from Et₂O or flash chromatography.

Example E-1

4-Hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one

20 The title compound was prepared as described in General Method 8 using the crude aldol product isolated in Example D-1, 20 mL of f THF, and 200 mL of 1.0N NaOH. The reaction mixture was stirred at room temperature for 2 hours, then poured over ice and acidified to pH 5.1. The solution was extracted with EtOAc, dried (MgSO₄), and concentrated. Purification by silica gel
25 chromatography, eluting with 5:95 MeOH:CH₂Cl₂, gave the title compound.

¹H NMR (DMSO-d₆): δ 0.87-0.90 (m, 6 H), 1.82-1.96 (m, 2 H), 2.07-2.18 (m, 1 H), 2.25 (s, 3 H), 2.31 (d of ABX q, 1 H), 2.60 (d of ABX q, 1 H), 2.74-2.81 (m, 2 H), 4.96 (s, 1 H), 8.78 (s, 1 H), 11.39 (s, 1 H).

-95-

The following dihydropyrones were prepared from the corresponding aldol products from Examples D-1 to D-51. The 4-hydroxy-dihydropyrones exist in different forms depending upon the solvent.












TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones

Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-2		i-Propyl	Gen. Method 8	MS (APCI): 268 (M+H)
E-3		i-Propyl	Gen. Method 8	MS (APCI): 268 (M+H)
E-4		i-Propyl	Gen. Method 8	MS (APCI): 268 (M+H)
E-5		i-Propyl	Gen. Method 8	MS (APCI): 282 (M+H)
E-6		i-Propyl	Gen. Method 8	MS (APCI): 282 (M+H)
E-7		i-Propyl	Gen. Method 8	MS (APCI): 282 (M+H)





-96-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-8		i-Propyl	Gen. Method 8	MS (APCI): 282 (M+H)
E-9		i-Propyl	Gen. Method 8	MS (APCI): 310 (M+H)
E-10		i-Propyl	Gen. Method 8	MS (APCI): 310 (M+H)
E-11		i-Propyl	Gen. Method 8	MS (APCI): 310 (M+H)
E-12		i-Propyl	Gen. Method 8	MS (APCI): 310 (M+H)
E-13		i-Propyl	Gen. Method 8	MS (APCI): 325 (M+H)
E-14		i-Propyl	Gen. Method 8	MS (APCI): 262 (M+H)

-97-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-15		i-Propyl	Gen. Method 8	(CDCl ₃): δ 1.05 (d, 6 H), 1.7-1.9 (m, 1 H), 2.0-2.2 (m, 2 H), 2.6-2.9 (AB q, 2 H), 2.7-2.8 (m, 2 H), 3.42 (s, 2 H), 7.05 (d, 2 H), 8.5 (d, 2 H).
E-16		i-Propyl	Gen. Method 8	(CDCl ₃): δ 1.00 (d, 6 H), 1.7-1.9 (m, 1 H), 1.95-2.2 (m, 2 H), 2.6-2.8 (AB q, 2 H), 2.6-2.8 (m, 2 H), 3.40 (s, 2 H), 7.15-7.25 (m, 1 H), 7.4-7.5 (m, 1 H), 8.4-8.5 (s,m, 2 H).
E-17		i-Propyl	Gen. Method 8	(CDCl ₃): δ 1.1 (dd, 6 H), 1.7-1.9 (m, 1 H), 2.0-2.2 (m, 2 H), 2.6-2.8 (AB q, 2 H), 2.7-2.9 (m, 2 H), 3.40 (s, 2 H), 6.0 (m, 1 H), 6.30 (m, 1 H), 7.3 (m, 1 H).

-98-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)













				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-18		i-Propyl	Gen. Method 8	(CDCl ₃): δ 1.1 (dd, 6 H), 1.7-1.9 (m, 1 H), 1.95-2.2 (m, 2 H), 2.5-2.7 (m, 2 H), 2.6-2.8 (AB q, 2 H), 3.40 (s, 2 H), 6.25 (s, 1 H), 7.1 (m, 1 H), 7.18 (m, 1 H).
E-19		i-Propyl	Gen. Method 8	(CDCl ₃): δ 0.8-2.2 (m, 15 H), 2.65 (AB q, 2 H), 3.40 (AB q, 2 H), 3.60-3.90 (m, 3 H).
E-20		i-Propyl	Gen. Method 8	Mp 140-142.5°C
E-21		i-Propyl	Gen. Method 8	(CDCl ₃): δ 1.05 (d, 6 H), 1.8-2.0 (m, 1 H), 2.0-2.2 (m, 2 H), 2.6-2.8 (AB q, 2 H), 2.9-3.1 (m, 2 H), 3.40 (s, 2 H), 6.8 (m, 1 H), 6.9 (m, 1 H), 7.15 (m, 1 H).

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-22		i-Propyl	Gen. Method 7 then Gen. Method 8	(CDCl ₃): δ 1.1 (d, 6 H), 1.8-2.0 (m, 1 H), 2.0-2.2 (m, 2 H), 2.6-2.8 (AB q, 2 H), 2.8-3.0 (m, 2 H), 3.40 (s, 2 H), 4.75 (s, 2 H), 6.65 (d, 1 H), 6.8 (d, 1 H).
E-23		i-Propyl	Gen. Method 7 then Gen. Method 8	(CDCl ₃): δ 1.0 (dd, 6 H), 1.7-1.9 (m, 2 H), 2.0-2.2 (m, 2 H), 2.6-2.8 (m, Ab q, 4 H), 3.40 (s, 2 H), 4.8 (d, 2 H), 6.8 (s, 1 H), 6.9 (s, 1 H).
E-24		i-Propyl	Gen. Method 7 then Gen. Method 8	MS (APCI): 295 (M-H)







-100-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-25		i-Propyl	Gen. Method 8	(DMSO-d ₆): δ 0.90 (d, 3 H), 0.92 (d, 3 H), 1.73-1.90 (m, 2 H), 2.10-2.19 (m, 4 H), 2.28 (d of ABX q, 1 H), 2.58-2.65 (m, 3 H), 4.97 (s, 1 H), 6.92-6.98 (m, 2 H), 7.08-7.15 (m, 1 H), 11.39 (bs, 1 H).
E-26		i-Propyl	Gen. Method 8	(DMSO-d ₆): δ 0.89 (d, 3 H), 0.91 (d, 3 H), 1.81-1.95 (m, 2 H), 2.08-2.17 (m, 1 H), 2.28 (d of ABX q, 1 H), 2.58-2.64 (m, 3 H), 4.96 (s, 1 H), 7.07-7.14 (m, 2 H), 7.19-7.28 (m, 2 H), 11.37 (bs, 1 H).












-101-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-27		i-Propyl	Gen. Method 8	(DMSO-d ₆): δ 0.88 (d, 3 H), 0.90 (d, 3 H), 1.82-1.91 (m, 2 H), 2.05-2.14 (m, 1 H), 2.32 (d of ABX q, 1 H), 2.54-2.62 (m, 3 H), 4.96 (s, 1 H), 7.04-7.11 (m, 2 H), 7.18-7.22 (m, 2 H), 11.35 (bs, 1 H).
E-28		i-Propyl	Gen. Method 8	MS (APCI): 250 (M)
E-29		i-Propyl	Gen. Method 8	MS (APCI): 250 (M+H)
E-30		i-Propyl	Gen. Method 8	MS (APCI): 493 (M+H)
E-31		i-Propyl	Gen. Method 8	MS (APCI): 263 (M+H)










-102-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-32		i-Propyl	Gen. Method 8	MS (APCI): 278 (M+H)
E-33		i-Propyl	Gen. Method 8	MS (APCI): 286 (M+H)
E-34		i-Propyl	Gen. Method 8	MS (APCI): 290 (M)
E-35		i-Propyl	08/883,743	
E-36		Cyclopentyl		
E-37		Cyclohexyl		
E-38		i-Propyl		
E-39		Cyclopentyl		
E-40		Cyclohexyl		
E-41		i-Propyl		





-103-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-42		Cyclopentyl	From 08/883,743	
E-43		Cyclohexyl	From 08/883,743	
E-44		i-Propyl	From 08/883,743	
E-45		Cyclopentyl	From 08/883,743	
E-46		Cyclohexyl	From 08/883,743	
E-47		i-Propyl	Gen. Method 8	MS (APCI): 300 (M+H)
E-48		i-Propyl	Gen. Method 8	MS (APCI): 379 (M+H)
E-49		i-Propyl	Gen. Method 8	MS (APCI): 281 (M+H)




-104-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-50		i-Propyl	Gen. Method 8	MS (APCI): 281 (M+H)
E-51		i-Propyl	Gen. Method 8	(CDCl ₃): δ 1.02-1.06 (dd, 6 H), 1.80- 2.00 (m, 1 H), 2.05-2.25 (m, 2 H), 2.61-2.79 (AB q, 2 H), 2.85-3.05 (m, 2 H), 3.34-3.52 (AB q, 2 H), 4.60 (d, 2 H), 6.97 (d, 1 H), 7.10 (d, 1 H).
E-52		i-Propyl	Gen. Method 8	(CDCl ₃) δ 0.88 (d, 3 H), 0.91 (d, 3 H), 1.74-1.94 (m, 2 H), 2.10-2.19 (m, 1 H), 2.28 (d of ABX q, 1 H), 2.58-2.65 (m, 3 H), 4.97 (s, 1 H), 6.92-6.98 (m, 2 H), 7.08-7.15 (m, 1 H), 11.39 (bs, 1 H)




-105-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-53		i-Propyl	Gen. Method 8	(CDCl ₃) δ 0.88 (d, 3 H), 0.91 (d, 3 H), 1.87-1.93 (m, 2 H), 2.04-2.13 (m, 1 H), 2.33 (d of ABX q, 1 H), 2.60-2.69 (m, 3 H), 4.96 (s, 1 H), 7.00-7.04 (m, 3 H), 11.35 (bs, 1 H)
E-54		i-Propyl	Gen. Method 8	(DMSO) δ 0.89 (d, 3 H), 0.91 (d, 3 H), 1.77-1.94 (m, 2 H), 2.07-2.16 (m, 1 H), 2.29 (d of ABX q, 1 H), 2.57-2.63 (m, 3 H), 4.95 (s, 1 H), 6.95-7.01 (m, 1 H), 7.11-7.19 (m, 1 H), 7.27-7.35 (m, 1 H), 11.36 (bs, 1 H)




-106-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-55		i-Propyl	Gen. Method 8	(CDCl ₃) δ 0.87 (d, 3 H), 0.89 (d, 3 H), 1.86-1.92 (m, 2 H), 2.01-2.15 (m, 1 H), 2.34 (d of ABX q, 1 H), 2.53-2.61 (m, 3 H), 4.94 (s, 1 H), 7.15-7.24 (m, 2 H), 11.36 (bs, 1 H).
E-56		i-Propyl	Gen. Method 8	(DMSO) δ 0.89 (d, 3 H), 0.91 (d, 3 H), 1.77-1.85 (m, 2 H), 2.08-2.16 (m+s, 4 H), 2.29 (d of ABX q, 1 H), 2.53-2.64 (m, 3 H), 4.96 (s, 1 H), 6.84-6.96 (m, 2 H), 7.10-7.15 (m, 1 H), 11.36 (bs, 1 H)



-107-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-57		i-Propyl	Gen. Method 8	(DMSO) δ 0.89 (d, 3 H), 0.91 (d, 3 H), 1.09 (t, 3 H), 1.80-1.87 (m, 2 H), 2.08-2.17 (m, 1 H), 2.28 (d of ABX q, 1 H), 2.47-2.51 (m, partially obscured by DMSO, 2 H), 2.54-2.64 (m, 3 H), 4.96 (s, 1 H), 6.88-6.95 (m, 2 H), 7.15-7.18 (m, 1 H), 11.38 (bs, 1 H).
E-58		i-Propyl	Gen. Method 8	

-108-

TABLE F. Preparation of Racemic 4-Hydroxy-Dihydropyrones (cont'd)

				
Example	Aryl	R ₁₀	General Method	Analytical Data (¹ H NMR or MS)
E-59		i-Propyl	Gen. Method 7 (from E-58)	(DMSO) δ 0.95 (d, 3 H), 0.93 (d, 3 H), 1.84-1.94 (m, 2 H), 2.14-2.26 (m, 1 H), 2.35 (d of ABX q, 1 H), 2.53-2.62 (m, partially obscured by DMSO, 2 H), 2.66 (d of ABX q, 1 H), 4.52 (s, 2 H), 5.03 (s, 1 H), 5.32 (bs, 1 H), 6.99-7.05 (m, 1 H), 7.15-7.22 (m, 2 H), 11.42 (bs, 1 H)

The compounds from Table F above are named:

E-2: 4-Hydroxy-6-isopropyl-6-(2-thiazol-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

5 E-3: 4-Hydroxy-6-isopropyl-6-(2-thiazol-4-yl-ethyl)-5,6-dihydro-pyran-2-one;

E-4: 4-Hydroxy-6-isopropyl-6-(2-thiazol-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

E-5: 4-Hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

10 E-6: 4-Hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

-109-

E-7: 4-Hydroxy-6-isopropyl-6-[2-(2-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

E-8: 4-Hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

5 E-9: 4-Hydroxy-6-isopropyl-6-[2-(2-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

E-10: 4-Hydroxy-6-isopropyl-6-[2-(4-isopropyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

10 E-11: 4-Hydroxy-6-isopropyl-6-[2-(5-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

E-12: 4-Hydroxy-6-isopropyl-6-[2-(4-isopropyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

E-13: N-{4-[2-(4-Hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl)-ethyl]-thiazol-2-yl}-acetamide;

15 E-14: 4-Hydroxy-6-isopropyl-6-(2-pyridin-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

E-15: 4-Hydroxy-6-isopropyl-6-(2-pyridin-4-yl-ethyl)-5,6-dihydro-pyran-2-one;

20 E-16: 4-Hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

E-17: 6-(2-Furan-2-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

E-18: 6-(2-Furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

25 E-19: 4-Hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-5,6-dihydro-pyran-2-one;

E-20: 4-Hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

30 E-21: 4-Hydroxy-6-isopropyl-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

E-22: 4-Hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-110-

E-23: 4-Hydroxy-6-[2-(5-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

E-24: 4-Hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

5 E-25: 6-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

E-26: 6-[2-(2-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

10 E-27: 6-[2-(4-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

E-28: 4-Hydroxy-6-isopropyl-6-[2-(1H-pyrazol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

E-29: 4-Hydroxy-6-isopropyl-6-[2-(1H-pyrrol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

15 E-30: 4-Hydroxy-6-isopropyl-6-[2-(1-trityl-1H-imidazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

E-31: 4-Hydroxy-6-isopropyl-6-(2-pyrimidin-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

20 E-32: 6-[2-(2-Amino-pyrimidin-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

E-33: 4-[2-(4-Hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl)-ethyl]-benzonitrile;

E-34: 4-Hydroxy-6-isopropyl-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

25 E-47: 4-Hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

E-48: 2,2,2-Trifluoro-N-{4-[2-(4-hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl)-ethyl]-thiazol-2-yl}-acetamide;

30 E-49: 4-Hydroxy-6-isopropyl-6-[2-(4-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

E-50: 4-Hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

-111-

E-51: 4-Hydroxy-6-[2-(3-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

E-52: 6-[2-(2,6-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

5 E-53: 6-[2-(3,5-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

E-54: 6-[2-(2,4-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

10 E-55: 6-[2-(3,4,5-Trifluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

E-56: 6-[2-(5-Fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

E-57: 6-[2-(2-Ethyl-5-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

15 E-58: 6-{2-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-4-fluoro-phenyl]-ethyl}-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

E-59: 6-[2-(4-Fluoro-2-hydroxymethyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one.

20 **Alternate Synthesis of 4-Hydroxy-5,6-dihydro-pyran-2-ones via β -Hydroxyesters.**

General Method 9. Preparation of β -Hydroxyesters from Ketones (Table D) or Enones (Table B)

Diisopropylamine (1.25-1.4 equiv.) was cooled to -10°C and treated with nBuLi (1.25 equiv.) over 10 to 20 minutes. The solution was stirred for 15 to 25 45 minutes at -10°C and then cooled to -60°C to -78°C . The desired acetate (1.25 equiv.) was dissolved in THF and added dropwise to the LDA solution over 30 to 90 minutes. When addition was complete, the reaction mixture was stirred at -78°C to -40°C for another 30 to 90 minutes. The appropriate ketone (from Examples C-1 to C-39) or enone (from Examples B-1 to B-30) was dissolved in 30 THF and added over 15 to 30 minutes. The reaction mixture was warmed to room

-112-

temperature and stirred for 3 to 18 hours. The solution was poured into 1N HCl:ice; the product was extracted into EtOAc, dried (MgSO₄), and concentrated. The product could be purified via chromatography or recrystallization.

Example F-1







5 **3-Hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid tert-butyl ester**

The title compound was prepared as described in General Method 9 using t-butyl acetate (42.6 mmol), diisopropyl amine (42.6 mmol), nBuLi (42.6 mmol), 4-methyl-1-thiophen-3-yl-pentan-3-one (Example C-5; 21.3 mmol), and THF (100 mL). The title compound was purified by flash chromatography over silica
10 gel eluting with hexane:EtOAc 97:3. ¹H NMR (CDCl₃): δ 0.95 (dd, 6 H), 1.47 (s, 9 H), 1.7-2.0 (m, 3 H), 2.36-2.54 (AB q, 2 H), 2.6-2.8 (m, 2 H), 6.93-6.95 (m, 2 H), 7.23-7.25 (m, 1 H).

The following hydroxyesters were prepared in similar fashion from the analogous ketones from Examples C-1 to C-57 or enones from Examples B-1 to
15 B-36:






-113-

TABLE G. Preparation of Hydroxyesters

$ \begin{array}{c} R_{11} \quad OH \\ \quad \\ C - C - OR_{12} \\ \quad \\ CH_3 \quad O \end{array} $			
Example	R ₁₁	R ₁₂	Analytical Data (¹ H NMR or MS)
F-2		CH ₂ Ph	(CDCl ₃): δ 0.94 (d, 3 H), 0.96 (d, 3 H), 1.79-1.88 (m, 1 H), 2.71 (s, 2 H), 4.00 (br s, 1 H), 5.07 (d, 1H), 5.14 (d, 1 H), 6.19 (d, 1 H), 6.58 (d, 1 H), 6.90-6.70 (m, 2 H), 7.06-7.09 (m, 1 H), 7.24-7.29 (m, 5 H), 7.36-7.38 (m, 1 H).
F-3		CH ₂ Ph	
F-4		CH ₂ Ph	MS (APCI): 393 (M+H)
F-5		CH ₂ Ph	MS (APCI): 393 (M+H)
F-6		CH ₂ Ph	MS (APCI): 393 (M+H)
F-7		CH ₂ Ph	MS (APCI): 411 (M+H)




-114-

TABLE G. Preparation of Hydroxyesters (cont'd)

$ \begin{array}{c} R_{11} \quad OH \\ \quad / \\ C \quad C \quad OR_{12} \\ \quad \\ CH_3 \quad O \end{array} $			
Example	R ₁₁	R ₁₂	Analytical Data (¹ H NMR or MS)
F-8		CH ₂ Ph	MS (APCI): 411 (M+H)
F-9		CMe ₃	MS (APCI): 323 (M+H)
F-10		CMe ₃	(CDCl ₃): δ 0.95 (d, 3 H), 0.97 (d, 3 H), 1.48 (s, 9 H), 1.75-1.86 (m, 2 H), 1.88-1.96 (m, 1 H), 2.40 (d, 1 H), 2.52 (d, 1 H), 2.63-2.73 (m, 2 H), 3.92 (s, 1 H), 7.16-7.21 (m, 3 H), 7.26-7.31 (m, 2 H).
F-11		CH ₂ Ph	MS (APCI): 326 (M+H)
F-12		CH ₂ Ph	(CDCl ₃): δ 0.95 (d, 3 H), 0.96 (d, 3 H), 1.76-1.88 (m, 1 H), 2.71 (s, 2 H), 3.98 (bs, 1 H), 5.05 (d, 1 H), 5.13 (d, 1 H), 6.10 (d, 1 H), 6.57 (d, 1 H), 6.95-7.03 (m, 2 H), 7.23-7.31 (m, 7 H).



-115-

TABLE G. Preparation of Hydroxyesters (cont'd)

$ \begin{array}{c} R_{11} \quad OH \\ \quad / \\ C \quad C \quad O \\ \quad \\ CH_3 \quad OR_{12} \end{array} $			
Example	R ₁₁	R ₁₂	Analytical Data (¹ H NMR or MS)
F-13		Et	(CDCl ₃): δ 0.00 (s, 9 H), 0.83 (s, 6 H), 0.99 (d, 6 H), 1.3 (t, 3 H), 1.7-2.0 (m, 3 H), 2.50 (dd, 2 H), 2.60 (t, 2 H), 3.71 (s, 1 H), 4.19 (q, 2 H), 4.66 (s, 2 H), 6.91 (d, 1 H), 7.2 (d, 1 H).
F-14		CMe ₃	(CDCl ₃): δ 0.93-0.97 (t, 6 H), 1.47 (s, 9 H), 1.6-1.80 (m, 2 H), 1.91-1.95 (m, 1 H), 2.37 (s, 3 H), 2.37-2.40 (d, 1 H), 2.48-2.51 (d, 1 H), 2.56-2.61 (m, 2 H), 3.92 (s, 1 H), 6.80 (d, 1 H), 6.99 (d, 1 H).
F-15		CH ₂ Ph	

-116-

TABLE G. Preparation of Hydroxyesters (cont'd)

$ \begin{array}{c} R_{11} \quad OH \\ \quad \\ C - C - C - OR_{12} \\ \quad \\ CH_3 \quad O \end{array} $			
Example	R ₁₁	R ₁₂	Analytical Data (¹ H NMR or MS)
F-16		CH ₂ Ph	(CDCl ₃) δ 0.93 (d, 3 H), 0.95 (d, 3 H), 1.77-1.87 (m, 1 H), 2.70 (s, 2 H), 4.00 (s, 1 H), 5.06 (d, 1 H), 5.15 (d, 1 H), 6.18 (d, 1 H), 6.52 (d, 1 H), 6.64-6.71 (m, 1 H), 6.75- 7.82 (m, 2 H), 7.24-7.28 (m, 3 H), 7.34-7.38 (m, 2 H).
F-17		CMe ₃	(CDCl ₃) δ 0.94 (d, 3 H), 0.96 (d, 3 H), 1.46 (s, 9 H), 1.60-1.75 (m, 2 H), 1.89-1.97 (m, 1 H), 2.30 (s, 3 H), 2.39 (d, 1 H), 2.50 (d, 1 H), 2.56- 2.67 (m, 2 H), 3.97 (s, 1H), 6.77-6.86 (m, 2 H), 7.04-7.08 (m, 1 H).

The compounds of Table G above are named:

F-2: 5-(3-Fluoro-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;

F-3: 5-(3,4-Difluoro-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;

F-4: 3-Hydroxy-3-isopropyl-5-(2-trifluoromethyl-phenyl)-pent-4-enoic acid benzyl ester;

F-5: 3-Hydroxy-3-isopropyl-5-(3-trifluoromethyl-phenyl)-pent-4-enoic acid benzyl ester;

-117-

F-6: 3-Hydroxy-3-isopropyl-5-(4-trifluoromethyl-phenyl)-pent-4-enoic acid benzyl ester;

F-7: 5-(3-Fluoro-5-trifluoromethyl-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;

5 F-8: 5-(4-Fluoro-3-trifluoromethyl-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;

F-9: 3-Hydroxy-3-[2-(2-hydroxymethyl-phenyl)-ethyl]-4-methyl-pentanoic acid tert-butyl ester;

10 F-10: 3-Hydroxy-4-methyl-3-(2-phenyl-ethyl)-pentanoic acid tert-butyl ester;

F-11: 3-Hydroxy-3-isopropyl-5-phenyl-3-pent-4-enoic acid benzyl ester;

F-12: 5-(4-Fluoro-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;

15 F-13: 3-{2-[4-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-ethyl}-3-hydroxy-4-methyl-pentanoic acid ethyl ester;

F-14: 3-Hydroxy-4-methyl-3-[2-(2-methyl-thiophen-3-yl)-ethyl]-pentanoic acid tert-butyl ester;

F-15: 5-(3-Fluoro-2-methyl-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;

20 F-16: 5-(3,5-Difluoro-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester; and

F-17: 5-(4-Fluoro-2-methyl-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid tert-butyl ester.

Separation of β -Hydroxy Ester Enantiomers Via Chiral HPLC

25 Example F-1 (S)

(S)-3-Hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid tert-butyl ester

30 The title compound was prepared by resolution on a Chiralpak AD column eluting with 1:99 isopropanol:hexane to afford both enantiomers of 3-hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid tert-butyl ester. The S enantiomer eluted first.

-118-

¹H NMR (CDCl₃): δ 0.95 (dd, 6 H), 1.47 (s, 9 H), 1.7-2.00 (m, 3 H), 2.36-2.54 (AB q, 2 H), 2.6-2.8 (m, 2 H), 6.93-6.95 (m, 2 H), 7.23-7.25 (m, 1 H).

Example F-1 (R)

(R)-3-Hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid tert-butyl ester

5

The title compound was prepared as described above via resolution of Compound F-9 on a Chiralpak AD column eluting with 1:99 isopropanol:hexane to afford both enantiomers of 3-hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid tert-butyl ester. The R enantiomer eluted second.

10 ¹H NMR (CDCl₃): δ 0.95 (dd, 6 H), 1.47 (s, 9 H), 1.7-2.00 (m, 3 H), 2.36-2.54 (AB q, 2 H), 2.6-2.8 (m, 2 H), 6.93-6.95 (m, 2 H), 7.23-7.25 (m, 1 H).

Example F-9 (S)

(S)-3-Hydroxy-4-methyl-3-[2-(2-hydroxymethyl-phenyl)-ethyl]-pentanoic acid tert-butyl ester

15

The title compound was prepared by resolution of Compound F-9 on a Chiralpak AD column eluting with isopropanol:hexane (1:99) to afford both enantiomers of 3-hydroxy-4-methyl-3-[2-(2-hydroxymethyl-phenyl)-ethyl]-pentanoic acid tert-butyl ester. MS (APCI): 323 (M+H).

Example F-14 (S)

20 **(S)-3-Hydroxy-4-methyl-3-[2-(2-methyl-thiophen-3-yl)-ethyl]-pentanoic acid tert-butyl ester**

The title compound was prepared by resolution on a Chiralpak AD column eluting with 0.75:0.25:99 ethanol:isopropanol:hexane to afford both enantiomers of 3-hydroxy-4-methyl-3-[2-(2-methyl-thiophen-3-yl)-ethyl]-pentanoic acid tert-butyl ester. The S enantiomer eluted first.

25

¹H NMR (CDCl₃): δ 0.93-0.97 (t, 6 H), 1.47 (s, 9 H), 1.6-1.80 (m, 2 H), 1.91-1.95 (m, 1 H), 2.37 (s, 3 H), 2.37-2.40 (d, 1 H), 2.48-2.51 (d, 1 H), 2.56-2.61 (m, 2 H), 3.92 (s, 1 H), 6.80 (d, 1 H), 6.99 (d, 1 H).

-119-

General Method 10. Preparation of β -Hydroxyacids

The appropriate β -hydroxyester (1 equiv.) was dissolved in EtOH and treated with 1.4 to 2.0 equivalents of an alkoxide (KOH or LiOH). The mixture was stirred at room temperature or heated to reflux for 4 to 18 hours. The solution
5 was concentrated to dryness, and the residue was partitioned between H₂O and Et₂O. The aqueous layer was separated, acidified with 1N HCl, and extracted with Et₂O. The solution was dried (MgSO₄) and concentrated.

Alternatively, the benzyl esters (compounds F-2 through F-8 and compounds F-11 and F-12) were hydrogenated as described in General Method 3
10 to give the corresponding acid.

Example G-1 (S)**(S)-3-Hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid**





The title compound was prepared as described in General Method 10 using (S)-3-hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid tert-butyl ester
15 (Example F-1 (S); 8 mmol), LiOH (16 mmol), H₂O (5 mL), and MeOH (15 mL).

¹NMR (CDCl₃): δ 0.98 (t, 6 H), 1.8-2.0 (m, 3 H), 2.50-2.70 (Ab q, 2 H), 2.6-2.8 (m, 2 H), 6.93-6.95 (m, 2 H), 7.23-7.25 (m, 1 H).

The following β -hydroxyacids were prepared from the corresponding esters, either by hydrolysis (General Method 10) or hydrogenolysis (as outlined in
20 General Method 3) as noted:








-120-

TABLE H. Synthesis of β -hydroxyacids

$ \begin{array}{c} \text{R}_{13} \quad \text{OH} \\ \quad \\ \text{C} - \text{CH}_2 - \text{COOH} \\ \\ \text{CH}_3 \end{array} $				
Example	R ₁₃	General Method	Chirality	Analytical Data (¹ H NMR or MS)
G-1 (R)		Gen. Method 10	R	(CDCl ₃): δ 0.98 (t, 6 H), 1.8-2.0 (m, 3 H), 2.50-2.70 (AB q, 2 H), 2.6-2.8 (m, 2 H), 6.93-6.95 (m, 2 H), 7.23-7.25 (m, 1 H).
G-1		Gen. Method 10	\pm	(CDCl ₃): δ 0.92-0.95 (m, 6 H), 1.78-2.02 (m, 3 H), 2.49-2.76 (m, 4 H), 6.91 (m, 2 H), 7.22 (m, 1 H).
G-2		Gen. Method 3	\pm	(CDCl ₃): δ 0.95-1.00 (m, 6 H), 1.71-2.03 (m, 3 H), 2.49-2.56 (m, 1 H), 2.62-2.73 (m, 3 H), 3.75-3.79 (m, 1 H), 6.84-6.91 (m, 2 H), 6.96 (d, 1 H), 7.18-7.27 (m, 1 H).
G-3		Gen. Method 3	\pm	(CDCl ₃): δ 0.94-0.99 (m, 6 H), 1.78-1.89 (m, 2 H), 1.92-2.01 (m, 1 H), 2.52 (d, 1 H), 2.61-2.70 (m, 3 H), 3.75-3.79 (m, 1 H), 6.85-6.91 (m, 1 H), 6.94-7.09 (m, 2 H).





-121-

TABLE H. Synthesis of β -hydroxyacids

$ \begin{array}{c} \text{R}_{13} \quad \text{OH} \\ \quad \\ \text{C} - \text{CH}_2 - \text{COOH} \\ \\ \text{CH}_3 \end{array} $				
Example	R ₁₃	General Method	Chirality	Analytical Data (¹ H NMR or MS)
G-4		Gen. Method 3	±	MS (APCI): 303 (M-H)
G-5		Gen. Method 3	±	MS (APCI): 303 (M-H)
G-6		Gen. Method 3	±	MS (APCI): 607 (2M-H)
G-7		Gen. Method 3	±	MS (APCI): 321 (M-H)
G-8		Gen. Method 3	±	MS (APCI): 321 (M-H)
G-9		Gen. Method 10	S	MS (APCI): 266 (M) Rotation: -3.8° (c=2.0, EtOH)
G-10		Gen. Method 10 (using F-10) or Gen. Method 3 (using F-11)	±	(CDCl ₃): δ 0.97 (d, 3 H), 0.99 (d, 3 H), 1.80-1.93 (m, 2 H), 1.95-2.05 (m, 1 H), 2.56 (d, 1 H), 2.65-2.74 (m, 3 H), 7.17-7.21 (m, 3 H), 7.26-7.32 (m, 2 H).




-122-

TABLE H. Synthesis of β -hydroxyacids (cont'd)

$ \begin{array}{c} \text{R}_{13} \quad \text{OH} \\ \quad \\ \text{---} \text{C} \text{---} \text{CH}_2 \text{---} \text{C}(=\text{O})\text{OH} \\ \\ \text{CH}_3 \end{array} $				
Example	R ₁₃	General Method	Chirality	Analytical Data (¹ H NMR or MS)
G-11		Gen. Method 3	±	(CDCl ₃): δ 0.96 (d, 3 H), 0.99 (d, 3 H), 1.76-1.89 (m, 2 H), 1.91-2.05 (m, 1 H), 2.55 (d, 1 H), 2.63-2.70 (m, 3 H), 3.74-3.78 (m, 1 H), 6.93-6.91 (m, 2 H), 7.11-7.16 (m, 2 H).
G-12		Gen. Method 10	±	MS (APCI): 385 (M-H)
G-13		Gen. Method 10	S	MS (APCI): 255 (M-H)
G-14		Gen. Method 3	±	(CDCl ₃) δ 0.97 (d, 3 H), 0.99 (d, 3 H), 1.66- 1.86 (m, 2 H), 1.96- 2.05 (m, 1 H), 2.21 (d, 3 H), 2.59 (d, 1 H), 2.65-2.73 (m, 3 H), 6.84-6.93 (m, 2 H), 7.03-6.10 (m, 1 H).

-123-

TABLE H. Synthesis of β -hydroxyacids (cont'd)

				
Example	R ₁₃	General Method	Chirality	Analytical Data (¹ H NMR or MS)
G-15		Gen. Method 3	±	(CDCl ₃) δ 0.96 (d, 3 H), 0.98 (d, 3 H), 1.81-1.86 (m, 2 H), 1.87-2.01 (m, 1 H), 2.53 (d, 1 H), 2.64-2.78 (m, 3 H), 3.74-3.78 (m, 1 H), 6.59-6.67 (m, 1 H), 6.69-6.74 (m, 2 H).
G-16		Gen. Method 10	±	

The compounds of Table H above are named:

G-2: 3-[2-(3-Fluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;

G-3: 3-[2-(3,4-Difluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;

5 G-4: 3-Hydroxy-4-methyl-3-[2-(2-trifluoromethyl-phenyl)-ethyl]-pentanoic acid;

G-5: 3-Hydroxy-4-methyl-3-[2-(3-trifluoromethyl-phenyl)-ethyl]-pentanoic acid;

10 G-6: 3-Hydroxy-4-methyl-3-[2-(4-trifluoromethyl-phenyl)-ethyl]-pentanoic acid;

G-7: 3-[2-(3-Fluoro-5-trifluoromethyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;

G-8: 3-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;

-124-

G-9: (S)-3-Hydroxy-3-[2-(2-hydroxymethyl-phenyl)-ethyl]-4-methyl-pentanoic acid;

G-10: 3-Hydroxy-4-methyl-3-(2-phenyl-ethyl)-pentanoic acid;

G-11: 3-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;

5 G-12: 3-{2-[4-(tert-Butyl-dimethyl-silanyloxy-methyl)-thiophen-3-yl]-ethyl}-3-hydroxy-4-methyl-pentanoic acid;

G-13: 3-Hydroxy-4-methyl-3-[2-(2-methyl-thiophen-3-yl)-ethyl]-pentanoic acid;

10 G-14: 3-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;

G-15: 3-[2-(3,5-Difluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid; and

G-16: 3-[2-(4-Fluoro-2-methyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid.

15 **Resolution of Racemic β -hydroxyacids enantiomers**

Example G-2 (R)

(R)-3-[2-(3-Fluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

Racemic compound G-2 (42.0 g, 165 mmol) from Table H was dissolved in 410 mL EtOAc and treated with (S)- α -methylbenzylamine (10.01 g, 82.58 mmol). The resulting slurry was placed on a steam bath and heated to reflux
20 to dissolve the solids completely. The solution was allowed to cool to room temperature slowly overnight. The flask was then cooled to 5°C, and the solids were filtered and washed with cold EtOAc and Et₂O. The product was recrystallized from hot EtOAc (110 mL) on a steam bath, and then cooled slowly
25 to room temperature overnight. The flask was then cooled (5°C); the solids were filtered and washed with cold EtOAc followed by Et₂O to yield the desired compound. HPLC analysis of isolated solid (as the free acid): (AD chiralpak column, 1 mL/min., 97.5% Hexanes:2.5% IPA +0.1% TFA)

retention time: 19.60 min. (95.3%)

30 22.67 min. (4.7%)

-125-

Example G-2 (S)

(S)-3-[2-(3-Fluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

A second solid was isolated from the mother liquor from Example G-2 (R) after the solution was allowed to stand for 48 hours. The mixture was cooled to 0°C in ice bath for several hours. The solid was filtered off and washed with cold EtOAc followed by Et₂O. HPLC analysis of isolated solid (as the free acid):

(AD chiralpak column, 1 mL/min., 97.5% Hexanes:2.5% IPA +0.1% TFA)

retention time: 18.11 min. (11.76%)

19.28 min. (88.24%).

The isolated solid was recrystallized from hot EtOAc (~100 mL) on a steam bath and cooled slowly to room temperature overnight. The mixture then was cooled in an ice bath for 2 hours. The resulting solids were filtered and washed with cold EtOAc followed by Et₂O to give the desired enantiomer. The free acid was obtained by treating the salt with 1N HCl followed by extraction with EtOAc. The organic phased was washed with brine, dried (MgSO₄), filtered and concentrated to afford the resolved β-hydroxy acid. HPLC analysis of isolated solid (as the free acid): (AD chiralpak column, 1 mL/min., 97.5% Hexanes:2.5% IPA +0.1% TFA)

retention time: 18.20 min. (3.4%)

19.42 min. (96.6%)

Example G-3 (R)

(R)-3-[2-(3,4-Difluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

Compound G-3 (16.0 g, 58.8 mmol) from Table H was dissolved in 150 mL EtOAc and treated with (S)-α-methylbenzylamine (3.77 g, 31.14 mmol).

The thick suspension that resulted was placed on a steam bath and heated to reflux. Additional EtOAc (240 mL) was added, followed by hot isopropanol (approximately 80 mL), until the solids dissolved. The solution was allowed to cool to room temperature slowly overnight. The mixture was then cooled to approximately 0°C. The resulting solids were filtered and washed with cold EtOAc followed by Et₂O to yield the desired enantiomer. HPLC analysis of

-126-

isolated solid (as the free acid): (AD chiralpak column, 1 mL/min., 97.5% Hexanes:2.5% IPA + 0.1% TFA)

retention time: 23.80 min. (7.7%)
25.10 min. (92.3%).

The isolated solid was dissolved in hot IPA (120 mL) and EtOAc (600 mL) on a steam bath and then cooled slowly to room temperature overnight. The flask was cooled in an ice bath for several hours and then filtered. The solids were washed with cold EtOAc followed by Et₂O to yield the pure stereoisomer. The free acid was obtained by treating the salt with 1N HCl followed by extraction with EtOAc. The organic phase was washed with brine, dried (MgSO₄), filtered and concentrated to afford the resolved β-hydroxy acid. HPLC analysis of isolated solid (as the free acid): (AD chiralpak column, 1 mL/min., 97.5% Hexanes:2.5% IPA + 0.1% TFA) retention time: 25.90 min. (>99%).

15 Example G-3 (S)

(S)-3-[2-(3,4-Difluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

The mother liquor from the isolation of diastereomeric salt G-3 (R) was treated with 1N HCl and extracted with EtOAc. The organic phase was washed with brine, dried (MgSO₄), filtered and concentrated to liberate compound G-3. HPLC analysis indicated the isolated material was 78% (S):22% (R) based on comparison with previously isolated solid (diastereomeric salt G-3 (R)). β -Hydroxy acid G-3 (9.75 g, 35.8 mmol) was dissolved in 100 mL EtOAc and treated with (R)- α -methylbenzylamine (3.25 g, 26.86 mmol). The resulting thick suspension was placed on a steam bath and heated to reflux. Additional EtOAc (50 mL) was added, followed by hot IPA (100 mL), a little at a time until the solids dissolved. The solution was allowed to cool to room temperature overnight. The mixture was then cooled to 0°C in an ice bath. The resulting solids were filtered and washed with cold EtOAc followed by Et₂O. The free acid was obtained by treating the salt with 1N HCl followed by extraction with EtOAc. The organic phase was washed with brine, dried (MgSO₄), filtered and concentrated to afford the resolved β -hydroxy acid. HPLC analysis of isolated material (as the

-127-

free acid): (AD chiralpak column, 1 mL/min., 97.5% Hexanes:2.5% IPA + 0.1% TFA)

retention time: 22.06 min. (96.9%)

24.34 min. (3.1%).

5

Example G-10 (S)

(S)-3-Hydroxy-4-methyl-3-(2-phenyl-ethyl)-pentanoic acid

Compound G-10 (15.0 g, 63.48 mmol) was dissolved in 155 mL EtOAc and treated with (S)- α -methylbenzylamine (4.23 g, 34.91 mmol). The resulting slurry was placed on a steam bath and heated to reflux at which all solids
10 dissolved. The solution allowed to cool to room temperature slowly overnight, then cooled to -20°C. Cold EtOAc was added, and the solids were filtered and washed with cold EtOAc and Et₂O to yield the title compound. The free acid was obtained by treating the salt with 1N HCl followed by extraction with EtOAc. The organic phased was washed with brine, dried (MgSO₄), filtered and concentrated
15 to afford the resolved β -hydroxy acid. HPLC analysis of isolated solid (as the free acid): (OD chiralpak column, 1 mL/min., 97.5% Hexanes:2.5% IPA:+0.1% Formic acid)

retention time: 20.67 min. (4.4%)

22.24 min. (95.6%).

20

Example G-10 (R)

(R)-3-Hydroxy-4-methyl-3-(2-phenyl-ethyl)-pentanoic acid

The mother liquor from the isolation of diastereomeric salt G-10 (S) was treated with 1N HCl and extracted with EtOAc. The organic phase was washed with brine, dried (MgSO₄), filtered and concentrated to liberate the free
25 β -hydroxy acid. HPLC analysis indicated the isolated material was 73% (R):27% (S). β -Hydroxy acid G-10 (1 equiv.) was dissolved in 600 mL EtOAc and treated with (R)- α -methylbenzylamine (0.74 equiv.). The resulting suspension was placed on a steam bath and heated to reflux. The solution was allowed to cool to room temperature slowly and then cooled to 0°C in an ice bath.
30 The resulting solids were filtered off and washed with cold EtOAc followed by

-128-

Et₂O to yield the title compound. The free acid was obtained by treating the salt with 1N HCl followed by extraction with EtOAc. The organic phased was washed with brine, dried (MgSO₄), filtered and concentrated to afford the resolved β-hydroxy acid. HPLC analysis of isolated solid (as the free acid): (OD chiralpak column, 1 mL/min., 97.5% Hexanes:2.5% IPA:+0.1% Formic acid)
5 retention time: 21.86 min. (95.6%)
23.75 min. (4.4%).

Example G-11 (R)

(R)-3-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

10 Compound G-11 (37.1 g, 146 mmol) was dissolved in 375 mL EtOAc and treated with (S)-α-methylbenzylamine (4.23 g, 34.9 mmol). The resulting slurry was heated to reflux, diluted with an additional 50 to 100 mL EtOAc, and treated with hot isopropanol until the solids had completely dissolved. The solution was allowed to cool to room temperature slowly and then cooled to 0°C in an ice bath
15 for ~1 hour. Cold EtOAc was added; the solids were filtered off and washed with cold EtOAc and Et₂O to yield the title compound. HPLC analysis of isolated solid (as the free acid): (AS chiralpak column, 1 mL/min., 98% Hexanes:2% IPA:+0.1% Formic acid)
retention time: 16.81 min. (7.52%)
20 21.42 min. (92.48%).

The isolated solid and 450 mL EtOAc was heated to reflux and treated with hot isopropanol until the solids dissolved. The solution was allowed to cool to room temperature slowly overnight. The flask was then cooled to -10 to -15°C and treated with cold EtOAc. The solids were filtered off and washed with cold
25 EtOAc and Et₂O to yield the title compound. The free acid was obtained by treating the salt with 1N HCl followed by extraction with EtOAc. The organic phased was washed with brine, dried (MgSO₄), filtered and concentrated to afford the resolved β-hydroxy acid. HPLC analysis of isolated solid (as the free acid): (AS chiralpak column, 1 mL/min., 98% Hexanes:2% IPA:+0.1% Formic acid)
30 retention time: 16.74 min. (1.19%)
20.95 min. (98.81%).

-129-

Example G-11 (S)

(S)-3-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

The mother liquors from the isolation of the G-11 (R) enantiomer were combined, treated with 1N HCl and extracted with EtOAc. The organic phase was washed with brine, dried (MgSO₄), filtered and concentrated to liberate the free β -hydroxy acid. HPLC analysis indicated the isolated material was 79% (S):21% (R). β -Hydroxy acid G-11 (22.6 g, 89.0 mmol) was dissolved in 250 mL EtOAc and treated with (R)- α -methylbenzylamine (9.17 g, 75.64 mmol). The resulting mixture was placed on a steam bath, heated to reflux, diluted with additional ~100 mL EtOAc, and treated with hot isopropanol until solids completely dissolved. The solution was allowed to cool to room temperature slowly overnight and then cooled to -10 to -15°C. Cold EtOAc was added; the solids were filtered off and washed with cold EtOAc and Et₂O to yield the title compound. The free acid was obtained by treating the salt with 1N HCl followed by extraction with EtOAc. The organic phase was washed with brine, dried (MgSO₄), filtered and concentrated to afford the resolved β -hydroxy acid. HPLC analysis of isolated solid (as the free acid): (AS chiralpak column, 1 mL/min., 98% Hexanes:2% IPA:+0.1% Formic acid)

retention time: 16.61 min. (96.17%)

20.77 min. (3.83%).

Example G-14 (S)

(S)-3-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

Compound G-14 (28.3 g, 105 mmol) was dissolved in 270 mL EtOAc and treated with (R)- α -methylbenzylamine (6.52 g, 53.8 mmol). No precipitate formed. After 2 hours the solvent was evaporated. To the resulting residue was added 200 mL hexanes, and the flask was refluxed on a steam bath until the solids dissolved. Et₂O (20 mL) was added and the solution allowed to cool to room temperature slowly overnight. The resulting solids were filtered off and washed with cold hexanes/Et₂O. HPLC analysis of isolated solid (as the free acid): (AS chiralpak column, .8 mL/min., 98% Hexanes/2 % IPA/+0.1% Formic acid).

-130-

retention time: 16.38 min. (80.76%)

22.67 min. (19.24%)

The isolated solids, Et₂O (100 mL), and hexanes (200 mL) were heated to reflux on a steam bath. EtOAc was added until all solids dissolved (approximately 150 mL). An additional 200 mL of hexanes were added, and the mixture was brought back to reflux and then allowed to cool slowly to room temperature overnight. The flask was then cooled to approx. 5°C for 2 hours. The resulting solids were filtered and washed with cold Et₂O to yield the title compound. HPLC analysis of isolated solid (as the free acid): (AS chiralpak column, 1 mL/min., 98% Hexanes/2% IPA/+0.1% Formic acid).

retention time: 12.12 min. (94.8%)

24.10 min. (5.2%)

Example G-15 (S)

(S)-3-[2-(3,5-Difluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

Compound G-15 (24.6 g, 90.5 mmol) was dissolved in 235 mL EtOAc and treated with (R)- α -methylbenzylamine (6.03 g, 49.8 mmol). The resulting slurry was placed on a steam bath and heated to reflux until all solids dissolved. The solution was allowed to cool and hexanes were added (approximately 100 mL). The solution was returned to reflux, the steam turned off, and the mixture allowed to cool to room temperature overnight. The flask was then cooled to approx. 5°C for 1 hour. The resulting solids were filtered and washed with cold Et₂O. HPLC analysis of isolated solid (as the free acid): (AS chiralpak column, 1.0 mL/min., 98% Hexanes/2% IPA/+0.1% Formic acid).

retention time: 12.51 min. (92.3%)

14.83 min. (7.7%)

The solids were suspended in EtOAc (approximately 300 mL) and heated to reflux on a steam bath. When the solids did not dissolve, isopropanol (approximately 20 mL) was added gradually until the material completely dissolved. An additional 100 mL of EtOAc was added; the mixture was brought back to reflux and then allowed to cool to room temperature overnight. The flask was then cooled to approx. 5°C for 2 hours. The resulting solids were filtered and washed

-131-

with cold Et₂O to yield the title compound. HPLC analysis of isolated solid (as the free acid): (AS chirapak column, 1 mL/min., 98% Hexanes/2% IPA/+0.1% Formic acid).

retention time: 12.65 min. (98.0%)

5

15.10 min. (2.0%)

Example G-16 (S)

(S)-3-[2-(4-Fluoro-2-methyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid

Compound **G-16** (enriched - 62% (S)-enantiomer/38% (R) enantiomer) (11.1 g, 41.5 mmol) was dissolved in 120 mL EtOAc and treated with (S)- α -methylbenzylamine (6.52 g, 53.79 mmol). The resulting suspension was heated to reflux until all solids dissolved, at which time the solution was allowed to cool to room temperature overnight. The reaction flask was cooled to 0°C for 1 hour. The resulting solids were filtered off and washed with cold Et₂O. HPLC analysis of isolated solid (as the free acid): (AS chirapak column, .8 mL/min., 98% Hexanes/2% IPA/+0.1% Formic acid).

15

retention time: 15.86 min. (79.43%)

19.34 min. (20.57%)

The isolated solids were suspended in EtOAc (150 mL) and isopropanol (10 mL) and heated to reflux on a steam bath until all solids dissolved. The solution was allowed to cool to room temperature overnight and then cooled to 0°C for 1 hour. The resulting solids were filtered off and washed with cold Et₂O to afford the title compound. HPLC analysis of isolated solid (as the free acid): (AS chirapak column, .8 mL/min., 98% Hexanes/2 % IPA/+0.1% Formic acid).

20

25

retention time: 15.83 min. (92.0%)

19.34 min. (8.0%)

The isolated solids in EtOAc (130 mL) and isopropanol (6 mL) were again heated to reflux on a steam bath until the solids dissolved. The steam was shut off, and the solution was allowed to cool to room temperature overnight. The reaction flask was cooled to 0°C for 1 hour. The resulting solids were filtered off and washed with cold Et₂O to afford the title compound. HPLC analysis of isolated

30

-132-

solid (as the free acid): (AS chirapak column, .8 mL/min., 98% Hexanes/2% IPA/
+0.1% Formic acid).

retention time: 15.83 min. (96.7%)

19.34 min. (3.3%)

5

Example H

Bis [3-methoxy-3-oxopropanoato-(1-)-O,O'] magnesate

Monomethyl malonate (2.39 g, 20 mmol) and magnesium ethoxide
(1.16 g, 10 mmol) in THF (50 mL) were stirred for 3 hours at room temperature.
The solution was concentrated under vacuum to give the title compound. The
product was carried on crude to the next step.

10

General Method 11. Preparation of Chiral (or Racemic) β -Ketoester from β -Hydroxy acid for the Alternate Synthesis of intermediate 4-Hydroxy-5,6- dihydro-pyran-ones

A solution of the chiral β -hydroxyacid as isolated above (or the racemic
entity from Table H) (1.0 equiv.), THF (30 mL), and carbonyldiimidazole
(1.2-3 equiv.) was stirred for 3 to 18 hours at room temperature. Bis [3-methoxy-
3-oxopropanoato (1-)-O,O'] magnesate from Example H (1 to 2 equiv.) was
added, and the reaction was stirred for 6 to 72 hours at room temperature. The
reaction was concentrated and the residue partitioned between EtOAc and 1N
HCl. The organic layer was washed with aqueous NaHCO₃ and brine, dried
(MgSO₄) and concentrated. Purification was accomplished by silica gel
chromatography; alternatively, the material could be carried on crude.

15

20

Example I-1 (S)

(S)-5-Hydroxy-6-methyl-3-oxo-5-(2-thiophen-3-yl-ethyl)-heptanoic acid ethyl ester

25

The title compound was prepared as described in General Method 11 using
(S)-3-hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid (from
Example G-1 (S); 7.81 mmol), carbonyl diimidazole (8.59 mmol), bis
[3-methoxy-3-oxopropanoato(1-)-O,O']magnesate (15.62 mmol) and THF

-133-

(85 mL). Purification was accomplished using silica gel chromatography, eluting with 100% CH₂Cl₂.

¹H NMR (CDCl₃): δ 0.94 (dd, 6 H), 1.27 (t, 3 H), 1.75-1.90 (m, 2 H), 1.9-2.0 (m, 1 H), 2.65-2.75 (m, 3 H), 2.80 (d, 1 H), 3.48 (s, 2 H), 4.15-4.25 (m, 2 H), 6.93 (m, 2 H), 7.2-7.3 (m, 1 H).

Example I-2 (S)

5-[2-(3-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester, (S) enantiomer








The title compound was prepared following General Method 11 from compound G-2 (S) (5.85 g, 23.00 mmol) and CDI (8.21 g, 50.61 mmol) in 100 mL of THF. After 3.5 hours stirring under N₂ at room temperature, bis [3-methoxy-3-oxopropanoato(1-)-O,O']magnesate (13.94 g, 48.31 mmol) was added, and the mixture was stirred at room temperature overnight. The product was purified via silica gel chromatography, eluting with 4:1 hexanes:EtOAc to afford the title compound.

¹H NMR (CDCl₃): δ 0.93 (d, 3 H), 0.95 (d, 3 H), 1.27 (m, 3 H), 1.77-1.83 (m, 2 H), 1.91-2.01 (m, 1 H), 2.63-2.69 (m, 2 H), 2.71 (d, 1 H), 2.83 (d, 1 H), 3.48 (s, 2 H), 3.61 (s, 1 H), 4.19 (m, 2 H), 6.84-6.90 (m, 2 H), 6.95 (d, 1 H), 7.19-7.24 (m, 1 H).

The following β-ketoesters were prepared in similar fashion from the appropriate chiral (or racemic – Examples G-1 to G-16) β-ketoacids:






-134-

TABLE I. Preparation of β -Ketoesters

			
Example	R ₁₄	Chirality	Analytical Data (¹ H NMR or MS)
I-1 (R)		R	(CDCl ₃): δ 0.94 (dd, 6 H), 1.27 (t, 3 H), 1.75-1.90 (m, 2 H), 1.9-2.0 (m, 1 H), 2.65-2.75 (m, 3 H), 2.80 (d, 1 H), 3.48 (s, 2 H), 4.15-4.25 (m, 2 H), 6.93 (m, 2 H), 7.2-7.3 (m, 1 H).
I-2 (R)		R	
I-3 (R)		R	(CDCl ₃): δ 0.93 (d, 3 H), 0.95 (d, 3 H), 1.27 (m, 3 H), 1.77 (m, 2 H), 1.90-2.00 (m, 1 H), 2.62 (m, 2 H), 2.70 (d, 1 H), 2.83 (d, 1 H), 3.48 (s, 2 H), 3.61 (s, 1 H), 4.19 (m, 2 H), 6.85-6.90 (m, 1 H), 6.94-7.09 (m, 2 H).
I-3 (S)		S	
I-4		\pm	MS (APCI): 373 (M-H)
I-5		\pm	MS (APCI): 373 (M-H)





-135-

TABLE I. Preparation of β -Ketoesters (cont'd)

			
Example	R ₁₄	Chirality	Analytical Data (¹ H NMR or MS)
I-6		±	MS (APCI): 373 (M-H)
I-7		±	MS (APCI): 391 (M-H)
I-8		±	MS (APCI): 391 (M-H)
I-9 (S)		S	
I-10 (S)	Ph	S	(CDCl ₃): δ 0.93 (d, 3 H), 0.96 (d, 3 H), 1.26 (t, 3 H), 1.79-1.85 (m, 2 H), 1.93-2.02 (m, 1 H), 2.64-2.69 (m, 2 H), 2.72 (d, 1 H), 2.83 (d, 1 H), 3.48 (s, 2 H), 4.18 (q, 2 H), 7.15-7.20 (m, 3 H), 7.26-7.31 (m, 2 H).






-136-

TABLE I. Preparation of β -Ketoesters (cont'd)

			
Example	R ₁₄	Chirality	Analytical Data (¹ H NMR or MS)
I-10 (R)	Ph	R	(CDCl ₃): δ 0.93 (d, 3 H), 0.96 (d, 3 H), 1.26 (t, 3 H), 1.79-1.85 (m, 2 H), 1.93-2.02 (m, 1 H), 2.64-2.69 (m, 2 H), 2.72 (d, 1 H), 2.83 (d, 1 H), 3.47 (s, 2 H), 4.19 (q, 2 H), 7.15-7.20 (m, 3 H), 7.25-7.31 (m, 2 H).
I-11 (R)		R	(CDCl ₃): δ 0.93 (d, 3 H), 0.95 (d, 3 H), 1.26 (t, 3 H), 1.75-1.81 (m, 2 H), 1.91-2.01 (m, 1 H), 2.64-2.69 (m, 2 H), 2.71 (d, 1 H), 2.83 (d, 1 H), 3.48 (s, 2 H), 4.18 (q, 2 H), 6.92-6.98 (m, 2 H), 7.10-7.15 (m, 2 H).
I-11 (S)		S	(CDCl ₃): δ 0.93 (d, 3 H), 0.95 (d, 3 H), 1.26 (t, 3 H), 1.75-1.81 (m, 2 H), 1.91-2.00 (m, 1 H), 2.60-2.68 (m, 2 H), 2.71 (d, 1 H), 2.83 (d, 1 H), 3.48 (s, 2 H), 4.18 (q, 2 H), 6.92-6.98 (m, 2 H), 7.10-7.15 (m, 2 H).
I-12		\pm	

-137-

TABLE I. Preparation of β -Ketoesters (cont'd)

			
Example	R ₁₄	Chirality	Analytical Data (¹ H NMR or MS)
I-13 (S)		S	MS (APCI): 325 (M-H)
I-14 (S)		S	(CDCl ₃) δ 0.94 (d, 3 H), 0.96 (d, 3 H), 1.25 (q, 3 H), 1.69-1.76 (m, 2 H), 1.95-2.04 (m, 1 H), 2.12 (d, 3 H), 2.64-2.70 (m, 2 H), 2.72 (d, 1 H), 3.48 (s, 2 H), 3.61 (s, 1 H), 4.14-4.24 (m, 2 H), 6.82-6.92 (m, 2 H), 7.02-7.10 (m, 1 H).
I-15 (S)		S	
I-16 (S)		S	

The compounds of Table I above are named.

I-2: (R)-5-[2-(3-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

I-3: 5-[2-(3,4-Difluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester (R and S isomers);

I-4: 5-Hydroxy-6-methyl-3-oxo-5-[2-(2-trifluoromethyl-phenyl)-ethyl]-heptanoic acid ethyl ester;

I-5: 5-Hydroxy-6-methyl-3-oxo-5-[2-(3-trifluoromethyl-phenyl)-ethyl]-heptanoic acid ethyl ester;

-138-

I-6: 5-Hydroxy-6-methyl-3-oxo-5-[2-(4-trifluoromethyl-phenyl)-ethyl]-heptanoic acid ethyl ester;

I-7: 5-[2-(3-Fluoro-5-trifluoromethyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

5 I-8: 5-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

I-9: (S)-5-Hydroxy-5-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-methyl-3-oxo-heptanoic acid ethyl ester;

10 I-10: 5-Hydroxy-6-methyl-3-oxo-5-(2-phenyl-ethyl)-heptanoic acid ethyl ester (R and S isomers);

I-11: 5-[2-(4-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester (S and R isomers);

I-12: 5-{2-[4-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

15 I-13: (S)-5-Hydroxy-6-methyl-5-[2-(2-methyl-thiophen-3-yl)-ethyl]-3-oxo-heptanoic acid ethyl ester;

I-14: (S)-5-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

20 I-15: (S)-5-[2-(3,5-Difluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester; and

I-16: (S)-5-[2-(4-Fluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester.

**Alternate Synthesis of the Intermediate 4-Hydroxy-5,6-dihydro-pyran-2-ones
(for both chiral and racemic intermediates)**

25 Example J-1 (S)







(S)-4-Hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one

The title compound was prepared as described in General Method 8 using (S)-hydroxy-6-methyl-3-oxo-5-(2-thiophen-3-yl-ethyl)-heptanoic acid ethyl ester (6.4 mmol), 0.1N NaOH (400 mL), and THF (40 mL). The product was triturated
30 from Et₂O, mp 117-123°C.

The following chiral dihydropyrones were prepared in a similar fashion:

-139-

TABLE J. Synthesis of Chiral Dihydropyrones

			
Example	R ₁₅	Chirality	Analytical Data (¹ H NMR or MS)
J-1 (R)		R	mp 117-123°C
J-2 (S)		S	(DMSO-d ₆): δ 0.88 (d, 3 H), 0.90 (d, 3 H), 1.89-1.93 (m, 2 H), 2.08-2.15 (m, 1 H), 2.32 (d, 1 H), 2.56-2.62 (m, 3 H), 4.96 (s, 1 H), 6.95-7.04 (m, 3 H), 7.27-7.33 (m, 1 H), 11.37 (bs, 1 H).
J-2 (R)		R	(DMSO-d ₆): δ 0.88 (d, 3 H), 0.90 (d, 3 H), 1.88-1.93 (m, 2 H), 2.05-2.15 (m, 1 H), 2.33 (d, 1 H), 2.56-2.62 (m, 3 H), 4.96 (s, 1 H), 6.96-7.04 (m, 3 H), 7.26-7.33 (m, 1 H), 11.36 (bs, 1 H).
J-3 (S)		S	(DMSO-d ₆): δ 0.87 (d, 3 H), 0.90 (d, 3 H), 1.86-1.91 (m, 2 H), 2.04-2.13 (m, 1 H), 2.33 (d, 1 H), 2.55-2.61 (m, 3 H), 4.95 (s, 1 H), 6.99-7.03 (m, 1 H), 7.24-7.34 (m, 2 H).
J-3 (R)		R	(DMSO-d ₆): δ 0.87 (d, 3H), 0.90 (d, 3H), 1.86-1.91 (m, 2H), 2.04-2.13 (m, 1H), 2.33 (d, 1H), 2.55-2.61 (m, 3H), 4.95 (s, 1H), 6.99-7.03 (m, 1H), 7.24-7.34 (m, 2H).




-140-

TABLE J. Synthesis of Chiral Dihydropyrones (cont'd)

<div></div>			
Example	R ₁₅	Chirality	Analytical Data (¹ H NMR or MS)
J-4	<div></div>	±	MS (APCI): 329 (M+H)
J-5	<div></div>	±	MS (APCI): 329 (M+H)
J-6	<div></div>	±	MS (APCI): 329 (M+H)
J-7	<div></div>	±	MS (APCI): 347 (M+H)
J-8	<div></div>	±	MS (APCI): 347 (M+H)
J-9	<div></div>	S	(CDCl ₃): δ 1.05 (d, 3 H), 1.06 (d, 3 H), 1.8 (br.s, 1 H), 1.9 (m, 1 H), 2.03 (m, 1 H), 2.23 (m, 1 H), 2.66 (d, 1 H), 2.78 (d, 1 H), 2.81 (m, 2 H), 3.4 (d, 1 H), 3.53 (d, 1 H), 4.63 (d, 1 H), 4.72 (d, 1 H), 7.24 (m, 4 H),
J-10	<div></div>	S	FROM 08/883,743







-141-

TABLE J. Synthesis of Chiral Dihydropyrones (cont'd)

			
Example	R ₁₅	Chirality	Analytical Data (¹ H NMR or MS)
J-11 (S)	Ph	S	(DMSO-d ₆): δ 0.89 (d, 3 H), 0.91 (d, 3 H), 1.81-1.98 (m, 2 H), 2.07-2.16 (m, 1 H), 2.31 (d of ABX q, 1 H), 2.54-2.63 (m, 3 H), 4.96 (s, 1 H), 7.13-7.18 (m, 3 H), 7.23-7.28 (m, 2 H), 11.36 (bs, 1 H).
J-11 (R)	Ph	R	(DMSO-d ₆): δ 0.89 (d, 3 H), 0.91 (d, 3 H), 1.81-1.98 (m, 2 H), 2.07-2.16 (m, 1 H), 2.31 (d of ABX q, 1 H), 2.55-2.63 (m, 3 H), 4.97 (s, 1 H), 7.13-7.17 (m, 3 H), 7.23-7.28 (m, 2 H), 11.35 (bs, 1 H).
J-12 (R)		R	(DMSO-d ₆) 0.87 (d, 3 H), 0.89 (d, 3 H), 1.81-1.93 (m, 2 H), 2.05-2.14 (m, 1 H), 2.31 (d of ABX q, 1 H), 2.54-2.62 (m, 3 H), 4.95 (s, 1 H), 7.03-7.10 (m, 2 H), 7.17-7.22 (m, 2 H).
J-12 (S)		S	(DMSO-d ₆): δ 0.88 (d, 3 H), 0.90 (d, 3 H), 1.82-1.92 (m, 2 H), 2.05-2.14 (m, 1 H), 2.32 (d of ABX q, 1 H), 2.54-2.62 (m, 3 H), 4.96 (s, 1 H), 7.04-7.10 (m, 2 H), 7.17-7.22 (m, 2 H).

-142-

TABLE J. Synthesis of Chiral Dihydropyrones (cont'd)

			
Example	R ₁₅	Chirality	Analytical Data (¹ H NMR or MS)
J-13		±	MS (APCI): 295 (M-H)
J-14 (S)		S	MS (APCI): 279 (M-H)
J-15 (S)		S	(DMSO-d ₆) δ 0.89 (d, 3 H), 0.91 (d, 3 H), 1.75-1.87 (m, 2 H), 2.10-2.19 (m+d, 4 H), 2.69 (d, 1 H), 2.58-2.65 (m, 3 H), 4.96 (s, 1 H), 6.92-6.98 (m, 2 H), 7.08-7.15 (m, 1 H)
J-16 (S)		S	(DMSO-d ₆) δ 0.87 (d, 3 H), 0.89 (d, 3 H), 1.87-1.93 (m, 2 H), 2.04-2.13 (m, 1 H), 2.34 (d, 1 H), 2.54-2.65 (m, 3 H), 4.95 (s, 1 H), 6.91-7.04 (m, 3 H).
J-17 (S)		S	(DMSO-d ₆) δ 0.89 (d, 3 H), 0.91 (d, 3 H), 1.73-1.85 (m, 2 H), 2.08-2.17 (m, 1 H), 2.20 (s, 3 H), 2.28 (d, 1 H), 2.49-2.65 (m, partially obscured by DMSO, 3 H), 4.95 (s, 1 H), 6.86-6.98 (m, 2 H), 7.07-7.12 (m, 1 H).

-143-

The compounds of Table J above are named:

J-2: 6-[2-(3-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (R and S isomers);

5 J-3: 6-[2-(3,4-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (R and S isomers);

J-4: 4-Hydroxy-6-isopropyl-6-[2-(2-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

J-5: 4-Hydroxy-6-isopropyl-6-[2-(3-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

10 J-6: 4-Hydroxy-6-isopropyl-6-[2-(4-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

J-7: 6-[2-(3-Fluoro-5-trifluoromethyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

15 J-8: 6-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

J-9: (S)-4-Hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

J-11: 4-Hydroxy-6-isopropyl-6-(2-phenyl-ethyl)-5,6-dihydro-pyran-2-one (R and S isomers);

20 J-12: 6-[2-(4-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (R and S isomers);

J-13: 4-Hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

25 J-14: (S)-4-Hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

J-15: (S)-6-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

J-16: (S)-6-[2-(3,5-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

30 J-17: (S)-6-[2-(4-Fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one.

-144-

Synthesis of Tosylthiolate Side Chains**General Method 12. Preparation of Nitro Intermediates**

The appropriate alkylbenzene (1 equiv.) was dissolved in acetic acid and acetic anhydride. To the solution was added sulfuric acid, followed by 70% nitric acid or fuming nitric acid (1 to 1.5 equiv.; ratio of concentrated sulfuric acid to nitric acid is 1:2), dropwise at 0°C. After the addition, the reaction was warmed to room temperature. The reaction mixture was kept at room temperature to 80°C for 6 to 18 hours. The reaction mixture was poured into ice and extracted with EtOAc. The crude product was purified by either distillation or by flash silica gel chromatography.

Example K**1-Bromo-4-tert-butyl-2-nitro-benzene**

The reaction was done as described in General Method 12 using 1-bromo-4-tert-butyl-benzene (15 g), acetic acid (10 mL), nitric acid (6.5 g), and sulfuric acid (13 g). MS(APCI): 291.

General Method 13. Preparation of Benzthiazoles

To the aryl amine (1 equiv.) in MeOH or acetic acid was added sodium or potassium or ammonium thiocyanate (3-6 equiv.). The reaction was stirred at 0°C to room temperature. This reaction mixture was cooled to 0°C and treated dropwise with bromine (2-2.2 equiv.) or a solution of sodium bromide (2-2.2 equiv.) and bromine (2-2.2 equiv.). After the addition was complete, the reaction mixture was slowly warmed to room temperature and stirred at room temperature for 4 to 24 hours. The reaction mixture was added to sodium bicarbonate solution and extracted with EtOAc (2-4 times). The EtOAc layer was separated, washed with H₂O and brine, and dried (MgSO₄). The organic layer was concentrated, and the residue was purified either by crystallization or by flash silica gel chromatography.

-145-

· Example L-1

5-tert-Butyl-6-thiocyanato-benzothiazol-2-yl-amine

1-Bromo-4-tert-butyl-2-nitro-benzene (18 g) was reduced as described in General Method 3 using 10% palladium on charcoal (1 g) and MeOH (100 mL) in a hydrogen atmosphere (50 psi). The catalyst was filtered, and the solvents were evaporated. The residue was taken in EtOAc (100 mL) and washed with saturated sodium bicarbonate solution and H₂O. The solution was dried and concentrated. The residue was purified by flash chromatography to give 3-tert-butyl-phenylamine. MS (APCI): 150 (M+H).

The title compound was prepared according to General Method 13 using 3-tert-butyl-phenylamine (9.4 g, 63 mmol), sodium thiocyanate (30.7 g, 379 mmol), sodium bromide (14.3 g, 139 mmol), bromine (22.2 g, 139 mmol) and MeOH (300 mL). The crude reaction mixture was purified by flash chromatography (20% EtOAc in hexanes to 75% EtOAc in hexanes as eluents). MS (APCI): 264 (M+H).

Example L-2

5-iso-Propyl-6-thiocyanato-benzothiazol-2-yl-amine

The title compound was prepared according to General Method 13 using 3-isopropyl-phenylamine (20.8 g, 154 mmol), sodium thiocyanate (74.8 g, 923 mmol), sodium bromide (34.8 g, 338 mmol), bromine (54.1 g, 338 mmol), and MeOH (300 mL). The crude reaction mixture was purified by flash chromatography (CH₂Cl₂ to 25% EtOAc in CH₂Cl₂ as eluents). MS (APCI): 250 (M+H).

Examples L-3, L-4, and L-5

2-Isopropyl-4-isothiocyanato-5-methyl-phenylamine, 7-Isopropyl-4-methyl-6-thiocyanato-benzothiazol-2-yl-amine, and 4-Isopropyl-7-methyl-6-thiocyanato-benzothiazol-2-yl-amine

The title compounds were prepared according to General Method 13 using 2-isopropyl-5-methyl-phenylamine (contains approximately 10% of 3-isopropyl-6-methyl-phenylamine) (20.0 g, 134 mmol), sodium thiocyanate (65.3 g, 805 mmol), sodium bromide (30.4 g, 295 mmol), bromine (47.2 g, 295 mmol), and

-146-

MeOH (300 mL). The crude reaction mixture was purified by flash chromatography (20% EtOAc in hexanes to 75% EtOAc in hexanes as eluents), which yielded three products:

5 L-3: 2-Isopropyl-4-isothiocyanato-5-methyl-phenylamine: uppermost spot on tlc: MS (APCI): 207 (M+H).

L-4: 4-Isopropyl-7-methyl-6-thiocyanato-benzothiazol-2-yl-amine : middle sport on tlc: MS (APCI): 264 (M+H), 239.

L-5: 7-Isopropyl-4-methyl-6-thiocyanato-benzothiazol-2-yl-amine: lowest spot on tlc: MS (APCI): 264 (M+H)

10

Example L-6

6-tert-Butyl-4-thiocyanato-benzothiazol-2-yl-amine

The title compound was prepared according to General Method 13 using 4-tert-butyl-phenylamine (16.2 g, 109 mmol), sodium thiocyanate (53.0 g, 653 mmol), sodium bromide (24.6 g, 240 mmol), bromine (38.3 g, 240 mmol),
15 and MeOH (300 mL). The crude product was purified by silica gel chromatography (10-50% EtOAc in hexanes as eluents). MS (APCI): 263 (M).

General Method 14. Cleavage of SCN Group

A solution of the SCN intermediate in denatured EtOH was treated with dithiothreitol (3-5 equiv.) and 0.2 M KH_2PO_4 (in a ratio of 4:1 EtOH:buffer). The
20 mixture was stirred for 6 to 48 hours at 0°C to 50°C. The solvent was evaporated, and H_2O and CHCl_3 were added; the organic layer was washed with brine, dried (Na_2SO_4), filtered and concentrated. The resulting residue was flash chromatographed on silica gel.

General Method 14b. Alternate Cleavage of the SCN Group

25 The thiocyanate (1 eq.) was dissolved in THF and treated with sodium hydrogen sulfide (3 eq.) and sodium borohydride (6 eq.). A mixture of MeOH and water (2:1) was added dropwise at 0°C, and the reaction mixture was stirred at room temperature overnight. Water and toluene were added, and the reaction was quenched with glacial acetic acid at 0°C. The organic layer was separated, washed

-147-

with H₂O, and dried (MgSO₄.) The organic layer was concentrated, and the crude was used without any purification.

Example M-1

2-Amino-5-tert-butyl-benzothiazole-6-thiol

5 The title compound was prepared according to General Method 14 using 5-tert-butyl-6-thiocyanato-benzothiazol-2-yl-amine (Example L-1; 8 g, 30 mmol), dithiothreitol (18.7 g, 121 mmol), EtOH (200 mL), and phosphate buffer (50 mL). MS (APCI): 239 (M+H).

Example M-2

2-Amino-5-isopropyl-benzothiazole-6-thiol and 5-Isopropyl-benzothiazole-6-thiol

10 The title compound was prepared according to General Method 14 using 5-isopropyl-6-thiocyanato-benzothiazol-2-yl-amine (Example L-2; 23.5 g, 94.4 mmol), dithiothreitol (58.2 g, 377 mmol), EtOH (300 mL), and phosphate
15 buffer (150 mL). The reaction mixture was kept under reflux overnight. The reaction mixture contained a mixture of 2-amino-5-isopropyl-benzothiazole-6-thiol and 5-isopropyl-benzothiazole-6-thiol in 1:2 ratio. The crude reaction mixture was used as such without any further purification. 2-Amino-5-isopropyl-benzothiazole-6-thiol: MS (APCI): 225 (M+H); 5-Isopropyl-benzothiazole-
20 6-thiol: MS (APCI): 210 (M+H).

Example M-3

2-Amino-7-isopropyl-4-methyl-benzothiazole-6-thiol

 The title compound was prepared according to General Method 14 using 7-isopropyl-4-methyl-6-thiocyanato-benzothiazol-2-yl-amine (Example L-5;
25 11.5 g, 43.7 mmol), dithiothreitol (27.0 g, 175 mmol), EtOH (200 mL), and phosphate buffer (50 mL). MS (APCI): 239 (M+H).

-148-

Example M-4

2-Amino-6-tert-butyl-benzothiazole-4-thiol

The title compound was prepared according to General Method 14 using 6-tert-butyl-4-thiocyanato-benzothiazol-2-yl-amine (Example L-6; 2.0 g, 7.6 mmol), dithiothreitol (4.6 g, 30 mmol), phosphate buffer of pH 7 (10mL), and EtOH (50 mL). The crude product was used further without any purification. MS (APCI): 239 (M+H).

Example N

2-(Methyloxycarbonyl)amino-4-isopropyl-7-methyl-benzothiazole-6-thiol

To 4-isopropyl-7-methyl-6-thiocyanato-benzothiazol-2-yl-amine (Example L-4; 4.0 g, 17 mmol) dissolved in CH₂Cl₂ (25 mL) and pyridine (5 mL), was added methyl chloroformate at 0°C. The reaction mixture was stirred at room temperature for 2 hours. H₂O was added to the reaction mixture followed by EtOAc (100 mL). The organic layer was washed with 2N HCl, H₂O, saturated sodium bicarbonate solution and brine, and was dried (MgSO₄). The solvents were evaporated, and the residue was purified by flash chromatography (100% hexanes to 25% EtOAc in hexanes) to give 2-(methyloxycarbonyl)amino-4-isopropyl-7-methyl-6-thiocyanato-benzothiazole. MS (APCI): 322 (M+H).

The title compound was prepared according to General Method 14 using 2-(methyloxycarbonyl)amino-4-isopropyl-7-methyl-6-thiocyanato-benzothiazole (1.5 g, 4.7 mmol), dithiothreitol (2.9 g, 18.7 mmol), EtOH (50 mL), and phosphate buffer (15 mL). MS (APCI): 296 (M-H).

Preparation of Benzothiphenes

Example O

3-tert-Butyl-2-mercapto-benzo[b]thiophene

A solution of 3-tert-butylbenzo[b]thiophene (2.00 g, 10.4 mmol; *J. Chem. Soc., Perkin Trans. 1*, 1972;3:414-18) in Et₂O (30 mL, N₂ atmosphere) was treated with a 1.6 M hexane solution of nBuLi (3.75 mL, 12 mmol) and allowed to stir overnight. The mixture was then cooled to 0°C and treated with dry

-149-

precipitated sulfur (0.38 g, 12 mmol). The mixture was allowed to stir for 2 hours, then quenched with ice H₂O. The mixture was washed with Et₂O, and the aqueous layer was acidified with 1 N HCl and extracted with Et₂O. The organic layers were combined, dried (MgSO₄), and concentrated. The crude thiol was used without purification in the next step.

Example P

2-(3,5-Dibromo-thiophen-2-yl)-propan-2-ol

A solution of 2,3,5-tribromothiophene (8.17 g, 25 mmol) in Et₂O (N₂ atmosphere) was cooled to -78°C and treated with a 1.6 M hexane solution of nBuLi (15.6 mL, 25 mmol). The mixture was then stirred for 10 minutes, then treated with acetone (1.84 mL, 25 mmol). The mixture was then stirred for 30 minutes and quenched with H₂O. The mixture was extracted with Et₂O; the organic layers were combined, dried (MgSO₄), and the solvent removed in vacuo. The residue was filtered through a plug of silica gel (gradient elution from hexanes to EtOAc) to provide the title compound.

¹H NMR (CDCl₃): δ 1.70 (s, 6 H), 2.40 (br s, 1 H), 6.89 (s, 1 H).

Example Q

(4-Bromo-5-isopropyl-thiophen-2-yl)-methanol

A solution of 2-(3,5-dibromo-thiophen-2-yl)-propan-2-ol (Example P; 7.64 g, 25 mmol) in CH₂Cl₂ (100 mL) was cooled to 0°C and treated with triethylsilane (6.80 mL, 30.56 mmol) followed by trifluoroacetic acid (40 mL). The mixture was stirred for 15 minutes and then concentrated in vacuo. The residue was taken up in hexanes and filtered through a plug of silica gel. The solvent was removed in vacuo, and the residue was diluted with Et₂O and cooled to -78°C (N₂ atmosphere). The mixture was treated with a 1.6 M hexane solution of nBuLi (15.6 mL, 25 mmol), stirred 10 minutes, and treated with a solution of 4-formylmorpholine (2.88 mL, 25 mmol) in Et₂O (25 mL). The mixture was allowed to slowly come to room temperature overnight, then quenched with H₂O

-150-

and extracted with Et₂O. The organic layers were combined, dried (MgSO₄), and concentrated.

The residue was then taken up in MeOH (250 mL) and treated with an excess of sodium borohydride. After 30 minutes, the mixture was diluted with Et₂O, washed with H₂O, and dried (MgSO₄). The solvent was evaporated and the residue purified via silica gel chromatography (hexanes to 10% EtOAc:hexanes) to provide the title compound.

¹H NMR (CDCl₃): δ 1.28 (d, 6 H), 1.73 (bs, 1 H), 3.31 (sept, 1 H), 4.73 (d, 2 H), 6.82 (d, 1 H).

10

Example R

(4-Bromo-5-isopropyl-thiophen-2-yl-methoxy)-tert-butyl-dimethyl-silane

A solution of (4-bromo-5-isopropyl-thiophen-2-yl)-methanol (Example Q; 3.4 g, 14 mmol) in CH₂Cl₂ (4.0 mL) was treated with NEt₃ (2.8 mL, 20 mmol), p-dimethylaminopyridine (0.050 g), and t-butyldimethylsilylchloride (2.41 g, 16.0 mmol). The mixture was stirred for 14 hours, then diluted with Et₂O and washed with H₂O. The organic layer was dried (MgSO₄) and the solvent evaporated to provide the crude product, which was used without further purification.

¹H NMR (CDCl₃): δ 0.0 (s, 6 H), 0.82 (s, 9 H), 1.17 (d, 6 H), 3.20 (sept, 1 H), 4.66 (d, 2 H), 6.60 (s, 1 H).

20

Example S

5-(tert-Butyl-dimethyl-silanyloxymethyl)-2-isopropyl-3-mercapto-thiophene

A solution of (4-bromo-5-isopropyl-thiophen-2-ylmethoxy)-tert-butyl-dimethyl-silane (Example R; 4.7 g, 13.5 mmol) in Et₂O (100 mL, N₂ atmosphere) was cooled to -78°C and treated with a 1.6 M hexane solution of nBuLi (8.42 mL, 13.5 mmol). After 10 minutes, dry triturated sulfur (0.43 g, 13.5 mmol) was added, and the mixture was allowed to warm to room temperature. After 2 hours the mixture was diluted with Et₂O and washed with 0.5 M HCl. The organic layer

25

-151-

was dried (MgSO_4) and concentrated to give the title compound, which was used crude in the next step.

Example T

5-tert-Butyl-4-hydroxy-2-methylbenzaldehyde

5 Aluminum chloride (26.6 g, 200 mmol) was added to chlorobenzene (40 mL), and the mixture was cooled to -10°C . A solution of 2-tert-butyl-5-methylphenol (16.4 g, 100 mmol) in triethyl orthoformate (32.5 g, 220 mmol) was added dropwise. When addition was complete, the reaction mixture was warmed to 0°C to 5°C and kept at that temperature for 6 hours. Five percent HCl (100 mL) was added, and the mixture was extracted with 1:1 EtOAc:toluene. The organic phase was extracted with 23% KOH (3×50 mL) and water (25 mL), and the water extract was combined with the KOH extracts. The pH of the combined aqueous extracts was adjusted to 4 with 37% HCl. The resulting precipitate was filtered, washed with water (3×50 mL), and dried to give the title compound, 15 mp 171 - 171°C .

Example U

Dimethylthiocarbamic acid O-(2-tert-butyl-4-formyl-5-methylphenyl) ester

Ten percent KOH (80 mL) was added to a mixture of 5-tert-butyl-4-hydroxy-2-methylbenzaldehyde (Example T; 99.0 g, 515 mmol) in water (180 mL) and THF (135 mL.) The mixture was treated with a solution of 20 N,N-dimethylthiocarbamoyl chloride (86.1 g) in THF (108 mL) over a 2-hour period with simultaneous addition of 10% KOH in order to hold the pH of the reaction mixture between 12.0 and 12.3. After approximately 70% of the dimethylthiocarbamoyl chloride-THF solution had been added, vacuum was 25 applied to the reaction mixture to remove the THF; during this distillation, the simultaneous additions of 10% KOH and the dimethylthiocarbamoyl chloride-THF solution were continued while maintaining the pH range between 12.0 and 11.6 and the reaction temperature between 23°C to 25°C . The distillation was continued for 10 minutes after all the dimethylthiocarbamoyl chloride solution had 30 been added. Stirring at ambient pressure and temperature was continued for

-152-

another 90 minutes, during which time the pH was stable between 11.6 and 11.75. A solution of EtOAc (150 mL) and heptane (150 mL) was added to the reaction mixture, and after stirring and settling, the layers were separated and the aqueous layer extracted with 1:1 heptane:EtOAc. The combined organic extracts were
5 extracted with 10% KOH (2×100 mL) and water (2×90 mL) and concentrated. The residue was diluted with MeOH (120 mL). The solution was warmed to 45°C and H₂O (20 mL) was added. The solution was cooled to room temperature, 50% NaOH (170 mg) was added, and the solution was concentrated. This residue was redissolved in MeOH (120 mL) at 50°C and H₂O (15 mL) was added. The
10 solution was cooled slowly to -5°C with the addition of dimethylthiocarbamic acid O-(2-tert-butyl-4-formyl-5-methyl-phenyl) ester seed crystals to promote crystallization. The mixture was filtered and the solid washed with 4:1 MeOH:H₂O to give the title compound, mp 80-81°C.

Example V

15 (5-tert-Butyl-4-mercapto-5-methyl-phenyl)methanol

Dimethylthiocarbamic acid O-(2-tert-butyl-4-formyl-5-methylphenyl) ester (Example U; 24.0 g, 85.9 mmol) was treated with tetraethylene glycol dimethyl ether (65 g), and the mixture was stirred and heated under argon to 275°C. The solution was heated at this temperature for 30 minutes. The solution
20 was cooled, and H₂O was added. The mixture was cooled to 0°C and filtered; the solid was washed with H₂O to give dimethyl thiocarbamic acid S-(2-tert-butyl-4-formyl-5-methyl-phenyl) ester.

The crude solid was treated with MeOH (40 mL) and THF (30 mL) followed by slow addition (20 minutes) of NaBH₄ (14.1 g of a 12 weight percent solution in 14 M NaOH), while maintaining the temperature below 8°C. The
25 resulting solution was stirred at 20°C to 25°C for 90 minutes. Fifty percent NaOH (4 g) was added, and the solution was heated to reflux for 3 hours. The mixture was cooled to room temperature, treated with H₂O (110 mL) and toluene (70 mL), and acidified to pH 4 with 37% HCl. The aqueous layer was separated and

-153-

extracted with toluene (2×15 mL). Concentration gave the title compound which was used without further purification in the next step.

Example W

3-tert-Butyl-4-dimethylcarbamoylsulfanyl-benzoic acid methyl ester

5 A solution of 3.0 g (14.4 mmol) of 3-tert-butyl-4-hydroxy-benzoic acid methyl ester (*Aust. J. Chem.*, 1978;31:907-916), cesium carbonate (7.04 g, 21.6 mmol), and acetonitrile (50 mL) was heated to reflux and then treated with N,N-dimethylthiocarbamoyl chloride (2.67 g, 21.6 mmol) all at once. The mixture was refluxed for 2 hours, cooled to room temperature, and quenched with 1N HCl.

10 The solution was extracted with EtOAc; the organic layer was washed with 1N NaOH and brine, dried (MgSO_4), and concentrated. The residue was chromatographed over silica gel, eluting with 3:2 hexane:EtOAc, to give 3-tert-butyl-4-dimethylthiocarbamoyloxy-benzoic acid methyl ester.

^1H NMR (CDCl_3): δ 1.39 (s, 9 H), 3.44 (d, 6 H), 3.90 (s, 3 H), 7.09 (d, 1 H), 7.89 (dd, 1 H), 8.12 (m, 1 H).

15 Neat 3-tert-butyl-4-dimethylthiocarbamoyloxy-benzoic acid methyl ester (2.71 g, 9.2 mmol) was heated to 230°C for 2 hours and then cooled to room temperature. The material was chromatographed over silica gel, eluting with 7:3 hexane:EtOAc, to give the title compound.

20 ^1H NMR (CDCl_3): δ 1.50 (s, 9 H), 3.11 (br s, 6 H), 3.91 (s, 3 H), 7.53 (d, 1 H), 7.84 (dd, 1 H), 8.13 (d, 1 H).

Example X

(3-tert-Butyl-4-mercapto-phenyl)methanol

25 A solution of 2.1 g (7.1 mmol) of 3-tert-butyl-4-dimethylcarbamoylsulfanyl-benzoic acid methyl ester (Example W) in toluene (100 mL) was cooled to -78°C and treated with 42 mL of DIBAL (1.0 M in CH_2Cl_2 ; 42 mmol). The reaction mixture was allowed to warm to room temperature overnight. The reaction was quenched cautiously with saturated aqueous citric acid and extracted with EtOAc (2×100 mL). The combined

-154-

organic extracts were washed with brine, dried (MgSO₄), and concentrated.

Chromatography of the residue over silica gel, eluting with 3:2 hexane:EtOAc, gave the title compound.

¹H NMR (CDCl₃): δ 1.48 (s, 9 H), 1.97 (br s, 1 H), 3.61 (s, 1 H), 4.59 (s, 2 H),
5 7.02 (dd, 1 H), 7.20 (d, 1 H), 7.34 (d, 1 H).

Example Y

(5-tert-Butyl-4-hydroxy-2-methyl-phenyl)-acetic acid, methyl ester

A solution of 8.5 g (51 mmol) of (4-hydroxy-2-methyl-phenyl)-acetic acid
(*Indian J. Chem. Sect. B.* 26, 1987:679-682), tert-butanol (60 g), and conc.
10 sulfuric acid (2 mL) was stirred at 70°C for 5 days. The solution was cooled to
room temperature, poured into water, and extracted with EtOAc. The organic
extract was washed with brine, dried (MgSO₄), and concentrated to give (5-tert-
butyl-4-hydroxy-2-methyl-phenyl)-acetic acid.

¹H NMR (CDCl₃): δ 1.33 (s, 9 H), 2.14 (s, 3 H), 3.54 (s, 2 H), 6.47 (s, 1 H),
15 7.00 (s, 1 H).

A mixture of the (5-tert-butyl-4-hydroxy-2-methyl-phenyl)-acetic acid
isolated above (11.3 g, 51 mmol), MeOH (150 mL), and conc. sulfuric acid
(2 mL) was refluxed overnight. The solution was cooled to room temperature and
concentrated. The residue was partitioned between EtOAc and water; the organic
20 phase was washed with brine, dried (MgSO₄), and concentrated. The product was
chromatographed over silica gel, eluting with hexane:EtOAc, to give the title
compound.

¹H NMR (CDCl₃): δ 1.35 (s, 9 H), 2.16 (s, 3 H), 3.54 (s, 2 H), 3.67 (s, 3 H),
6.46 (s, 1 H), 7.01 (s, 1 H).

25

Example Z

(5-tert-Butyl-4-dimethylcarbamoylsulfanyl-2-methyl-phenyl)-acetic acid methyl ester

A solution of 3.0 g (13 mmol) of (5-tert-butyl-4-hydroxy-2-methyl)-phenyl
acetic acid, methyl ester (Example Y) in 20 mL of DMF was treated portionwise

-155-

with 0.60 g (15 mmol) of NaH and stirred at room temperature for 1 hour. N,N-dimethylthiocarbamoyl chloride (2.35 g, 19 mmol) was added all at once, and the reaction mixture was stirred at 70°C overnight. Water was added; the solution was extracted with EtOAc. The organic layer was separated, washed with
5 brine, dried (MgSO₄), and concentrated. The crude material was chromatographed over silica gel, eluting with hexane:EtOAc 3:1, to give (5-tert-butyl-4-dimethylthiocarbamoyloxy-2-methyl-phenyl)-acetic acid methyl ester.
¹H NMR (CDCl₃): δ 1.30 (s, 9 H), 2.22 (s, 3 H), 3.40 (d, 6 H), 3.58 (s, 2 H), 3.66 (s, 3 H), 6.80 (s, 1 H), 7.17 (s, 1 H).

10 The product isolated above (2.1 g, 6.5 mmol) was heated neat to 310°C for 1 hour and then cooled to room temperature. The residue was chromatographed on silica gel, eluting with hexane:EtOAc 2:1, to give the title compound.
¹H NMR (CDCl₃): δ 1.41 (s, 9 H), 2.22 (s, 3 H), 3.00-3.10 (br d, 6 H), 3.58 (s, 2 H), 3.65 (s, 3 H), 7.22-7.24 (m, 2 H).

15

Example AA

2-(5-tert-Butyl-4-mercapto-2-methyl-phenyl)-ethanol

A solution of 0.68 g (2.1 mmol) of (5-tert-butyl-4-dimethylcarbamoylsulfanyl-2-methyl-phenyl)-acetic acid methyl ester (Example Z) in 20 mL of toluene was cooled to -78°C and treated dropwise with
20 1.0 M DIBAL in CH₂Cl₂ (10 mL; 10 mmol). The cold bath was removed, and the mixture was stirred at room temperature for 2 hours. Saturated citric acid was added cautiously. H₂O and EtOAc were added, and the mixture was filtered. The filtrate was transferred to a separatory funnel where the aqueous phase was isolated and re-extracted with EtOAc. The organic extracts were combined,
25 washed with brine, and dried (MgSO₄). Concentration gave an oil which was chromatographed over silica gel, eluting with 1:1 hexane:EtOAc, to give the title compound.

¹H NMR (CDCl₃): δ 1.42 (s, 9 H), 2.20 (s, 3 H), 2.79 (t, 2 H), 3.49 (t, 1 H), 3.76 (t, 2 H), 7.01 (s, 1 H), 7.11 (s, 1 H).

-156-

General Method 15. Preparation of Tosyl Reagents

A solution of p-toluenesulfonyl bromide (1 equiv.), a base such as NEt₃ or pyridine (1-1.5 equiv.), and a solvent such as CCl₄ or EtOAc at 0°C to 24°C was treated with a solution of the appropriate thiol (1 equiv.) in CCl₄ or EtOAc in a dropwise fashion. Addition was complete in 0.5 to 11 hours. The reaction was allowed to warm to room temperature. H₂O was added; the organic layer was separated, washed with saturated aqueous NH₄Cl and brine, and dried (MgSO₄). Concentration gave a residue which could be purified either by silica gel chromatography or by trituration.

Example BB-1**Toluene-4-thiosulfonic acid S-(2-amino-5-tert-butyl-benzothiazol-6-yl)ester**

The title compound was prepared according to General Method 15 using 2-amino-5-tert-butyl-benzothiazole-6-thiol (7.0 g, 29 mmol), pyridine (2.3 g, 29 mmol), tosyl bromide (6.9 g, 29 mmol), carbon tetrachloride (100 mL), and EtOAc (100 mL). The crude reaction mixture was purified by flash chromatography (5% EtOAc in hexanes to 50% EtOAc in hexanes as eluents). MS(APCI): 393 (M+H).

Example BB-2**Toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl) ester**

A solution of (5-tert-butyl-4-mercapto-5-methyl-phenyl)methanol (from Example V), pyridine (6.12 G), and toluene (30 mL) was slowly added (over 3 hours) at 40°C to a prestirred solution of toluenesulfonyl chloride (16.6 g) and lithium bromide (8.0 g) in THF (75 mL). The resulting mixture was stirred at 40°C for 2 hours and then overnight at room temperature. H₂O (70 mL) and EtOAc (30 mL) were added. The organic layer was separated and washed with 10% HCl (2 × 50 mL) and H₂O (50 mL). The solution was concentrated to give a residue which was warmed to 60°C and treated with heptane. This solution was stirred and cooled to 10°C; the solids were filtered and washed with






-157-

1:1 toluene:heptane. Upon drying in vacuo, the title compound was obtained, mp 121-122°C.

The following tosylthiolates were prepared using the appropriate thiol and following General Method 15:






-158-

TABLE K. Thiotosylate Side Chains

Prep. of Starting Materials	Example	Structure	¹ H NMR or MS Data
Examples L-2 and M-2	BB-3		MS(APCI): 378
Examples L-2 and M-2	BB-4		MS(APCI): 364 (M+H)
Examples L-5 and M-3	BB-5		MS(APCI): 393 (M+H)
Examples L-4 and N	BB-6		MS(APCI): 451 (M+H)
Examples L-6 and M-4	BB-7		MS(APCI): 393 (M+H)





-159-

TABLE K. Thiotosylate Side Chains (cont'd)

Prep. of Starting Materials	Example	Structure	¹ H NMR or MS Data
Example O	BB-8		(CDCl ₃): δ 1.54 (s, 9 H), 2.43 (s, 3 H), 7.24 (d, 2 H), 7.32-7.39 (m, 2 H), 7.62 (d, 2 H), 7.69-7.75 (m, 1 H), 8.17-8.21 (m, 1 H).
Examples P-S	BB-9		(CDCl ₃): δ 0.0 (s, 6 H), 0.82 (s, 9 H), 0.91 (d, 6 H), 2.31 (s, 3 H), 2.93 (sept., 1 H), 4.65 (d, 2 H), 6.60 (s, 1 H), 7.13 (d, 2 H), 7.41 (d, 2 H).
	BB-10		USSN 08/883,743
	BB-11		USSN 08/883,743
	BB-12		USSN 08/883,743






-160-

TABLE K. Thiotosylate Side Chains (cont'd)

Prep. of Starting Materials	Example	Structure	¹ H NMR or MS Data
Examples W and X	BB-13		(CDCl ₃): δ 1.23 (s, 9 H), 2.40 (s, 3 H), 4.73 (s, 2 H), 7.23 (m, 3 H), 7.43 (m, 1 H), 7.46-7.50 (m, 2 H), 7.61 (d, 1 H).
Examples Y, Z, and AA	BB-14		(CDCl ₃): δ 1.21 (s, 9 H), 2.17 (s, 3 H), 2.39 (s, 3 H), 2.85 (t, 2 H), 3.80 (t, 2 H), 7.18-7.22 (m, 4 H), 7.47 (d, 2 H).
Example BB-11	BB-15		
Examples CC – FF	BB-16		(CDCl ₃): δ 1.28 (s, 9 H), 1.69 (s, 9 H), 2.4 (s, 3 H), 6.5 (d, 1 H), 7.2 (d, 2 H), 7.44 (d, 2 H), 7.61 (d, 1 H), 7.81 (s, 1 H), 8.28 (br s, 1 H).





-161-

TABLE K. Thiotosylate Side Chains (cont'd)

Prep. of Starting Materials	Example	Structure	¹ H NMR or MS Data
Examples GG - LL	BB-17		MS(APCI): 462 (M+H)
Examples MM - NN	BB-18		(CDCl ₃): δ 1.22 (s, 9 H), 2.44 (s, 3 H), 2.99 (t, 2 H), 4.03 (t, 2 H), 7.22 (d, 2 H), 7.39 (d, 2 H), 7.42 (s, 1 H), 7.66 (s, 1 H), 7.81 (d, 2 H), 7.94 (d, 2 H).
Examples OO-1 - VV-1	BB-19		MS(APCI): 347 (M+H).
Examples OO-2 - VV-2	BB-20		MS(APCI): 361 (M+H).
Examples OO-2 - VV-2	BB-21		MS(APCI): 375 (M- H).



-162-

TABLE K. Thiotosylate Side Chains (cont'd)

Prep. of Starting Materials	Example	Structure	¹ H NMR or MS Data
Examples WW - AAA	BB-22		MS(APCI): 432 (M+H)
Examples BBB - DDD	BB-23		MS(APCI): 405 (M+H)
Example BB-1	BB-24		MS(APCI): 558 (M+H)
Example BB-1	BB-25		MS(APCI): 471 (M+H)

-163-

TABLE K. Thiosylate Side Chains (cont'd)

Prep. of Starting Materials	Example	Structure	¹ H NMR or MS Data
Example BB-1	BB-26		MS(APCI): 435 (M+H)
Example VV-1	BB-27		MS(APCI): 363 (M+H)

The compounds from Table K above are named:

BB-3: Toluene-4-thiosulfonic acid S-(2-amino-5-isopropyl-benzothiazol-6-yl)ester;

BB-4: Toluene-4-thiosulfonic acid S-(5-isopropyl-benzothiazol-6-yl)ester;

5 BB-5: Toluene-4-thiosulfonic acid S-(2-amino-7-isopropyl-4-methyl-benzothiazol-6-yl)ester;

BB-6: Toluene-4-thiosulfonic acid S-(4-isopropyl-2-methoxycarbonylamino-7-methyl-benzothiazol-6-yl)ester;

10 BB-7: Toluene-4-thiosulfonic acid S-(2-amino-6-tert-butyl-benzothiazol-4-yl)ester;

BB-8: Toluene-4-thiosulfonic acid S-(3-tert-butyl-benzo[b]thiophen-2-yl)ester;

BB-9: Toluene-4-thiosulfonic acid S-[5-(tert-butyl-silanyloxymethyl)-2-isopropyl-thiophen-3-yl]ester;

15 BB-13: Toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-phenyl)ester;

-164-

BB-14: Toluene-4-thiosulfonic acid S-[2-tert-butyl-4-(2-hydroxy-ethyl)-5-methyl-phenyl]ester; and

BB-15: Benzenesulfonylthioic acid, 4-methyl-, S-[2-(1,1-dimethylethyl)-5-methyl-4-[[[(5-trifluoromethyl)-2-pyridinyl]sulfonyl]amino]phenyl]ester;

5 BB-16: 6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-indole-1-carboxylic acid tert-butyl ester;

BB-17: 6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester;

10 BB-18: Toluene-4-thiosulfonic acid S-[6-tert-butyl-1-(4-cyanobenzenesulfonyl)-2,3-dihydro-1H-indol-5-yl] ester;

BB-19: Toluene-4-thiosulfonic acid S-(6-isopropyl-3H-benzimidazol-5-yl) ester;

BB-20: Toluene-4-thiosulfonic acid S-(6-tert-butyl-3H-benzimidazol-5-yl) ester;

15 BB-21: Toluene-4-thiosulfonic acid S-(6-tert-butyl-2-oxo-2,3-dihydro-1H-benzimidazol-5-yl) ester;

BB-22: 6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-1H-indole-2-carboxylic acid ethyl ester;

20 BB-23: Toluene-4-thiosulfonic acid S-(7-tert-butyl-2,3-dioxo-1,2,3,4-tetrahydro-quinoxalin-6-yl) ester;

BB-24: Toluene-4-thiosulfonic acid S-[5-tert-butyl-2-(4-cyanobenzenesulfonyl-amino)-benzothiazol-6-yl]] ester;

BB-25: Toluene-4-thiosulfonic acid S-(6-tert-butyl-2-methanesulfonylamino-benzothiazol-6-yl) ester;

25 BB-26: Toluene-4-thiosulfonic acid S-(2-acetylamino-5-tert-butyl-benzothiazol-6-yl) ester; and

BB-27: Toluene-4-thiosulfonic acid S-(6-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazol-5-yl) ester.

-165-

Example BB-15

Benzenesulfonylthioic acid, 4-methyl-, S-[2-(1,1-dimethylethyl)-5-methyl-4-[[[(5-trifluoromethyl)-2-pyridinyl]sulfonyl]amino]phenyl]ester

Compound BB-11 (0.5 g, 1.43 mmol) was dissolved in pyridine (5 mL) and CH₂Cl₂ (2 mL). To this solution was added 0.53 g (2.16 mmol) of 5-(trifluoromethyl)pyridine-2-sulfonyl chloride (*J. Amer. Chem. Soc.*, 1997;119(15):3627-3628), and the reaction mixture was stirred under N₂ at room temperature overnight. 1N HCl was then added; the mixture was extracted with EtOAc, washed with 1 N HCl and brine, dried (MgSO₄), filtered, and concentrated. The resulting residue was chromatographed over silica gel, eluting with 9:1 CH₂Cl₂:EtOAc to afford the desired product.

¹H NMR (CDCl₃): δ 1.13 (s, 9 H), 2.19 (s, 3 H), 2.42 (s, 3 H), 6.83 (s, 1 H), 7.22 (d, 2 H), 7.40 (s, 1 H) 7.45 (d, 2 H), 8.08 (d, 1 H), 8.15 (m, 1 H), 8.99 (s, 1 H).

15

Example BB-24

Toluene-4-thiosulfonic acid S-[5-tert-butyl-2-(4-cyanobenzenesulfonyl-amino)-benzothiazol-6-yl] ester

The title compound was prepared as described in Example BB-15 using toluene-4-thiosulfonic acid S-(2-amino-5-tert-butyl-benzothiazol-6-yl)ester (prepared in Example BB-1; 1.0 g, 3.37 mmol), 4-cyanobenzenesulfonyl chloride (0.68 g, 3.37 mmol), pyridine (10 mL) and CH₂Cl₂ (3 mL). The crude product was purified by flash silica gel chromatography (20%-70% EtOAc in hexanes as eluents). MS(APCI): 558 (M+H).

20

Example BB-25

25

Toluene-4-thiosulfonic acid S-(6-tert-butyl-2-methanesulfonylamino-benzothiazol-6-yl) ester

The title compound was prepared as described in Example BB-15 using toluene-4-thiosulfonic acid S-(2-amino-5-tert-butyl-benzothiazol-6-yl)ester (prepared in Example BB-1; 1.75 g, 4.47 mmol), methanesulfonyl chloride (1.02 g, 8.94 mmol), pyridine (3 mL), DMAP (0.3 g) and CH₂Cl₂ (20 mL). The

30

-166-

crude product was purified by flash silica gel chromatography (20%-70% EtOAc in hexanes as eluents). MS(APCI): 471 (M+H).

Example BB-26

Toluene-4-thiosulfonic acid S-(2-acetylamino-5-tert-butyl-benzothiazol-6-yl) ester

5 The title compound was prepared as described in Example BB-15 using toluene-4-thiosulfonic acid S-(2-amino-5-tert-butyl-benzothiazol-6-yl)ester (prepared in Example BB-1; 1.75 g, 4.47 mmol), acetyl chloride (4 mL), pyridine (5 mL), DMAP (0.3 g) and CH₂Cl₂ (20 mL). The crude product was purified by
10 flash silica gel chromatography (20%-70% EtOAc in hexanes as eluents). MS(APCI): 435 (M+H).

Example BB-27

Toluene-4-thiosulfonic acid S-(6-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazol-5-yl) ester

15 Toluene-4-thiosulfonic acid S-(1,2-diamino-5-isopropyl-phenyl)ester (prepared in Example VV-1 below; 0.5 g, 1.5 mmol) and triphosgene (0.15 g, 0.5 mmol) were dissolved in THF (30 mL). NEt₃ (1 mL) was added, and the reaction mixture was stirred at 90°C for 1.5 hours. The crude reaction mixture was subjected to flash silica gel chromatography (EtOAc to 2%-10% MeOH in
20 CH₂Cl₂ as eluents). MS(APCI): 363 (M+H).

Example CC

(5-tert-Butyl-2-methyl-phenyl)-carbamic acid tert-butyl ester

5-tert-Butyl-2-methyl-phenylamine (50 g, 307 mmol) in 500 mL of hexanes was treated with di-t-butyl dicarbonate (80.3 g, 368 mmol.). The reaction
25 mixture was kept under reflux overnight. After cooling to room temperature, the reaction mixture was diluted with EtOAc, washed with saturated NaHCO₃ solution and brine, and dried (MgSO₄). After evaporation of the solvents, the crude reaction mixture was purified by flash silica gel chromatography to give the title compound. MS(APCI): 262 (M-H).

-167-

Example DD

(5-tert-Butyl-2-methyl-4-thiocyano-phenyl)-carbamic acid tert-butyl ester

Thiocyanation was performed as described in the General Method 13 using (5-tert-butyl-2-methyl-phenyl)-carbamic acid tert-butyl ester (prepared in Example CC; 4.32 g, 15.82 mmol), sodium thiocyanate (7.7 g, 94.95 mmol), sodium bromide (1.95 g, 18.98 mmol), bromine (3.03 g, 18.98 mmol), and methanol (50 mL). MS(APCI): (M+H).

Example EE

(5-tert-Butyl-4-mercapto-2-methyl-phenyl)-carbamic acid tert-butyl ester

The title compound was prepared according to General Method 14 using (5-tert-butyl-2-methyl-4-thiocyano-phenyl)-carbamic acid tert-butyl ester (prepared in Example DD; 30 g, 93.75 mmol), dithiothreitol (29 g, 187.5 mmol) and phosphate buffer (50 mL). MS(APCI): 294 (M-H).

Example FF

6-tert-Butyl-5-mercapto-indole-1-carboxylic acid tert-butyl ester

(5-tert-Butyl-4-mercapto-2-methyl-phenyl)-carbamic acid tert-butyl ester (prepared in Example EE; 20 g, 68 mmol) was dissolved in 100 mL of THF, cooled to -40°C, and treated dropwise with sec-butyllithium (1.3 M, 156 mL, 203 mmol). After the addition was complete, the reaction mixture was stirred at -30°C to -45°C for 30 minutes. To the mixture was added DMF (14.8 g, 203.39 mmol), and the solution was stirred at -35°C for 30 minutes. The reaction mixture was slowly warmed to room temperature and was quenched with saturated NH₄Cl solution. EtOAc (200 mL) was added; the organic layer was washed with brine, dried (MgSO₄) and concentrated. The residue was dissolved in 100 mL of THF and 2 mL of concentrated HCl and stirred at room temperature for 1 hour. The solution was diluted with EtOAc, washed with NaHCO₃ solution and brine, and dried (MgSO₄). The crude product was purified by flash silica gel chromatography to give the title compound. MS(APCI): 304 (M-H).

-168-

Example BB-16

6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-indole-1-carboxylic acid tert-butyl ester

5 The title compound was prepared according to General Method 15 using 6-tert-butyl-5-mercapto-indole-1-carboxylic acid tert-butyl ester (prepared in Example FF), pyridine, tosyl bromide, carbon tetrachloride, and EtOAc.

Example GG

6-tert-Butyl-indole-1-carboxylic acid tert-butyl ester

10 (5-tert-Butyl-2-methyl-phenyl)-carbamic acid tert-butyl ester (prepared in Example CC; 19.7 g, 75 mmol) in 100 mL of THF was cooled to -40°C. To it was added tert-butyllithium (1.5 M, 100 mL, 150 mmol), dropwise with stirring, at -45°C to -35°C. After the addition was complete, the reaction was stirred at -35°C for another 15 minutes. DMF (10.97 g, 150 mmol) was added dropwise, and the reaction mixture was stirred at -35°C for 30 minutes, followed by slow warming
15 to room temperature. The reaction was quenched with saturated NH₄Cl solution and diluted with EtOAc. The organic layer was washed with brine, dried (MgSO₄), and concentrated. The residue was dissolved in 100 mL of THF and 2 mL of concentrated HCl and stirred at room temperature for 1 hour. EtOAc was added, and the solution was washed with saturated NaHCO₃ solution and brine
20 and dried (MgSO₄). Concentration gave a residue that was purified by flash silica gel chromatography (5%-50% EtOAc in hexanes). MS(APCI): 274 (M+H).

Example HH

6-tert-Butyl-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester

25 6-tert-Butyl indole-1-carboxylic acid tert-butyl ester (prepared in Example GG; 13.0 g, 48 mmol) was taken up in 500 mL of AcOH. 20% Pd/C (2 g) was added, and the mixture was shaken under a hydrogen atmosphere of 50 psi. The catalyst was filtered, and the filtrate was concentrated. The residue was dissolved in EtOAc, washed with saturated NaHCO₃ solution and brine, and dried (MgSO₄). Concentration gave the title compound which was used without
30 purification in the next step.

-169-

Example II

6-tert-Butyl-2,3-dihydro-1H-indole

6-tert-Butyl-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester

(Prepared in Example HH; 12.0 g) in CH₂Cl₂ (100 mL) was cooled to 0°C and treated with HCl gas for 10 minutes. The reaction mixture was warmed to room temperature and stirred for 1 hour. The solvents were evaporated; the residue was dissolved in EtOAc, washed with saturated NaHCO₃ solution and brine, and dried (MgSO₄.) Concentration gave the title compound which was used without any further purification. MS(APCI): 176 (M+H).

Example JJ

6-tert-Butyl-5-thiocyanato-2,3-dihydro-1H-indole

The title compound was prepared according to General Method 13 using 6-tert-butyl-2,3-dihydro-1H-indole (prepared in Example II; 8.3 g, 47 mmol), MeOH (100 mL), sodium thiocyanate (23.1 g, 284.57 mmol), NaBr (5.86 g, 56.92 mmol) and bromine (9.1 g, 56.92 mmol). The crude reaction mixture was purified by flash silica gel chromatography (5%-75% EtOAc in hexanes). MS(APCI): 233 (M+H).

Example KK

6-tert-Butyl-5-thiocyanato-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester

The title compound was prepared in the manner described in Example CC using 6-tert-butyl-5-thiocyanato-2,3-dihydro-1H-indole (prepared in Example JJ; 2.0 g, 8.6 mmol), di-t-butyl dicarbonate (2.35 g, 10.8 mmol) and hexanes (150 mL). The crude product was purified by flash silica gel chromatography.

Example LL

6-tert-Butyl-5-mercapto-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester

The title compound was prepared according to General Method 14 using 6-tert-butyl-5-thiocyanato-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester (prepared in Example KK; 2.0 g, 6.6 mmol), dithiothreitol (1.21 g, 7.9 mmol),

-170-

EtOH (100 mL), and phosphate buffer (pH: 7.5, 10 mL). The crude product was used without any purification.

Example BB-17

5 **6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester**

The title compound was prepared according to General Method 15 using 6-tert-Butyl-5-mercapto-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester (prepared in Example FF), pyridine, tosyl bromide, carbon tetrachloride, and EtOAc.

10

Example MM

4-(6-tert-Butyl-5-thiocyanato-2,3-dihydro-indole-1-sulfonyl)-benzonitrile

6-tert-Butyl-5-thiocyanato-2,3-dihydro-1H-indole (prepared in Example JJ; 4.5 g, 19 mmol) was taken in CH₂Cl₂ (50 mL) and treated with 4-cyanophenylsulfonyl chloride (3.91 g, 19.4 mmol) and pyridine (5 mL). The reaction mixture was stirred overnight, quenched with saturated NaHCO₃ solution, and diluted with EtOAc. The organic layer was separated, dried (MgSO₄), and concentrated. The crude product was purified by flash chromatography (20%-75% EtOAc in hexanes as eluents). MS(APCI): 397 (M⁺).

15

Example NN

20 **4-(6-tert-Butyl-5-mercapto-2,3-dihydro-indole-1-sulfonyl)-benzonitrile**

The title compound was prepared according to the General Method 14b using 4-(6-tert-butyl-5-thiocyanato-2,3-dihydro-indole-1-sulfonyl)-benzonitrile (prepared in Example MM; 6.0 g, 15 mmol), sodium hydrogen sulfide (2.53 g, 45.3 mmol), sodium borohydride (3.43 g, 90.7 mmol), MeOH (10 mL), water (5 mL), and AcOH (5 mL). The crude product was used further without any purification.

25

-171-

Example OO-1

2,2,2-Trifluoro-N-(5-isopropyl-2-nitro-phenyl)-acetamide

Trifluoroacetic anhydride (50 mL) was cooled to 0°C and treated dropwise with 3-isopropyl-phenylamine (5 g, 37.03 mmol). To the resulting solution was added potassium nitrate (4.12 g, 101.11 mmol). After 1 hour, the reaction mixture was warmed to room temperature and stirred overnight. The reaction mixture was diluted carefully with ice water and extracted with EtOAc (4 × 100 mL). The organic layer was washed with saturated NaHCO₃ solution and brine, dried (MgSO₄), and concentrated. The crude product was used further without any purification. MS(APCI): 275 (M-H).

Example OO-2

N-(5-tert-Butyl-2-nitro-phenyl)-2,2,2-trifluoro-acetamide

The title compound was prepared following the procedure used in Example OO-1 using trifluoroacetic anhydride (300 mL), 3-tert-butyl-phenylamine (144 g, 951 mmol), and potassium nitrate (106 g, 1047 mmol). The crude product was used further without any purification. MS(APCI): 289 (M-H).

Example PP-1

5-Isopropyl-2-nitro-phenylamine

2,2,2-Trifluoro-N-(5-isopropyl-2-nitro-phenyl)-acetamide (prepared in Example OO-1; 9.4 g, 34 mmol) was dissolved in MeOH (30 mL) and treated with 7% aqueous potassium carbonate solution. The reaction mixture was stirred overnight at room temperature. The reaction was diluted with water (30 mL) and extracted with EtOAc (3 × 50 mL). The organic layer was washed with saturated NaHCO₃ solution and brine, dried (MgSO₄), and concentrated. The crude product was used further without any purification. MS(APCI): 181 (M+H).

Example PP-2

5-tert-Butyl-2-nitro-phenylamine

The title compound was prepared as outlined in Example PP-1 from N-(5-tert-butyl-2-nitro-phenyl)-2,2,2-trifluoro-acetamide (prepared in Example OO-2;

-172-

248 g, 855 mmol), MeOH (500 mL), and 7% aqueous potassium carbonate solution. The crude product was used without any purification. MS(APCI): 195 (M+H).

Example QQ-1

5 **5-Isopropyl-2-nitro-4-thiocyanato-phenylamine**

The title compound was prepared according to General Method 13 using 5-isopropyl-2-nitro-phenylamine (prepared in Example PP-1; 5.4 g, 30 mmol), sodium thiocyanate (14.6 g, 180 mmol), sodium bromide (3.7 g, 35.99 mmol), bromine (5.75 g, 36.0 mmol), and MeOH (100 mL). The crude reaction mixture was purified by flash silica gel chromatography (5%-50% EtOAc in hexanes as eluents). MS(APCI): 238 (M+H).

Example QQ-2

5-tert-Butyl-2-nitro-4-thiocyanato-phenylamine

The title compound was prepared according to the General Method 13 using 5-tert-butyl-2-nitro-phenylamine (prepared in Example PP-2; 170 g, 876 mmol), ammonium thiocyanate (400 g, 5260 mmol), bromine (140 g, 876 mmol), and MeOH (500 mL). The crude reaction mixture was purified by flash silica gel chromatography (5%-50% EtOAc in hexanes as eluents). MS(APCI): 252 (M+H).

20 **Example RR-1**

4-Isopropyl-5-thiocyanato-benzene-1,2-diamine

5-Isopropyl-2-nitro-4-thiocyanato-phenylamine (prepared in Example QQ-1; 7.83 g, 33 mmol) was hydrogenated using Raney nickel (2 g) and THF (100 mL) in the presence of hydrogen (50 psi) as described in General Method 3. MS(APCI): 208(M+H).

-173-

Example RR-2

4-tert-Butyl-5-thiocyanato-benzene-1,2-diamine

5 5-tert-Butyl-2-nitro-4-thiocyanato-phenylamine (prepared in Example QQ-2; 3 g, 12 mmol) was hydrogenated using Raney nickel (0.75 g) in the presence of hydrogen as described in General Method 3. MS(APCI): 222(M+H).

Example SS-1

(2-tert-Butyloxycarbonylamino-4-isopropyl-5-thiocyanato-phenyl)-carbamic acid tert-butyl ester

10 Boc protection was performed as described in Example CC using 4-isopropyl-5-thiocyanato-benzene-1,2-diamine (prepared in Example RR-1; 4.49 g, 21.7 mmol), di-t-butyl dicarbonate (11.3 g, 52.0 mmol), hexanes (150 mL), and EtOAc (50 mL). The crude product was purified by flash silica gel chromatography. MS(APCI): 406 (M-H).

Example SS-2

15 **(2-tert-Butoxycarbonylamino-4-tert-Butyl-5-thiocyanato-phenyl)-carbamic acid tert-butyl ester**

20 Boc protection was performed as described in Example CC using 4-tert-butyl-5-thiocyanato-benzene-1,2-diamine (prepared in Example RR-2; 9.78 g, 44.2 mmol), di-t-butyl dicarbonate (23.2 g, 106 mmol), hexanes (150 mL), and EtOAc (50 mL). The crude product was purified by flash silica gel chromatography. MS(APCI): 422 (M+H).

Example TT-1

(2-tert-Butyloxycarbonylamino-4-isopropyl-5-mercapto-phenyl)-carbamic acid tert-butyl ester

25 The title compound was prepared according to General Method 14b using (2-tert-butyloxycarbonylamino-4-isopropyl-5-thiocyanato-phenyl)-carbamic acid tert-butyl ester (prepared in Example SS-1; 9 g, 22 mmol), sodium hydrogen sulfide (3.7 g, 66 mmol), sodium borohydride (5.02 g, 132 mmol), MeOH (20 mL), H₂O (5 mL), and AcOH (2 mL). The crude product was used without
30 any purification. MS(APCI): 381 (M-H).

-174-

Example TT-2

(2-tert-Butoxycarbonylamino-4-tert-Butyl-5-mercapto-phenyl)-carbamic acid tert-butyl ester

5 The title compound was prepared according to General Method 14b using (2-tert-butoxycarbonylamino-4-tert-butyl-5-thiocyanato-phenyl)-carbamic acid tert-butyl ester (prepared in Example SS-2; 18 g, 43 mmol), sodium hydrogen sulfide (7.2 g, 128 mmol), sodium borohydride (9.7 g, 256 mmol), MeOH (150 mL), H₂O (50 mL), and AcOH (10 mL). The crude product was used without purification.

10

Example UU-1

Toluene-4-thiosulfonic acid S-(4,5-bis-tert-butoxycarbonylamino-2-isopropyl-phenyl) ester

15 The title compound was prepared according to the General Method 15 using (2-tert-butoxycarbonylamino-4-isopropyl-5-mercapto-phenyl)-carbamic acid tert-butyl ester (prepared in Example TT-1; 22.1 mmol), tosyl bromide (5.2 g, 22 mmol), pyridine (5 mL), and EtOAc (50 mL). The crude product was purified by flash chromatography (5%-60% EtOAc in hexanes as eluents). ¹H-NMR (CDCl₃): δ 0.97 (s, 6 H), 1.53 (s, 18 H), 2.4 (s, 3 H), 3.2 (m, 1 H), 7.11 (br s, 1 H), 7.22 (d, 2 H), 7.28 (s, 1 H), 7.36 (s, 1 H), 7.5 (d, 2 H), 7.75 (br s, 1 H).

20

Example UU-2

Toluene-4-thiosulfonic acid S-(4,5-bis-tert-butoxycarbonylamino-2-tert-butyl-phenyl) ester

25 The title compound was prepared according to the General Method 15 using (2-tert-butoxycarbonylamino-4-tert-butyl-5-mercapto-phenyl)-carbamic acid tert-butyl ester (prepared in Example TT-2; 16.9 g, 42.8 mmol), tosyl bromide (10.04 g, 42.76 mmol), pyridine (10 mL), and EtOAc (150 mL). The crude product was purified by flash chromatography (5%-60% EtOAc in hexanes as eluents). ¹H-NMR (CDCl₃): δ 1.19 (s, 9 H), 1.53 (s, 18 H), 2.42 (s, 3 H), 6.42 (br s, 1 H), 7.22 (d, 2 H), 7.28 (s, 1 H), 7.5 (m, 3 H), 7.87 (br s, 1 H).

-175-

Example VV-1

Toluene-4-thiosulfonic acid S-(4,5-diamino-2-isopropyl-phenyl) ester

Toluene-4-thiosulfonic acid S-(4,5-bis-tert-butoxycarbonylamino-2-isopropyl-phenyl) ester (prepared in Example UU-1; 8.88 g, 16.6 mmol) was dissolved in CH₂Cl₂ (25 mL), and HCl gas was bubbled through the solution for 15 minutes. The reaction mixture was stirred at room temperature for 90 minutes. The solvents were evaporated, and the residue was dissolved in MeOH (10 mL) and treated with pH 7.5 buffer. The precipitate that formed was filtered and dried. MS(APCI): 337 (M+H).

10

Example VV-2

Toluene-4-thiosulfonic acid S-(4,5-diamino-2-tert-butyl-phenyl) ester

The title compound was prepared as described in Example VV-1 from toluene-4-thiosulfonic acid S-(4,5-bis-tert-butoxycarbonylamino-2-tert-butyl-phenyl) ester (prepared in Example UU-2; 15 g, 28 mmol), CH₂Cl₂ (100 mL), and HCl gas. MS(APCI): 351 (M+H).

15

Example BB-19

Toluene-4-thiosulfonic acid S-(6-isopropyl-3H-benzoimidazol-5-yl)

Toluene-4-thiosulfonic acid S-(1,2-diamino-5-isopropyl-phenyl)ester (prepared in Example VV-1; 0.5 g) was dissolved in 96% formic acid (0.5 mL) and refluxed for 3 hours. After cooling, toluene (10 mL) and water (5 mL) were added. The organic layer was concentrated; the residue was taken in EtOAc (20 mL), washed with H₂O, and dried (MgSO₄). Concentration gave the crude title compound which was purified by flash chromatography (20%-100% EtOAc in hexanes to 2%-8% MeOH in CH₂Cl₂ as eluents). MS(APCI): 347 (M+H).

20

25

Example BB-20

Toluene-4-thiosulfonic acid S-(6-tert-butyl-3H-benzoimidazol-5-yl) ester

The title compound was prepared using the procedure outlined in Example BB-19 from toluene-4-thiosulfonic acid S-(1,2-diamino-5-tert-butyl-phenyl)ester (prepared in Example VV-2; 0.5 g), and 96% formic acid (0.5 mL). The crude

-176-

product was purified by flash chromatography (20%-100% EtOAc in hexanes to 2%-8% MeOH in CH₂Cl₂ as eluents). MS(APCI): 361 (M+H).

Example BB-21

Toluene-4-thiosulfonic acid S-(6-tert-butyl-2-oxo-2,3-dihydro-1H-benzimidazol-5-yl) ester

5

Toluene-4-thiosulfonic acid S-(4,5-diamino-2-tert-butyl-phenyl)ester (prepared in Example VV-2; 0.4 g, 1.2 mmol) and triphosgene (0.12 g, 0.41 mmol) were taken in THF (30 mL). To it triethylamine (1 mL) was added. The reaction mixture was kept at 90°C for 1.5 hours. The crude reaction mixture was subjected to flash silica gel chromatography (EtOAc to 2%-10% MeOH in CH₂Cl₂ as eluents). MS(APCI): 375 (M-H).

10

Preparation of 2-substitued-Indoles for the Synthesis of BB-22

Example WW

4-tert-Butyl-1-methyl-2-nitrobenzene

15

Nitration was performed as described in General Method 12 using tert-butyl-4-methylbenzene (25 g), concentrated H₂SO₄ (92 g), 70% HNO₃ (82 g), and H₂O (25 g). The crude product was purified via distillation under vacuum.

Example XX

3-(4-tert-Butyl-2-nitro-phenyl)-2-oxo-propionic acid ethyl ester

20

The title compound was prepared by adapting the procedure described by Gagliardi S., *et al.*, in *J. Med. Chem.*, 1998;41:1568 using potassium (1.66 g, 42.4 mmol), Et₂O (200 mL), EtOH (10 mL), diethyl oxalate (7.56 g, 51.8 mmol), and 4-tert-butyl-1-methyl-2-nitrobenzene (prepared in Example WW; 10.0 g, 51.8 mmol). MS(APCI): 294 (M+H).

-177-

Example YY

6-tert-Butyl-1H-indole-2-carboxylic acid ethyl ester

A mixture of 3-(4-tert-butyl-2-nitro-phenyl)-2-oxo-propionic acid ethyl ester (prepared in Example XX; 9.1 g, 31 mmol), iron powder (15.3 g, 274 mmol),
5 EtOH (70 mL), and glacial AcOH (70 mL) was refluxed for 2 hours. After cooling, the resulting mixture was evaporated. THF (100 mL) was added to the residue, and the suspension was filtered on florisil, eluting with a large amount of THF. The solvents were evaporated, and the residue was purified by flash silica gel chromatography (5%-30% EtOAc in hexanes as eluents). MS(APCI):
10 246 (M+H).

Example ZZ

6-tert-Butyl-5-thiocyanato-1H-indole-2-carboxylate ethyl ester

The title compound was prepared according to General Method 13 using 6-tert-butyl-1H-indole-2-carboxylic acid ethyl ester (prepared in Example YY;
15 3.5 g, 14 mmol), ammonium thiocyanate (1.63 g, 21.4 mmol), bromine (3.42 g, 21.4 mmol), and MeOH (50 mL). The crude product was purified by flash silica gel chromatography (5%-50% EtOAc in hexanes as eluents). MS(APCI): 301 (M-H).

Example AAA

20 (6-tert-Butyl-5-mercapto-1H-indole-2-carboxylic acid ethyl ester

The title compound was prepared according to the general Method 14b using (6-tert-butyl-5-thiocyanato-1H-indole-2-carboxylic acid ethyl ester (prepared in Example ZZ; 0.25 g, 0.83 mmol), sodium hydrogen sulfide (0.13 g, 2.48 mmol), sodium borohydride (0.188 g, 4.98 mmol), MeOH (5 mL), and
25 AcOH (1 mL). MS(APCI): 278 (M+H).

-178-

Example BB-22

6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-1H-indole-2-carboxylic acid ethyl ester

5 The title compound was prepared according to General Method 15 using (6-tert-Butyl-5-mercapto-1H-indole-2-carboxylic acid ethyl ester (prepared in Example AAA), pyridine, tosyl bromide, carbon tetrachloride, and EtOAc.

Preparation of Quinoxalones for the Synthesis of BB-23

Example BBB

(5-tert-Butyl-2-nitro-4-thiocyanatophenyl)-oxalamic acid methyl ester

10 5-tert-Butyl-2-nitro-4-thiocyanato-phenylamine (prepared in Example QQ-2; 2.5 g, 10.0 mmol) and methyl chloroacetate (5 mL) were dissolved in CH₂Cl₂ (50 mL) and treated with pyridine (5 mL.) The reaction mixture was stirred at room temperature overnight, quenched with saturated NaHCO₃ solution, and diluted with EtOAc (100 mL). The organic layer was washed with dil HCl
15 and brine, dried (MgSO₄), and concentrated. The crude product was used without purification. MS(APCI): 336 (M-H).

Example CCC

6-tert-Butyl-7-thiocyanato-1,4-dihydro-quinoxaline-2,3-dione

20 (5-tert-Butyl-2-nitro-4-thiocyanatophenyl)-oxalamic acid methyl ester (prepared in Example BBB; 3.91 g, 11.6 mmol) was taken up in THF (100 mL) and treated with Raney nickel (1 g). The reaction mixture was shaken in a hydrogen atmosphere (50 psi). The catalyst was filtered; the filtrate was concentrated, and the residue was triturated with EtOAc/hexanes to obtain the title compound. MS(APCI): 274 (M-H).

25

Example DDD

6-tert-Butyl-7-mercapto-1,4-dihydro-quinoxaline-2,3-dione

The title compound was prepared according to General Method 14b using 6-tert-butyl-7-thiocyanato-1,4-dihydro-quinoxaline-2,3-dione (prepared in Example CCC; 1.5 g, 9.1 mmol), sodium hydrogen sulfide (0.51 g, 27 mmol),

-179-

sodium borohydride (2.1 g, 55 mmol), MeOH (30 mL), H₂O (3 mL), and glacial AcOH (5 mL). The crude product was used without purification. MS(APCI): 251 (M+H).

Example BB-23

5 **Toluene-4-thiosulfonic acid S-(7-tert-butyl-2,3-dioxo-1,2,3,4-tetrahydro-
quinoxalin-6-yl) ester**

 The title compound was prepared according to the General Method 15
using 6-tert-butyl-7-mercapto-1,4-dihydro-quinoxaline-2,3-dione (prepared in
Example DDD, 9.1 mmol), tosyl bromide (2.1 g, 9.1 mmol), pyridine (2 mL),
10 EtOAc (50 mL). The crude product was purified by flash silica gel
chromatography (5%-10% MeOH in CH₂Cl₂) as eluents. MS(APCI): 405 (M+H).

General Method 16a. Coupling of the Dihydropyrone and the Thiotosylate

 The appropriate dihydropyrone intermediate (1 equiv.) from Table F or
Table J was added to a reaction flask followed by DMF (1-12 mL per mmol of
15 dihydropyrone). Potassium carbonate (K₂CO₃) (4-8 equiv.) was added followed
by the appropriate thiotosylate reagent (1.1-1.5 eq.) from Table K. The reaction
was stirred at room temperature (2.5 hours to overnight). The reaction was worked
up by pouring into a mixture of EtOAc and either 1N HCl or saturated aqueous
NH₄Cl. The layers were separated and the aqueous layer extracted again with
20 EtOAc. The combined organic extracts were washed with brine, dried (MgSO₄),
and concentrated. The final compounds were purified via recrystallization,
trituration, or silica gel chromatography.

General Method 16b. Coupling of the Dihydropyrone and the Thiotosylate

 The appropriate dihydropyrone intermediate (1 equiv.) from Examples E-1
25 to E-46 or Examples J-1 to J-12 and the appropriate thiotosylate reagent
(1.0-1.5 equiv.) from Examples BB-1 to BB-14 were added to a reaction flask,
followed by acetonitrile (1-12 mL per mmol of dihydropyrone) and triethylamine
(NEt₃; 2-4 equiv.) The reaction was stirred at room temperature (2.5 hours to
overnight.) The NEt₃ and solvent were evaporated, and the reaction was poured

-180-

into a mixture of EtOAc and either saturated aqueous NH₄Cl or water. The aqueous layer was extracted with EtOAc; the combined organic extracts were dried (MgSO₄), and concentrated. The final compounds were purified via recrystallization, trituration, or silica gel chromatography.

5

Example 1

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-4-yl-ethyl)-5,6-dihydro-pyran-2-one

10

The title compound was prepared as described in General Method 16a using 4-hydroxy-6-isopropyl-6-(2-pyridin-4-yl-ethyl)-5,6-dihydro-pyran-2-one (Example E-15; 0.8 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 0.9 mmol), potassium carbonate (3.1 mmol) in DMF (5 mL). The product was trituated from Et₂O, mp 140-155°C. MS (APCI): 470 (M+H).

Example 2

15

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one

20

The title compound was prepared as described in General Method 16a using 4-hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one (Example E-16; 0.92 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 1.01 mmol), potassium carbonate (3.67 mmol) in DMF (5 mL). The product was trituated from Et₂O, mp 122-125°C. MS (APCI): 470 (M+H).

Example 3

25

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-2-yl-ethyl)-5,6-dihydro-pyran-2-one

The title compound was prepared as described in General Method 16a using 4-hydroxy-6-isopropyl-6-(2-pyridin-2-yl-ethyl)-5,6-dihydro-pyran-2-one (Example E-14; 0.8 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 0.9 mmol), potassium

-181-

carbonate (3.1 mmol) in DMF (5 mL). The product was triturated from Et₂O, mp 100-110°C. MS (APCI): 470 (M+H).

Example 4

5 **3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one**

The title compound was prepared as described in General Method 16a using 6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (Example E-18; 1 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 1.1 mmol), potassium carbonate (4 mmol) in DMF (4 mL). The product was flash chromatographed over silica gel using 70:20:10 hexane:EtOAc:CH₂Cl₂, mp 62-70°C. MS (APCI): 457 (M-H).

Example 5

15 **3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-2-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one**

The title compound was prepared as described in General Method 16a using 6-(2-furan-2-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (Example E-17; 1 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 1.1 mmol), potassium carbonate (4 mmol) in DMF (5 mL), mp 64-110°C. MS (APCI): 457 (M-H).

Example 6

25 **3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one**

The title compound was prepared as described in General Method 16a using 4-hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one (Example E-19; 0.98 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-; 1.18 mmol), potassium carbonate (3.93 mmol) in DMF (5 mL). Flash chromatographed over silica gel using 10:90 EtOAc:CH₂Cl₂, mp 57-59°C. MS (APCI): 461 (M-H).

Example 7

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one

5 The title compound was prepared as described in General Method 16a using 4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one (Example E-20; 0.84 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 0.92 mmol), potassium carbonate (3.35 mmol) in DMF (4 mL). The product was purified via flash chromatography over silica gel eluting with 10:90 EtOAc:CH₂Cl₂, mp 63-70°C.

10 MS (APCI): 475 (M+H).

Example 8

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one

15 The title compound was prepared as described in General Method 16a using 4-hydroxy-6-isopropyl-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one (Example E-21; 1 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 1.2 mmol), potassium carbonate (4 mmol) in DMF (3 mL). The product was isolated via flash chromatography over silica gel using 10:90 EtOAc:CH₂Cl₂, mp 73-77°C.

20 MS (APCI): 475 (M+H).

Example 9

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one

25 The title compound was prepared as described in General Method 16a using 4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one (Example E-23; 0.67 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 0.74 mmol), potassium carbonate (2.7 mmol) in DMF (5 mL), mp 76-80°C.

30 MS (APCI): 503 (M-H).

Example 10

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one

5 The title compound was prepared as described in General Method 16b using 4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one (Example E-24; 1 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 1.05 mmol), NEt₃ (2.2 mmol) in acetonitrile (5 mL). The product was
10 chromatographed over silica gel using 97.5:2.5 CH₂Cl₂:MeOH, mp 87-90°C. MS (APCI): 503 (M-H).

Example 11

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one

15 The title compound was prepared as described in General Method 16b using 4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one (Example E-24; 1 mmol), toluene-4-thiosulfonic acid S-(4-amino-2-tert-butyl-5-methyl-phenyl)ester (Example BB-11; 1.05 mmol), NEt₃ (2.2 mmol) in acetonitrile (5 mL). The product was triturated from Et₂O,
20 mp 107-115°C. MS (APCI): 488 (M-H).

Example 12 (S)

(S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one

25 The title compound was prepared as described in General Method 16a using (S)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one (Example J-1 (S); 2.68 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 3.22 mmol), potassium carbonate (10.74 mmol) in DMF (5 mL), mp 64-70°C. MS (APCI): 475 (M+H).

Example 13

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one

The title compound was prepared according to the General Method 16a using toluene-4-thiosulfonic acid S-(2-amino-5-tert-butyl-benzothiazol-6-yl)ester (Example BB-1; 0.6 g, 1.53 mmol), (S)-4-hydroxy-6-isopropyl-6-phenylethyl-5,6-dihydro-pyran-2-one (Example J-11 (S); 0.4 g, 1.53 mmol), potassium carbonate (0.8 g), and DMF (DMF; 5 mL). The crude reaction mixture was purified by flash chromatography (25% EtOAc in hexanes to 50% EtOAc in hexanes to 10% MeOH in EtOAc as eluents), mp 186-188°C. MS (APCI): 497 (M+H).

Example 14

(S)-3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one

The title compound was prepared according to the General Method 16b using (S)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one (Example J-11 (S); 0.5 g, 1.92 mmol), toluene-4-thiosulfonic acid S-(2-amino-5-isopropyl-benzothiazol-6-yl)ester (Example BB-3; 0.73 g, 1.92 mmol), NEt₃ (0.39 g, 3.84 mmol), and acetonitrile (10 mL). The crude compound was purified by flash silica gel chromatography (25% EtOAc in hexanes to 100% EtOAc to 5% MeOH in EtOAc as eluents), mp 197-199°C. MS (APCI): 483 (M+H).

Example 15

6-[-2-(4-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one

The title compound was prepared according to General Method 16b using 6-[(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydropyran-2-one (Example E-27; 0.5 g, 1.8 mmol), toluene-4-thiosulfonic acid S-(5-isopropyl-benzothiazol-6-yl)ester (Example BB-4; 0.65 g, 1.8 mmol), NEt₃ (0.22 g, 2.16 mmol), and acetonitrile (10 mL). The crude compound was purified by flash

silica gel chromatography (25% EtOAc in hexanes to 100% EtOAc to 5% MeOH in EtOAc as eluents), mp 104-105°C. MS (APCI): 486.

Example 16

5 **3-(2-Amino-5-isopropyl-benzothiazol-6-yl-sulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-phenethyl-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one**

10 The title compound was prepared according to General Method 16b using 6-[(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (Example E-27; 0.5 g, 1.8 mmol), toluene-4-thiosulfonic acid S-(2-amino-5-isopropyl-benzothiazol-6-yl)ester (Example BB-3; 0.68 g, 1.8 mmol), NEt₃ (0.36 g, 3.6 mmol), and acetonitrile (20 mL). The crude compound was purified by flash silica gel chromatography (25% EtOAc in hexanes to 100% EtOAc to 5% MeOH in EtOAc as eluents), mp 204-206°C. MS (APCI): 501 (M+H).

Example 17

15 **(S)-3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-yl-sulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one**

20 The title compound was prepared according to General Method 16a using toluene-4-thiosulfonic acid S-(2-amino-7-isopropyl-4-methyl-benzothiazol-6-yl)ester (Example BB-5; 0.75 g, 1.92 mmol), (S)-4-hydroxy-6-isopropyl-6-phenylethyl-5,6-dihydro-pyran-2-one (Example J-11 (S); 0.5 g, 1.92 mmol), potassium carbonate (1.0 g), and DMF (5 mL). The crude reaction mixture was purified by flash chromatography (25% EtOAc in hexanes to 50% EtOAc in hexanes to 5% MeOH in EtOAc as eluents), mp 160-162°C. MS (APCI): 497 (M+H).

25

Example 18

30 **(S)-3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one**

30 The title compound was prepared according to General Method 16a using toluene-4-thiosulfonic acid S-(2-amino-7-isopropyl-4-methyl-benzothiazol-6-yl)ester (Example BB-5; 0.35 g, 1.27 mmol), (S)-4-hydroxy-6-[2-(4-hydroxy-

phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one (Example J-10; 0.35 g, 1.27 mmol), potassium carbonate (0.7 g), and DMF (5 mL). The crude reaction mixture was purified by flash chromatography (20% EtOAc in hexanes to 50% EtOAc in hexanes to 5% MeOH in EtOAc as eluents), mp 232-234°C.

5 MS (APCI): 513 (M+H).

Example 19

(S)-(6-{4-Hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-isopropyl-7-methyl-benzothiazol-2-yl)-carbamic acid methyl ester

10 The title compound was prepared according to General Method 16a using toluene-4-thiosulfonic acid S-[2-(methyloxycarbonyl)amino]-4-isopropyl-7-methyl-benzothiazol-6-yl)ester (Example BB-6; 0.65 g, 1.5 mmol), (S)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one (Example J-10 (S); 0.4 g, 1.5 mmol), potassium carbonate (0.7 g), and DMF
15 (5 mL). The crude reaction mixture was purified by flash chromatography (30% EtOAc in hexanes to 100% EtOAc as eluents), mp 150-151°C. MS (APCI): 571 (M+H).

Example 20

(S)-3-(2-Amino-6-tert-butyl-benzothiazol-4-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one

20 The title compound was prepared according to General Method 16a using toluene-4-thiosulfonic acid S-(2-amino-6-tert-butyl-benzothiazol-4-yl)ester (Example BB-7; 0.28 g, 0.73 mmol), (S)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one (Example J-10 (S); 0.2 g, 0.73 mmol),
25 potassium carbonate (0.7 g), and DMF (5 mL). The crude reaction mixture was purified by flash chromatography (30% EtOAc in hexanes to 100% EtOAc to 10% MeOH in CH₂Cl₂ in as eluents), mp 168-179°C. MS (APCI): 513 (M+H).

Example 21

(S)-3-(3-tert-Butyl-benzo[b]thiophen-2-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one

30

-187-

The title compound was prepared according to General Method 16a using (S)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one (Example J-10 (S); 0.052 g, 0.19 mmol), toluene-4-thiosulfonic acid S-(3-tert-butyl-benzo[b]thiophen-2-yl) ester (Example BB-8; 0.080 g, 0.21 mmol),
5 potassium carbonate (0.029 g, 0.21 mmol), and DMF (1.0 mL). The product was purified via column chromatography (eluting with 1:1 hexanes:EtOAc) to provide a solid, mp 207°C.

¹H NMR (DMSO-d₆): δ 0.86-0.92 (m, 6 H), 1.56 (s, 9 H), 1.85-1.95 (m, 2 H),
2.09-2.21 (m, 3 H), 2.52 (d, 1 H), 2.92 (d, 1 H), 6.62 (d, 2 H), 6.93 (d, 2 H),
10 7.12 (t, 1 H), 7.21 (t, 1 H), 7.40 (d, 1 H), 7.89 (d, 1 H), 9.14 (s, 1 H).

Example 22

3-(3-tert-Butyl-benzo[b]thiophen-2-yl-sulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one

The title compound was prepared according to General Method 16a using
15 4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one (Example E-34; 0.20 g, 0.73 mmol), toluene-4-thiosulfonic acid S-(3-tert-butyl-benzo[b]thiophen-2-yl)ester (Example BB-8; 0.302 g, 0.803 mmol), potassium carbonate (0.11 g, 0.80 mmol), and DMF (2.0 mL). The product was purified via column chromatography (eluting with EtOAc) to the title compound,
20 mp 92-94°C.

¹H NMR (DMSO-d₆): δ 0.93-0.99 (m, 6 H), 1.61 (s, 9 H), 2.13-2.28 (m, 3 H),
2.47-2.71 (m, 3 H), 2.99 (d, 1 H), 4.50 (dd, 2 H), 7.09-7.28 (m, 5 H), 7.35 (d, 1 H), 7.42 (d, 1 H), 7.93 (d, 1 H).

Example 23

4-Hydroxy-3-(5-hydroxymethyl-2-isopropyl-thiophen-3-ylsulfanyl)-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one

The title compound was prepared according to General Method 16b using
25 4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one (Example E-20; 0.200 g, 0.75 mmol), toluene-4-thiosulfonic acid S-[5-(tert-butyl-dimethyl-silyloxymethyl)-2-isopropyl-thiophen-3-yl]ester (Example BB-9;
30

-188-

0.377 g, 0.83 mmol), NEt₃ (0.209 mL, 1.50 mmol), and acetonitrile (1.5 mL). As part of the purification, the silylether was cleaved by dissolving the crude coupled product in THF (50 mL) and then treated with a 1.0 M THF solution of tetrabutylammoniumfluoride (1.5 mL, 1.5 mmol). After 30 minutes, the mixture was diluted with H₂O and extracted with EtOAc. The organic layers were then combined, dried (MgSO₄), and concentrated. The resulting residue was then submitted to column chromatography (eluting with 3% MeOH in CH₂Cl₂) to provide the title compound, mp 52-54°C.

¹H NMR (CDCl₃): δ 0.96-1.03 (m, 6 H), 1.32 (t, 6 H), 1.92-2.09 (m, 2 H), 2.17-2.26 (m, 1 H), 2.49 (d, 1 H), 2.66 (t, 2 H), 2.93 (d, 1 H), 3.71 (sept., 1 H), 4.57 (s, 2 H), 6.71 (s, 1 H), 6.79 (d, 1 H), 6.84 (bs, 1 H), 7.23-7.24 (m, 1 H), 7.83 (s, 1 H).

Example 24

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one

The title compound was prepared according to General Method 16a using 4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one (Example E-1; 0.22 g, 0.78 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl) ester (Example BB-2; 0.33 g, 0.90 mmol), potassium carbonate (1.0 g, 7.2 mmol), and DMF (4 mL). The product was chromatographed on silica gel, eluting with 10% MeOH in CH₂Cl₂, to give the title compound, mp 138-141°C.

¹H NMR (DMSO-d₆): δ 0.90-0.96 (m, 6 H), 1.46 (s, 9 H), 1.85 (s, 3 H), 1.95-2.00 (m, 2 H), 2.15-2.23 (m, 4 H), 2.70-2.82 (m, 3 H), 2.93 (d of ABX q, 1 H), 4.34 (s, 2 H), 4.92 (br s, 1 H), 6.66 (s, 1 H), 7.24 (s, 1 H), 8.79 (s, 1 H).

-189-

Example 25

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-2-yl-ethyl)-5,6-dihydro-pyran-2-one

The title compound was prepared according to General Method 16a using
5 4-hydroxy-6-isopropyl-6-(2-thiazol-2-yl-ethyl)-5,6-dihydro-pyran-2-one
(Example E-2; 0.25 g, 0.95 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 0.42 g, 1.15 mmol), potassium carbonate (0.52 g, 3.76 mmol), and DMF (5 mL). The product was chromatographed on silica gel, eluting with 5% MeOH in CH₂Cl₂, to give the
10 title compound, mp 108-110°C.

¹H NMR (DMSO-d₆): δ 0.91-0.95 (m, 6 H), 1.46 (s, 9 H), 1.92 (s, 3 H),
2.18-2.24 (m, 3 H), 2.77 (d of ABX q, 1 H), 2.96 (d of ABX q, 1 H), 3.06 (br t, 2 H), 4.34 (s, 2 H), 4.90 (br s, 1 H), 6.68 (s, 1 H), 7.24 (s, 1 H), 7.57 (d, 1 H), 7.67 (d, 1 H).

15

Example 26

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one

The title compound was prepared according to General Method 16a using
4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-
20 2-one (Example E-6; 0.33 g, 1.2 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl) ester (Example BB-2; 0.51 g, 1.4 mmol), potassium carbonate (0.48 g, 3.5 mmol), and DMF (5 mL). The product was chromatographed on silica gel, eluting with 5% MeOH in CH₂Cl₂, to give the title compound, mp 98-100°C.

25

¹H NMR (DMSO-d₆): δ 0.86-0.92 (m, 6 H), 1.42 (s, 9 H), 1.87 (s, 3 H),
2.15-2.20 (m, 3 H), 2.25 (s, 3 H), 2.71 (d of ABX q, 1 H), 2.83-2.95 (m, 3 H),
4.30 (s, 2 H), 4.89 (br s, 1 H), 6.64 (s, 1 H), 7.04 (s, 1 H), 7.20 (d, 1 H).

-190-


Example 27

N-(4-{2-[5-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl]-ethyl}-thiazol-2-yl)-acetamide

5 The title compound was prepared according to General Method 16a using N-{4-[2-(4-hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl)-ethyl]-thiazol-2-yl}-acetamide (Example E-13; 0.49 g, 1.5 mmol), toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenyl)ester (Example BB-2; 0.66 g, 1.8 mmol), potassium carbonate (0.83 g, 6.0 mmol), and DMF (5 mL). The
10 product was chromatographed on silica gel, eluting with 5% MeOH in CH₂Cl₂, to give the title compound, mp 128-130°C.

¹H NMR (DMSO-d₆): δ 0.86-0.92 (m, 6 H), 1.42 (s, 9 H), 1.85 (s, 3 H), 2.05 (m, 5 H), 2.17 (m, 1 H), 2.59-2.67 (m, 3 H), 2.92 (d of ABX q, 1 H), 4.30 (s, 2 H), 4.89 (br s, 1 H), 6.25 (s, 1 H), 6.75 (s, 1 H), 7.20 (d, 1 H), 11.92 (br s, 1 H),
15 12.00 (s, 1 H).

TABLE L. Final Dihydropyrones



Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI) (M+H)
28		±		16a	103-105	490 (M+H)
29		±		16a	120-124	476 (M+H)

-192-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
30		±		16a	118-121	490 (M+H)
31		±		16a	105-108	490 (M+H)
32		±		16b	97-100	518 (M+H)

-193-

TABLE L. Final Dihydropyrones (cont'd)



















Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
33		±		16b	109-112	518 (M+H)
34		±		16b	108-112	518 (M+H)
35		±		16a	93-98	460 (M+H)

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
36		±		16a	96-101	453 (M+H)
37		±		16a	90-96	
38		±		16a	104-106	490 (M+H)

-195-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
39		±		16a	110-117	
40		±		16a	136-138	459 (M+H)
41		±		16a	134-135	444 (M+H)






-196-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
42		±		16a	221-223	471 (M+H)
43		±		16a	120-122	456 (M+H)
44		±		16a	145-148	486 (M+H)

-197-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI) (M+H)
45		±		16a	137-140	471 (M+H)
46		±		16a	115-118	476 (M+H)
47	Ph	S		16b	197-199	468 (M+H)

-198-

TABLE L. Final Dihydropyrones (cont'd)













Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
48		S		16b	142	483 (M)
49		±		16a	158-160	
50		S		16a	232-234	513 (M+H)

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
51		S		16b	204-206	
52		R		16a	73-85	475 (M+H)
53		±		16b	107-109	551 (M+H)

-200-

TABLE L. Final Dihydropyrones (cont'd)



















Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
54		±		16a	118-123	
55		±		16a	80-87	
56		±		16a	70-73	

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
57		±		16a		461 (M+H)
58		±		16a	89-91	489 (M+H)
59		±		16a		669







-202-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
60				16a		663
61		±		16a	95-96	508 (M+H)
62		±		16b	107-113	488 (M-H)







-203-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
63		S		16a	115	487 (M-H)
64		±		16a	86-90	487 (M-H)
65		±		16a	98-101	472 (M-H)

-204-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
66		±		16a		489 (M+H)
67		±		16a	104-106	488 (M-H)
68				16a	104-105	498 (M+H)

-205-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
69		±		16a followed by NaOH deprotection	125-126	494 (M+H)
70		S		16a	138-140	647 (M+H)

-206-

TABLE L. Final Dihydropyrones (cont'd)













Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
71		S		16a	171-173	580 (M+H)
72		S		16a followed by NaOH deprotection	103-105	480 (M+H)
73		S		16a followed by NaOH deprotection	123-125	464 (M+H)

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
74		S		16a	122-124	515 (M+H)
75		S		16a	222-224	451 (M+H)
76		S		16a	158-160	467 (M+H)

-208-

TABLE L. Final Dihydropyrones (cont'd)















Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
77		S		16a	203-205	465 (M+H)
78		S		16a	232-234	481 (M+H)
79		S		16a	152-153	573 (M-H)

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
80		S		16a	192-194	536 (M+H)
81		S		Saponification of Example 80 with LiOH	123-125	506 (M-H)





-210-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
82		±		16a	190-192	503 (M+H)
83		±		16a	193-195	515 (M+H)





-211-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
84		±		16a	180-182	529 (M+H)
85		S		16a	173-175	533 (M+H)

-212-

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar ₂	Chirality at C-6	Ar ₁	General Method	Melting Point (°C)	Mass Spe (APCI)
86		S		16a	175-177	539 (M+H)
87		S		16a	221-222	668 (M+H)

-213-

TABLE L. Final Dihydropyrones (cont'd)











Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
88		S		16a	>173	529 (M+H)
89		±		16a followed by NH ₃ /MeOH deprotection	208-210	491 (M+1)
90		+		16a followed by HCl (g) deprotection		482 (M+1)

TABLE L. Final Dihydropyrones (cont'd)

Example	Ar2	Chirality at C-6	Ar1	General Method	Melting Point (°C)	Mass Spe (APCI)
91		+		16a followed by HCl (g) deprotection		472 (M+1)
92		±		16a	155-157	659 (M-1)

The Table L above shows the compounds of the invention as racemic; however, the R and S forms are within the scope of the invention. The S form is the preferred.

-215-

The compound names corresponding to Examples 28-60 in Table L above are:

Example 28: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 29: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

Example 30: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 31: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 32: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 33: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-isopropyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 34: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 35: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

Example 36: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 37: 3-(2-tert-Butyl-4-hydroxy-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 38: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-216-

Example 39: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

5 Example 40: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(1H-pyrazol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 41: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-[2-(1H-pyrazol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

10 Example 42: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyrimidin-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

Example 43: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyrimidin-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

Example 44: 6-[2-(2-Amino-pyrimidin-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

15 Example 45: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(2-amino-pyrimidin-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 46: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-4-yl-ethyl)-5,6-dihydro-pyran-2-one;

20 Example 47: (S)-4-Hydroxy-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-6-phenethyl-5,6-dihydro-pyran-2-one;

Example 48: (S)-4-Hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

25 Example 49: 3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 50: (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 51: (S)-3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 Example 52: (R)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

-217-

Example 53: 3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

5 Example 54: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

Example 55: 3-(2-tert-Butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 56: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

10 Example 57: 3-(2-tert-Butyl-4-hydroxymethyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

Example 58: 3-[2-tert-Butyl-4-(2-hydroxy-ethyl)-5-methyl-phenylsulfanyl]-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

15 Example 59: 5-Trifluoromethyl-pyridine-2-sulfonic acid {5-tert-butyl-4-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-2-methyl-phenyl}-amide; and

Example 60: 5-Trifluoromethyl-pyridine-2-sulfonic acid {5-tert-butyl-4-[4-hydroxy-6-isopropyl-2-oxo-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-2-methyl-phenyl}-amide;

20 Example 61: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 62: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

25 Example 63: (S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 64: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

30

-218-

Example 65: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 66: 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

Example 67: 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 68: 4-Hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

Example 69: 4-Hydroxy-3-(6-tert-butyl-indol-5-ylsulfanyl)-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 70: (S)-N-[6-tert-Butyl-5-(4-hydroxy-6-isopropyl-2-oxo-6-[2-(4-hydroxy-phenyl)ethyl]-5,6-dihydro-2H-pyran-3-ylsulfanyl)-indolin-1-yl]-(4-cyanophenyl)sulfonamide;

Example 71: (S)-6-tert-Butyl-5-{4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-indole-1-carboxylic acid tert-butyl ester;

Example 72: (S)-3-(6-tert-Butyl-indol-5-yl-sulfanyl)-4-hydroxy 6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 73: (S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-4-hydroxy- 6-isopropyl-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

Example 74: (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 75: (S)-4-Hydroxy-6-isopropyl-3-(6-isopropyl-3H-benzoimidazol-5-ylsulfanyl)-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

Example 76: (S)-5-(4-Hydroxy-6-isopropyl-2-oxo-6-phenethyl-5,6-dihydro-2H-pyran-3-ylsulfanyl)-6-isopropyl-1,3-dihydro-benzoimidazol-2-one;

Example 77: (S)-3-(6-tert-Butyl-3H-benzoimidazol-5-yl-sulfanyl)-4-hydroxy-6-isopropyl-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

Example 78: (S)-5-tert-Butyl-6-(4-hydroxy-6-isopropyl-2-oxo-6-(2-phenethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl)-1,3-dihydro-benzoimidazol-2-one;

-219-

Example 79: (S)-N-[5-tert-Butyl-6-(4-hydroxy-6-isopropyl-2-oxo-6-phenethyl-5,6-dihydro-2H-pyran-3-ylsulfanyl)-benzothiazol-2-yl]-methanesulfonamide;

5 Example 80: 6-tert-Butyl-5-(4-hydroxy-6-isopropyl-2-oxo-6-phenethyl-5,6-dihydro-2H-pyran-3-ylsulfanyl)-1H-indole-2-carboxylic acid ethyl ester;

Example 81: 6-tert-Butyl-5-(4-hydroxy-6-isopropyl-2-oxo-6-phenethyl-5,6-dihydro-2H-pyran-3-ylsulfanyl)-1H-indole-2-carboxylic acid;

Example 82: 3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-[(2-(3-thiophenyl)-ethyl)-5,6-dihydro-pyran-2-one];

10 Example 83: (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 84: 3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3-fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

15 Example 85: (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3,5-difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 86: (S)-N-[5-tert-Butyl-6-(4-hydroxy-6-isopropyl-2-oxo-6-(2-phenethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl)-benzothiazol-2-yl]-acetamide;

20 Example 87: (S)-N-{5-tert-Butyl-6-(4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl)-benzothiazol-2-yl}-4-cyanobenzenesulfonamide;

Example 88: (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

25 Example 89: 6-[2-(2-Amino-thiazol-4-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

Example 90: (S)-3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 Example 91: (S)-3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-isopropyl-6-[(2-(3-thiophenyl)-ethyl)-5,6-dihydro-pyran-2-one]; and

Example 92: 4-(6-tert-Butyl-5-{4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl})-2,3-dihydro-indole-1-sulfonyl)-benzonitrile.

-220-

Other compounds which can be prepared by the above methods include:

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(3-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one
3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-Cyclopentyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-cyclopentyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-cyclopentyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-furan-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(3-hydroxymethyl-furan-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-furan-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-furan-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

-221-

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-furan-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-Cyclopentyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

5 3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

10 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

15 6-[2-(2-Amino-thiazol-4-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(2-Amino-thiazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-4-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

25 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-oxazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-222-

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(1H-benzoimidazol-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

5 6-[2-(1H-Benzoimidazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(3-amino-1H-indazol-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

10 6-[2-(3-Amino-1H-indazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

15 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(3-amino-1H-indazol-4-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(3-Amino-1H-indazol-4-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-6-hydroxymethyl-5-methyl-pyridin-3-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

25 3-(2-tert-Butyl-5-methyl-pyridin-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(3-tert-Butyl-2-hydroxymethyl-6-methyl-pyridin-4-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

30 3-(5-tert-Butyl-2-methyl-pyridin-4-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-tert-Butyl-5-hydroxymethyl-thiophen-3-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

-223-

- 3-(4-tert-Butyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 6-[2-(4-Amino-phenyl)-ethyl]-3-(2-tert-butyl-5-methyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 5 3-(2-tert-Butyl-5-methyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 6-[2-(4-Amino-phenyl)-ethyl]-3-(4-tert-butyl-5-hydroxymethyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 6-[2-(4-Amino-phenyl)-ethyl]-3-(3-tert-butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 10 3-(3-tert-Butyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(3-tert-Butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 15 3-(3-tert-Butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 20 3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 25 3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- N-(7-tert-Butyl-6-{6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-acetamide;
- 30 N-(7-tert-Butyl-6-{4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-acetamide;

-224-

N-(7-tert-Butyl-6-{6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-methanesulfonamide;

5 N-(7-tert-Butyl-6-{4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-methanesulfonamide;

3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

10 3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

15 3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-6-hydroxymethyl-5-methyl-pyridin-3-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(3-tert-Butyl-2-hydroxymethyl-6-methyl-pyridin-4-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

20 3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

25 3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

N-{7-tert-Butyl-6-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-4-methyl-benzothiazol-2-yl}-acetamide;

30 N-{7-tert-Butyl-6-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-4-methyl-benzothiazol-2-yl}-methanesulfonamide;

3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

-225-

- 3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- (S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-cyclohexyl-4-hydroxy-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 5 (S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-cyclopentyl-4-hydroxy-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- (S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-n-pentyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-cyclohexyl-4-hydroxy-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 10 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-cyclopentyl-4-hydroxy-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-n-pentyl-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 15 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-cyclohexyl-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-cyclopentyl-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 20 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-n-pentyl-5,6-dihydro-pyran-2-one;
- (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-n-pentyl-6-phenethyl-5,6-dihydro-pyran-2-one;
- 25 (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-cyclohexyl-4-hydroxy-6-phenethyl-5,6-dihydro-pyran-2-one;
- (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-cyclopentyl-4-hydroxy-6-phenethyl-5,6-dihydro-pyran-2-one;
- (S)-4-Hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-propyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;
- 30 (S)-6-Cyclohexyl-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

-226-

(S)-6-Cyclopentyl-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

(S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-4-hydroxy-6-propyl-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

5 (S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-6-cyclohexyl-4-hydroxy-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

(S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-6-cyclopentyl-4-hydroxy-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

10 (S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-4-hydroxy-6-pentyl-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;


(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-6-propyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-6-pentyl-5,6-dihydro-pyran-2-one;

15 (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-cyclohexyl-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-5,6-dihydro-pyran-2-one; and

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-cyclopentyl-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-5,6-dihydro-pyran-2-one.

20 The compounds of the present invention were evaluated for their in vitro inhibition of HIV protease and for their antiviral efficacy in HIV infected lymphocytes and the results shown in Table 1. Two reference agents are included and are shown below.



25 The two Reference Agents 1 and 2 are the two best compounds as disclosed by Hagen, et al., *J. Med. Chem.*, 1997;40:3707-3711). These reference compounds were recognized to have particularly good antiviral activity due in part to the polar substituents placed on the phenyl and phenethylmoieties. In a subsequent disclosure by S. Vander Roest, et al., 37th Interscience Conference on

-227-

Antimicrobial Agents and Chemotherapy, September 28-October 1, 1997, Toronto, Canada Abstract I-84), it was revealed that Reference Agents 1 and 2 had comparatively good pharmacokinetics in mice relative to a large series of related polar substituted phenyl and phenethyl derivatives. Thus, Reference Agents 1 and 2, represent excellent comparative agents for the compounds of the current invention both in vitro and in vivo.

The following sections provide the experimental methodology for the in vitro and in vivo assays employed to demonstrate the efficacy and advantages of the compounds of the current invention.

10 **HIV Protease Assay:**

Materials

Recombinant HIV-1 protease (>96% purity) and HIV protease substrate III (the undecapeptide H-His-Lys-Ala-Arg-Val-Leu-Nph-Glu-Ala-Nle-Ser-NH₂, 97% purity) were purchased from Bachem Bioscience, Inc. (King of Prussia, PA).

15 **Method**

The methods employed follow the procedures of Tunmino, et al., *Archives of Biochemistry and Biophysics*, 1995;316:523). For determination of K_i values, HIV-1 protease, 6.0 nM final concentration, was added to a solution containing inhibitor, 40 μM substrate III and 1.0% Me₂SO in assay buffer: 80 mM MES, 160 mM NaCl, 1.0 mM EDTA 0.1% polyethylene glycol (Mr 8000) pH 6.2 at 37°C (total volume, 100 & 1). Polyethylene glycol was used in the assay in place of glycerol since the former was reported to be a more effective stabilizing agent in the protease (Jordan, et al., *J. Biol. Chem.*, 1992;267:20028). The final inhibitor concentrations used were 0 (0.1, 0.2, in some experiments), 0.5, 1, 2, 5, 10, 20, 50, and 100 μM. The solution was mixed, incubated for 5 minutes and the reaction quenched by addition of trifluoroacetic acid, 2% final. The (leu-*p*-nitro-phe) bond of the substrate is cleaved by the enzyme and substrate and products separated by reverse-phase HPLC. Absorbance was measured at 220 nm, peak areas determined, and percentage conversion to product used to calculate percentage control ($\text{control} = \frac{[\% \text{ conversion (+ inhibitor)}]}{[\% \text{ conversion (- inhibitor)}]} \times 100$).

-228-

Cellular Infection Assay

Compounds were tested in lymphocyte derived CEM cells using the XTT cytopathic procedures and were performed at Southern Research Institute (Buckheit, et al., *Antiviral Res.*, 1993;21:247; see also Weislow, et al., *J. Nat. Cancer Inst.*, 1989;81:577). Compound concentrations were 0.32, 1, 3.2, 10, 32, and 100 μ M. The EC₅₀ represents the concentration of agent which reduces HIV cytopathic effects 50% relative to untreated control. Cellular toxicity of the agents are estimated from the TC₅₀ which represents the concentration of the agent which inhibits 50% of the viability of uninfected cells.

Table 1 contains the results of the HIV protease assay Ki and the antiviral efficacy (EC₅₀, TC₅₀, TI) screening, where TI is the therapeutic index (TC₅₀/EC₅₀).

Mouse Blood Levels

Charles River CD-1 mice were dosed with a 25 mg/kg drug solution orally by gavage. For oral dosing the compounds were dissolved in 0.1N NaOH (20% final volume) and then 0.5% methylcellulose (80% final volume) was added. Blood was drawn and pooled from 5 mice at each time point via heart puncture and was placed in a 15 mL centrifuge tube containing heparin. The plasma stored in 200 μ L aliquots at -80°C until assayed.

Plasma was thawed at room temperature and then 400 μ L of acetonitrile was added to each vial and vortexed. Following centrifugation the supernatant was removed and evaporated in a nitrogen evaporator at 48°C. Residue was then resolubilized in 200 μ L of a 75% water/25% CH₃CN solution for HPLC analysis. HPLC analysis used a Zorbax RX-C8 column (4.6 \times 150 mm) with a 25 mM potassium phosphate buffer with 0.1% TFA, pH 3.0. Mobile phases consisted of varying percentages of two buffer mixtures: Buffer A - 90% buffer, 10% CH₃CN, and Buffer B - 30% buffer, 70% CH₃CN. Absorbance was determined at a wavelength of 260 nm.

A standard curve ranging from 0.5 to 50 μ g/mL was used for each drug tested. Ten times final concentration solutions of drug in water were prepared

-229-

from stock solutions of 1 mg/mL drug in DMSO. The final 1:10 dilution was then done in mouse plasma obtained from mice described above.

The results from testing the reference agents and the compounds of this invention are shown in Table 2.

5 **Rat and Dog Blood Levels**

The procedure for pharmacokinetic analysis in dogs and rats was similar. For dog studies, two dogs received a single oral 10 mg/kg dose of the HIV protease inhibitor as a solution. The protease inhibitor was dissolve in 0.1N NaOH followed by buffering or diluting with isotonic phosphate buffer or water
10 depending on solubility. For rat studies, two male rats received a single oral 10 mg/kg gavage dose of the HIV protease inhibitor suspended in 0.5% methylcellulose. For the IV administration, the compounds were dissolved in 0.1N sodium hydroxide at a concentration of 10 mg/ml. The IV dose was given bolus. Blood samples were collected prior to dosing and at 0.25, 0.5, 1, 1.5, 2, 3, 4, 8, 12,
15 and 24 hours postdose. Plasma samples were analyzed for the HIV protease inhibitor using a HPLC procedure. The method involved plasma protein precipitation with acetonitrile, reverse-phase chromatographic separation of the HIV protease inhibitor and internal standard, and quantitation by UV detection. Some plasma samples were concentrated by evaporation prior to injection.
20 Pharmacokinetic parameters were determined by noncompartmental analysis of individual rat-plasma concentration-time curves using WinNonlin. Maximum concentration (C_{max}) and time for these to occur (t_{max}) were recorded as observed. Area under concentration-time curve (AUC) values were obtained using the linear trapezoidal rule. The results of the pharmacokinetic studies in dogs and
25 rats are shown in Table 3.

Table 1 indicates that the compounds of the present invention have good to excellent activity toward inhibiting the HIV protease enzyme (K_i's) as well as activity in HIV infected cells, protecting the cells from HIV pathogenicity at μ M and sub- μ M concentrations. The two reference agents in Table 1, which represent
30 the best of the hydroxylated and aminated compounds from the art, are wholly comparable to the compounds of the current invention at the antiviral level. Yet the current invention relies not only on the antiviral activity but on the unexpected

-230-

finding that certain heterocycles could sufficiently mimic the polar groups of the reference agents while effecting significant gains in pharmacokinetic parameters due in part to reduced metabolism of the amines and phenols of the reference agents.

5 In Table 2, the results from mouse pharmacokinetic studies are shown for the reference agents and certain of the compounds of the present invention. In Table 3, the oral pharmacokinetics of certain compounds and the reference agents in rats and dogs are shown. Table 4 shows the pharmacokinetics in rats compared to the reference agents.

10 The results in Table 2 show a clear improvement in C_{max} and T_{1/2} of the compounds of the invention relative to the Reference Agents. The results from Table 3 show that the pharmacokinetic advantages of the compounds of the current invention are not limited to mice but are manifest in rats and dogs. Such data all support the unexpected observation that select heterocycles mimic the
15 polar hydroxylated and aminated phenyl groups, yet possess unexpectedly good pharmacokinetics relative to the best Reference Agents.

-231-

TABLE 1

Inhibition of HIV-Protease and Antiviral Efficacy in a CEM Cell Infection Assay
(Page 1 of 2)

Example	Inhib of HIV Protease KI (nM)	Antiviral Efficacy		
		EC ₅₀ (μM)	TC ₅₀ (μM)	TI
1	1.5	1.5	296	197
2	0.9	0.9	216	240
4	0.35	1.0	199	199
5	0.78	0.75	190	253
7	0.17	0.3	110	366
8	0.35	0.5	80.5	161
9	0.19	1.8	210	117
11	0.23	0.29	133	458
12	0.07	0.14	91	650
13		0.28	42	152
14	0.67	2.5	75	30
16		11	114	10.4
17		2.7	42	16
18		1.1	117	105
22	0.85	1.5	66	43
24		0.37	164	443
25	0.57	0.7	216	308
26	0.45	0.8	171	213
28	0.24	1.9	229	120
32	1.6	23	91	4
33		1.4	101	72
35	0.88	1.5	72	48
36	1.7	2.2	170	77
37	0.37	4.2	100	24
38	0.33	1.1	124	113
40	0.28	5.6	>320	>57
41	6.5	20	195	10
42		1.3	>200	>159
48	0.4	0.37	29	78
55	1.7	8.3	66	8
56	11	10	210	21
59	0.08	0.6	66	110
61	0.98	0.68	46	68
65	1.2	1.1	60	54
69	0.67	0.18	55	306
70	0.82	1.3	8	6
73	0.54	0.61	42	69

-232-

TABLE 1

Inhibition of HIV-Protease and Antiviral Efficacy in a CEM Cell Infection Assay
(Page 2 of 2)

Example	Inhib of HIV Protease KI (nM)	Antiviral Efficacy		
		EC ₅₀ (μM)	TC ₅₀ (μM)	TI
79	2.1	2.2	125	56
82	1.1	0.7	76	109
83	0.44	0.26	61	235
84	0.8	0.47	93	198
89	0.95	3.5	>200	>57
Ref 1	0.43	0.5	>100	>200
Ref 2	0.22	0.6	>100	>166

TABLE 2

Pharmacokinetic Evaluation of HIV Protease Inhibitors in Mice Following
an Oral Dose of 25 mg/kg

Example	C _{max} ^a (μM)	T 1/2 ^b (hr)
Ref 1	23	2.2
Ref 2	17	2.0
2	46	7.3
4	54	5.6
7	40	3.5
12	95	9.5
25	33	3.3

^a The maximum concentration reached within the plasma.

^b The half life of the agents.

-233-

TABLE 3
Pharmacokinetic Evaluation of HIV Protease Inhibitors in Rats and Dogs
Following an Oral Dose of 10 mg/kg

Example	Rats		Dogs	
	C _{max} ^a (μ M)	T 1/2 ^b (hr)	C _{max} (μ M)	T 1/2 ^b (hr)
Ref 1	16	3.4	140	1.25
Ref 2	3.4	0.55	40	0.52
12	30	11.5	149	1.75
13	5.1	8.6	--	--
25	6.0	12.0	--	--

^a The maximum concentration reached within the plasma.

^b The half life of the agents.

TABLE 4
Pharmacokinetic Evaluation of HIV Protease Inhibitors in Rats Following
an IV Dose of 10 mg/kg


Example	C(0) (μ M)	T 1/2 ^a (hr)
Ref 1	86	3.7
Ref 2	100	3.9
12	160	12.0
25	120	8.0
69	335	4.5
83	199	4.0

^a The half life of the agents.

-234-

CLAIMS

1. A compound of Formula I



or a pharmaceutically acceptable salt thereof wherein

- 5 R_1 is H, a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;
- R_2 is H, a straight or branched alkyl of 1-5 carbons;
- R_3 is H, $(CR'_2)_nOR$, $(CR'_2)_nN(R)_2$, $(CR'_2)_nNR'COR$, $(CR'_2)_nCO_2R$,
 $(CR'_2)_nOCOR$, $(CR'_2)_nCON(R)_2$, $(CR'_2)_nOCON(R)_2$, $(CR'_2)_nR$,
 10 $(CR'_2)_nNR'CON(R)_2$, $(CR'_2)_nNR'CO_2R$, $(CR'_2)_nOSO_2N(R)_2$,
 $(CR'_2)_nNR'SO_2OR$, $(CR'_2)_nNR'SO_2N(R)_2$, $(CR'_2)_nOSO_2R$,
 $(CR'_2)_nNR'SO_2R$, $(CR'_2)_nSO_pR$, $(CR'_2)_nNR'CSN(R)_2$,
 $(CR'_2)_nNR'C(NR')N(R)_2$, $(CR'_2)_nSO_2N(R)_2$,
 $(CR'_2)_nC(NR')N(R)_2$, $(CR'_2)_nCOR$, $O(CR'_2)_mOR$,
 15 $NR(CR'_2)_mOR$, F, Cl, Br, CF_3 , CN, or =O;
- R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of 1-6 carbons, a cycloalkyl of 3-6 carbons, $(CR'_2)_nOR$,
 $(CR'_2)_nN(R)_2$, F, Cl, Br, CN, CF_3 , =O, $(CR'_2)_pNR'COR$,
 $(CR'_2)_pSO_pR$, $(CR'_2)_pR$, $(CR'_2)_pOCOR$, $O(CR'_2)_mOR$,
 20 $NR(CR'_2)_mOR$, $(CR'_2)_pNR'CON(R)_2$, $(CR'_2)_pOCON(R)_2$,
 $(CR'_2)_pNR'CO_2R$, $(CR'_2)_pCOR$, $(CR'_2)_pCO_2R$,
 $(CR'_2)_pCON(R)_2$, $(CR'_2)_pNR'SO_2R$, $(CR'_2)_pSO_2N(R)_2$,
 $(CR'_2)_pNR'SO_2OR$, $(CR'_2)_pOSO_2N(R)_2$, $(CR'_2)_pNR'SO_2N(R)_2$,
 $(CR'_2)_pC(NR')N(R)_2$, $(CR'_2)_pNR'C(NR')N(R)_2$, $(CR'_2)_pHet$;

-235-

any two of R₁-R₃ or R₄-R₆ may together form a ring of 5-6 total atoms

which may contain 0-3 heteroatoms;

n is an integer of from 0 to 3;

m is an integer of from 2 to 4;

5 p is an integer from 0 to 2;

R₇ is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R is independently H, a straight or branched alkyl of 1-4 carbons,

(CH₂)_nPh, or a (CH₂)_nheterocycle of 5-6 atoms containing

10 1-2 heteroatoms and wherein the (R)₂ in N(R)₂ may form a

heterocycle containing the nitrogen, all optionally substituted by F,

Cl, Br, OR', CN, CO₂R', N(R')₂, NR'COR', CF₃, or =O;

R' is independently H, a straight or branched alkyl of 1-4 carbons, or phenyl;

15 R'' is independently H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR', or N(R')₂;

Ar₁ and Ar₂ are independently phenyl or Het with the proviso that at least one Ar is Het;

Het is a heterocycle of from 5-6 atoms having from 1-4 heteroatoms or a

20 fused heterocycle of from 9-10 atoms having 1-3 heteroatoms.

2. A compound according to Claim 1 wherein:

R₁ is a straight or branched alkyl of from 1 to 4 carbons or a carbocycle of from 3 to 5 carbons and

R₂ is H or a straight or branched alkyl of from 1 to 3 carbons.

25 3. A compound according to Claim 1 wherein:

R₁ is isopropyl or t-butyl;

R₂ is H, methyl, or ethyl;

R₃ is H, (CR'₂)_nOR, (CR'₂)_nN(R)₂, (CR'₂)_nNR'COR, (CR'₂)_nCO₂R, (CR'₂)_nOCOR, (CR'₂)_nCON(R)₂, (CR'₂)_nOCON(R)₂,

-236-

- $(CR'_2)_nNR'CON(R)_2$, $(CR'_2)_nNR'CO_2R$, $(CR'_2)_nOSO_2N(R)_2$,
 $(CR'_2)_nNR'SO_2OR$, $(CR'_2)_nNR'SO_2N(R)_2$, $(CR'_2)_nOSO_2R$,
 $(CR'_2)_nNR'SO_2R$, $(CR'_2)_nSO_pR$, $(CR'_2)_nNR'CSN(R)_2$,
 $(CR'_2)_nCOR$, $O(CR'_2)_mOR$, $NR(CR'_2)_mOR$, F, Cl, Br, CF_3 , CN,
 5 or =O;
 R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of
 1-6 carbons, a cycloalkyl of 3-6 carbons, $(CH_2)_nOR$,
 $(CH_2)_nN(R)_2$, F, Cl, Br, CN, CF_3 , =O, $(CR'_2)_pNR'COR$,
 $(CR'_2)_pSO_pR$, $(CR'_2)_pNR'CON(R)_2$, $(CR'_2)_pOCON(R)_2$,
 10 $(CR'_2)_pNR'CO_2R$, $(CR'_2)_pCOR$, $(CR'_2)_pCO_2R$,
 $(CR'_2)_pCON(R)_2$, $(CR'_2)_pNR'SO_2R$, $(CR'_2)_pSO_2N(R)_2$,
 $(CR'_2)_pNR'SO_2OR$, $(CR'_2)_pOSO_2N(R)_2$, $(CR'_2)_pHet$ and any
 two of R_1 - R_3 or R_4 - R_6 may together form a ring of 5-6 total atoms
 which may contain 0-2 heteroatoms;
 15 n is an integer of 0 to 3;
 m is an integer of 2 to 4;
 p is 0 to 2;
 R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of
 3-6 carbons;
 20 R is H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a
 $(CH_2)_n$ heterocycle of 5-6 atoms containing 1-2 heteroatoms and
 wherein the $(R)_2$ in $N(R)_2$ may form a heterocycle containing the
 nitrogen, all optionally substituted by F, Cl, Br, OR' , $N(R')_2$,
 $NR'COR'$, CF_3 , or =O;
 25 R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;
 R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or
 $N(R')_2$;
 Ar_1 is phenyl or Het; and

-237-

Ar₂ is Het wherein Het is a heterocycle of from 5-6 atoms having

1-4 heteroatoms selected from: furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine, piperidine, morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of from 9-10 atoms having from 1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

4. A compound according to Claim 1 wherein:

R₁ is isopropyl or t-butyl;

R₂ is H, methyl or ethyl;

R₃ is H, (CH₂)_nOR, (CH₂)_nN(R)₂, (CH₂)_nNR'COR, (CH₂)_nCON(R)₂, (CH₂)_nOCON(R)₂, (CH₂)_nNR'CON(R)₂, (CH₂)_nNR'CO₂R, (CH₂)_nOSO₂N(R)₂, (CH₂)_nNR'SO₂OR, (CH₂)_nNR'SO₂N(R)₂, (CH₂)_nOSO₂R, (CH₂)_nNR'SO₂R, (CH₂)_nSO₂R, (CH₂)_nNR'CSN(R)₂, (CH₂)_nCOR, O(CH₂)_mOR', NR(CH₂)_mOR', or C(CH₃)₂OR';

R₄, R₅, and R₆ are independently H, a straight or branched alkyl of 1-6 carbons, a cycloalkyl of 3-6 carbons, (CH₂)_nOR, (CH₂)_nN(R)₂, F, Cl, Br, CN, CF₃, =O, (CH₂)_pNR'COR, (CH₂)_pNR'CON(R)₂, (CH₂)_pOCON(R)₂, (CH₂)_pNR'CO₂R, (CH₂)_pCOR, (CH₂)_pCON(R)₂, (CH₂)_pNR'SO₂R, (CH₂)_pNR'SO₂OR, (CH₂)_pOSO₂N(R)₂, wherein p is 0 to 2, or R₄ and R₅ may together form a ring of 5-6 total atoms which may contain 0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

-238-

R₇ is a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, (CH₂)_nPh, or a (CH₂)_nheterocycle of 5-6 atoms containing 1-2 heteroatoms and wherein the (R)₂ in N(R)₂ may be a heterocycle containing the nitrogen, all optionally substituted by F, Cl, Br, OR', N(R')₂, NR'COR', or =O;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR', or N(R')₂;

Ar₁ is phenyl; and

Ar₂ is a heterocycle of 5-6 atoms having from 1-4 heteroatoms selected from: furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine, pyridazine, pyrimidine, pyrazine, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine, piperidine, morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of 9-10 atoms having from 1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

5. A compound according to Claim 1 wherein:

R₁ is H, a straight or branched alkyl of 1-6 carbons or a carbocycle of 3-6 carbons;

R₂ is H, a straight or branched alkyl of 1-5 carbons;

R₃ is H, (CH₂)_nOR, (CH₂)_nN(R)₂, (CH₂)_nNR'COR, (CH₂)_nCON(R)₂, (CH₂)_nOCON(R)₂, (CH₂)_nNR'CON(R)₂, (CH₂)_nNR'CO₂R, (CH₂)_nOSO₂N(R)₂, (CH₂)_nNR'SO₂OR, (CH₂)_nNR'SO₂N(R)₂, (CH₂)_nOSO₂R, (CH₂)_nNR'SO₂R, (CH₂)_nSO₂R, (CH₂)_nCOR,

-239-

$O(CH_2)_mOR$, $NR(CH_2)_mOR$, $C(CH_3)_2OR'$, F, Cl, Br, CF_3 , or $=O$;

R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of

1-6 carbons, a cycloalkyl of 3-6 carbons, $(CH_2)_nOR$,

5 $(CH_2)_nN(R)_2$, F, Cl, Br, $=O$, $(CH_2)_pNR'COR$,

$(CH_2)_pNR'CON(R)_2$, $(CH_2)_pOCON(R)_2$, $(CH_2)_pNR'CO_2R$,

$(CH_2)_pCOR$, $(CH_2)_pCON(R)_2$, $(CH_2)_pNR'SO_2R$,

$(CH_2)_pNR'SO_2OR$, $(CH_2)_pOSO_2N(R)_2$, wherein p is 0 to 2;

R_4 and R_5 may together form a ring of 5-6 total atoms which may contain

10 0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

R_7 is methyl, ethyl, isopropyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl;

15 R is H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a

$(CH_2)_n$ heterocycle of 5-6 atoms containing 1-2 heteroatoms and

wherein the $(R)_2$ in $N(R)_2$ may be a heterocycle containing the

nitrogen, all optionally substituted by F, Cl, Br, OR' , $N(R')_2$,

$NR'COR'$, or $=O$;

20 R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or

$N(R')_2$;

Ar_1 is phenyl or Het; and

Ar_2 is Het wherein Het is a heterocycle of 5-6 atoms having from

25 1-4 heteroatoms selected from: furan, pyrrole, thiophene, oxazole,

isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole,

tetrazole, pyridine, pyridazine, pyrimidine, pyrazine,

tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine,

piperidine, morpholine, thiomorpholine, oxolane, dioxane,

30 sulfolane; or a fused heterocycle of from 9-10 atoms having

-240-

1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

6. A compound according to claim 1 wherein:

5 R_1 is H, methyl, ethyl, isopropyl, or t-butyl;

R_2 is H or methyl;

R_3 is H, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, $(CH_2)_nNR'COR$, $(CH_2)_nCON(R)_2$,

$(CH_2)_nOCON(R)_2$, $(CH_2)_nNR'CON(R)_2$, $(CH_2)_nNR'CO_2R$,

$(CH_2)_nOSO_2N(R)_2$, $(CH_2)_nNR'SO_2OR$, $(CH_2)_nNR'SO_2N(R)_2$,

10 $(CH_2)_nOSO_2R$, $(CH_2)_nNR'SO_2R$, $(CH_2)_nSO_2R$, $(CH_2)_nCOR$,

$O(CH_2)_mOR'$, $NR(CH_2)_mOR'$, $C(CH_3)_2OR'$, F, Cl, Br, CF_3 , or $=O$;

R_4 , R_5 , and R_6 are independently H, a straight or branched alkyl of

1-6 carbons, a cycloalkyl of 3-6 carbons, $(CH_2)_nOR$,

15 $(CH_2)_nN(R)_2$, F, Cl, Br, CF_3 , $(CH_2)_pNR'COR$,

$(CH_2)_pNR'CON(R)_2$, $(CH_2)_pOCON(R)_2$, $(CH_2)_pNR'CO_2R$,

$(CH_2)_pCOR$, $(CH_2)_pCON(R)_2$, $(CH_2)_pNR'SO_2R$,

$(CH_2)_pNR'SO_2OR$, $(CH_2)_pOSO_2N(R)_2$, $(CH_2)_pHet$ wherein p is

0 to 2;

20 any two of R_4 - R_6 may together form a ring of 5-6 total atoms which may contain 0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of

25 3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a

$(CH_2)_n$ heterocycle of 5-6 atoms having from 1-2 heteroatoms and

wherein the $(R)_2$ in $N(R)_2$ may form a heterocycle having

-241-

nitrogen, all optionally substituted by F, Cl, Br, OR', N(R')₂,

NR'COR', or =O;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR', or

5 N(R')₂;

Ar₁ is a heterocycle of 5-6 atoms having from 1-4 heteroatoms selected

from: furan, pyrrole, thiophene, oxazole, isoxazole, thiazole,

pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine,

pyridazine, pyrimidine, pyrazine, tetrahydrofuran,

10 tetrahydrothiophene, pyrrolidine, piperazine, piperidine,

morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a

fused heterocycle of 9-10 atoms having 1-3 heteroatoms selected

from: benzofuran, indole, indoline, benzothiophene,

benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline,

15 cinnoline, quinazoline and quinoxaline; and

Ar₂ is phenyl.

7. A compound according to Claim 1 wherein:

R₁ is isopropyl or t-butyl;

R₂ is H or methyl;

20 R₃ is H, (CH₂)_nOR, (CH₂)_nN(R)₂, (CH₂)_nOCON(R)₂,

(CH₂)_nNR'CON(R)₂, (CH₂)_nNR'CO₂R, (CH₂)_nOSO₂N(R)₂,

(CH₂)_nNR'SO₂OR, (CH₂)_nNR'SO₂N(R)₂, (CH₂)_nNR'SO₂R,

(CH₂)_nSO₂R, O(CH₂)_mOR', NR(CH₂)_mOR', or C(CH₃)₂OR';

R₄, R₅, and R₆ are independently H, a straight or branched alkyl of

25 1-6 carbons, a cycloalkyl of 3-6 carbons, (CH₂)_nOR,

(CH₂)_nN(R)₂, F, Cl, Br, =O, (CH₂)_pNR'COR,

(CH₂)_pNR'CON(R)₂, (CH₂)_pOCON(R)₂, (CH₂)_pNR'CO₂R,

(CH₂)_pCOR, (CH₂)_pCON(R)₂, (CH₂)_pNR'SO₂R,

(CH₂)_pNR'SO₂OR, (CH₂)_pOSO₂N(R)₂, wherein p is 0 to 2;

-242-

R₄ and R₅ may together form a ring of 5-6 total atoms which may contain
0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

5 R₇ is a straight or branched alkyl of 1-6 carbons or a carbocycle of
3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, (CH₂)_nPh, or a
(CH₂)_nheterocycle of 5-6 atoms having from 1-2 heteroatoms and
wherein the (R)₂ in N(R)₂ may be a heterocycle containing the
10 nitrogen, all optionally substituted by F, Cl, Br, OR', N(R')₂,
NR'COR', or =O;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR', or
N(R')₂;

15 Ar₁ is phenyl; and

Ar₂ is furan, pyrrole, thiophene, oxazole, isoxazole, thiazole, pyrazole,
tetrazole, or pyridine.

8. A compound according to Claim 1 wherein:

20 R₁ is H, a straight or branched alkyl of 1-6 carbons or a carbocycle of
3-6 carbons;

R₂ is H, methyl or ethyl or isopropyl;

R₃ is H, (CH₂)_nOR, (CH₂)_nN(R)₂, (CH₂)_nNRCOR, (CH₂)_nCON(R)₂,
(CH₂)_nOCON(R)₂, (CH₂)_nNR'CON(R)₂, (CH₂)_nNR'CO₂R,
(CH₂)_nOSO₂N(R)₂, (CH₂)_nNR'SO₂OR, (CH₂)_nNR'SO₂N(R)₂,
25 (CH₂)_nOSO₂R, (CH₂)_nNR'SO₂R, (CH₂)_nSO₂R, O(CH₂)_mOR,
NR(CH₂)_mOR', C(CH₃)₂OR', or =O;

R₄, R₅, and R₆ are independently H, a straight or branched alkyl of
1-6 carbons, a cycloalkyl of 1-6 carbons, (CH₂)_nOR,

-243-

$(\text{CH}_2)_n\text{N}(\text{R})_2$, F, Cl, Br, =O, $(\text{CH}_2)_p\text{NR}'\text{COR}$,
 $(\text{CH}_2)_p\text{NR}'\text{CON}(\text{R})_2$, $(\text{CH}_2)_p\text{OCON}(\text{R})_2$, $(\text{CH}_2)_p\text{NR}'\text{CO}_2\text{R}$,
 $(\text{CH}_2)_p\text{COR}$, $(\text{CH}_2)_p\text{CON}(\text{R})_2$, $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{R}$,
 $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{OR}$, $(\text{CH}_2)_p\text{OSO}_2\text{N}(\text{R})_2$, wherein p is 0 to 2;

5 R_4 and R_5 may together form a ring of 5-6 total atoms which may contain
 0-2 heteroatoms;

n is an integer of 0 to 3;

m is an integer of 2 to 4;

10 R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of
 3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, $(\text{CH}_2)_n\text{Ph}$, or a
 $(\text{CH}_2)_n$ heterocycle of 5-6 atoms containing 1-2 heteroatoms and
 wherein the $(\text{R})_2$ in $\text{N}(\text{R})_2$ may be a heterocycle having nitrogen,
 all optionally substituted by F, Cl, Br, OR' , $\text{N}(\text{R}')_2$, $\text{NR}'\text{COR}'$, or
 15 =O;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or
 $\text{N}(\text{R}')_2$;

20 Ar_1 is thiophene, thiazole, pyridine, benzothiophene, benzthiazole,
 benzoxazole, quinoline or isoquinoline; and

25 Ar_2 is phenyl or Het wherein Het is a heterocycle of 5-6 atoms having
 from 1-4 heteroatoms selected from: furan, pyrrole, thiophene,
 oxazole, isoxazole, thiazole, pyrazole, 1,2,3-triazole, 1,2,4-triazole,
 tetrazole, pyridine, pyridazine, pyrimidine, pyrazine,
 tetrahydrofuran, tetrahydrothiophene, pyrrolidine, piperazine,
 piperidine, morpholine, thiomorpholine, oxolane, dioxane,
 sulfolane; or a fused heterocycle of 9-10 atoms having
 1-3 heteroatoms selected from: benzofuran, indole, indoline,
 benzothiophene, benzimidazole, benzthiazole, benzoxazole,
 30 quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

-244-

9. A compound according to Claim 1 wherein:

R_1 is isopropyl or t-butyl;

R_2 is H, methyl, or ethyl;

R_3 is CH_2OH , NH_2 , $\text{OCH}_2\text{CH}_2\text{OH}$, NHCOR , $\text{OSO}_2\text{N(R)}_2$, $\text{NR}'\text{SO}_2\text{OR}$,

5 $\text{NR}'\text{SO}_2\text{R}$, or OSO_2R ;

R_4 , R_5 and R_6 are independently H, a straight or branched alkyl of

1-6 carbons, a cycloalkyl of 1-6 carbons, $(\text{CH}_2)_n\text{OR}$,

$(\text{CH}_2)_n\text{N(R)}_2$, F, Cl, Br, =O, $(\text{CH}_2)_p\text{NR}'\text{COR}$,

$(\text{CH}_2)_p\text{NR}'\text{CON(R)}_2$, $(\text{CH}_2)_p\text{OCON(R)}_2$, $(\text{CH}_2)_p\text{NR}'\text{CO}_2\text{R}$,

10 $(\text{CH}_2)_p\text{COR}$, $(\text{CH}_2)_p\text{CON(R)}_2$, $(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{R}$,

$(\text{CH}_2)_p\text{NR}'\text{SO}_2\text{OR}$, $(\text{CH}_2)_p\text{OSO}_2\text{N(R)}_2$, wherein p is 0 to 2;

R_4 and R_5 may together form a ring of 5-6 total atoms which may contain

0-2 heteroatoms;

R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of

15 3-6 carbons,

R is H, a straight or branched alkyl of 1-4 carbons, $(\text{CH}_2)_n\text{Ph}$, or a

$(\text{CH}_2)_n$ heterocycle of 5-6 atoms having from 1-2 heteroatoms and

wherein the $(\text{R})_2$ in N(R)_2 may be a heterocycle having nitrogen,

all optionally substituted by F, Cl, Br, OR' , N(R')_2 , $\text{NR}'\text{COR}'$, or

20 =O;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of from 1-4 carbons, F, Cl, Br, OR' ,

or N(R')_2 ;

Ar_1 is phenyl; and

25 Ar_2 is a heterocycle of 5-6 atoms having from 1-4 heteroatoms selected

from: furan, pyrrole, thiophene, oxazole, isoxazole, thiazole,

pyrazole, 1,2,3-triazole, 1,2,4-triazole, tetrazole, pyridine,

pyridazine, pyrimidine, pyrazine, tetrahydrofuran,

tetrahydrothiophene, pyrrolidine, piperazine, piperidine,

-245-

morpholine, thiomorpholine, oxolane, dioxane, sulfolane; or a fused heterocycle of 9-10 atoms having from 1-3 heteroatoms selected from: benzofuran, indole, indoline, benzothiophene, benzimidazole, benzthiazole, benzoxazole, quinoline, isoquinoline, cinnoline, quinazoline and quinoxaline.

5

10. A compound according to Claim 1 wherein:

R_1 is H, methyl, ethyl, isopropyl, or t-butyl;

R_2 is H or methyl;

R_3 is H, $(CH_2)_nOR$, $(CH_2)_nN(R)_2$, $(CH_2)_nNR'COR$, $(CH_2)_nCON(R)_2$,

10

$(CH_2)_nOCON(R)_2$, $(CH_2)_nNR'CON(R)$, $(CH_2)_nNR'CO_2R$,

$(CH_2)_nOSO_2N(R)_2$, $(CH_2)_nNR'SO_2OR$, $(CH_2)_nNR'SO_2N(R)_2$,

$(CH_2)_nOSO_2R$, $(CH_2)_nNR'SO_2R$, $(CH_2)_nSO_2R$,

$(CH_2)_nNR'CSN(R)_2$, $(CH_2)_nCOR$, $O(CH_2)_mOR$,

$NR(CH_2)_mOR'$, $C(CH_3)_2OR'$, F, Cl, Br, or =O;

15

R_4 , R_5 and R_6 are independently H, methyl, ethyl, OH, CH_2OH ,

CH_2CH_2OH , F, Cl, NH_2 ;

any two R_4 - R_6 may form a ring of 5-6 atoms having from

1-2 heteroatoms;

n is an integer of 0 to 3;

20

m is an integer of 2 to 4;

R_7 is a straight or branched alkyl of 1-6 carbons or a carbocycle of

3-6 carbons;

R is H, a straight or branched alkyl of 1-4 carbons, $(CH_2)_nPh$, or a

$(CH_2)_nheterocycle$ of 5-6 atoms having from 1-2 heteroatoms or a

25

heterocycle having a nitrogen, all optionally substituted by F, Cl,

Br, OR' , $N(R')_2$, $NR'COR'$, or =O;

R' is H, a straight or branched alkyl of 1-4 carbons, or phenyl;

R'' is H, a straight or branched alkyl of 1-4 carbons, F, Cl, Br, OR' , or

$N(R')_2$;

-246-

Ar₁ is furan, thiophene, thiazole, pyridine, imidazole, benzofuran, benzothiophene, benzimidazole, benzthiazole, quinoline, or isoquinoline; and
Ar₂ is phenyl.

- 5 11. A compound according to Claim 1 wherein:
 R₁ is H, methyl, ethyl, isopropyl, or t-butyl;
 R₂ is H or methyl;
 R₃ is H, (CH₂)_nOR, (CH₂)_nN(R)₂, or =O;
 R₄, R₅, and R₆ are independently H, methyl, OH, CH₂OH, CH₂CH₂OH,
10 NH₂, or F;
 R₇ is H, isopropyl, butyl, sec-butyl, cyclobutyl, cyclopentyl, or cyclohexyl;
 R is H, methyl, ethyl, phenyl, or CH₂Ph and wherein the (R)₂ of N(R)₂ may be a heterocycle having a nitrogen;
15 R'' is H, F, or CH₃;
 Ar₁ is furan, thiophene, thiazole, pyridine, imidazole, benzofuran, benzothiophene, benzimidazole, benzthiazole, quinoline, or isoquinoline; and
 Ar₂ is phenyl.
- 20 12. A compound according to Claim 1 wherein:
 R₁ is H, methyl, ethyl, isopropyl, or t-butyl;
 R₂ is H, methyl, ethyl, or isopropyl;
 R₃ is H, NH₂, OH, CH₂OR, CH₂N(R)₂, CH₂CON(R)₂,
 CH₂OSO₂N(R)₂, CH₂NHSO₂OR, CH₂NHSO₂R, CH₂OSO₂R,
25 Cl, Br, or OCH₂CH₂OH;
 R₄, R₅, and R₆ are independently H, methyl, ethyl, isopropyl, OH, NH₂, CH₂OR, CH₂N(R)₂, =O, F, Cl, Br, or CH₂NRCOR;
 R₇ is methyl, ethyl, isopropyl, cyclopropyl, cyclobutyl, or cyclopentyl;

-247-

R is H, methyl, ethyl, Ph, CH₂Ph, and wherein the (R)₂ in N(R)₂ may be a heterocycle having a nitrogen;

R'' is H, F, or CH₃;

Ar₁ is thiophene, thiazole, furan, pyridine, benzothiophene, benzofuran,

5 benzthiazole, benzoxazole, quinoline, or isoquinoline; and

Ar₂ is furan, thiophene, oxazole, isoxazole, imidazole, thiazole, pyrazole,

pyridine, benzofuran, benzothiophene, benzimidazole,

benzthiazole, quinoline, or isoquinoline.

13. A compound according to Claim 1 wherein:

10 R₁ is isopropyl or t-butyl;

R₂ is H, methyl, or ethyl;

R₃ is H, NH₂, OH, CH₂OR, CH₂N(R)₂, CH₂CON(R)₂, OSO₂N(R)₂,

NHSO₂OR, NHSO₂R, OSO₂R, or OCH₂CH₂OH;

R₄, R₅, and R₆ are independently H, methyl, ethyl, isopropyl, OH, NH₂,

15 CH₂OR, CH₂N(R)₂, =O, F, Cl, Br, or CH₂NRCOR;

R₇ is methyl, ethyl, isopropyl, cyclopropyl, cyclobutyl, or cyclopentyl;

R is H, methyl, ethyl, Ph, CH₂Ph, and wherein the (R)₂ in N(R)₂ may be a

heterocycle containing the nitrogen;

R'' is H, F, or CH₃;

20 Ar₁ is phenyl; and

Ar₂ is furan, thiophene, oxazole, isoxazole, imidazole, thiazole, pyrazole,

pyridine, benzofuran, benzothiophene, benzimidazole,

benzthiazole, quinoline, or isoquinoline.

14. A compound according to Claim 1 wherein:

25 R₁ is isopropyl or t-butyl;

R₂ is H, methyl, or ethyl;

R₃ is NH₂, CH₂OH, OCH₂CH₂OH, or CH₂CH₂OH;

-248-

- R₄, R₅, and R₆ are independently H, NH₂, CH₂OH, =O, methyl, ethyl, or isopropyl;
R₇ is isopropyl;
R'' is H, F, or CH₃;
5 Ar₁ is phenyl; and
Ar₂ is furan, thiophene, imidazole, thiazole, pyrazole, or pyridine.
15. A compound according to Claim 1 wherein:
R₁ is H, methyl, ethyl, isopropyl, or t-butyl;
R₂ is H or methyl;
10 R₃ is H, CH₂OH, NH₂, or =O;
R₄, R₅, and R₆ are independently H, OH, CH₂OH, NH₂, or F;
R₇ is isopropyl, sec-butyl, isobutyl, or cyclopentyl;
R is H, methyl, ethyl, Ph, CH₂Ph, and wherein the (R)₂ in N(R)₂ may be a heterocycle having a nitrogen;
15 R'' is H, F, or CH₃;
Ar₁ is furan, thiophene, imidazole, thiazole, pyrazole, or pyridine; and
Ar₂ is phenyl.
- 20 16. A pharmaceutical composition for the treatment of infection or disease caused by a retrovirus, which comprises an amount of the compound of Claim 1 sufficient to provide an antivirally effective dosage of the compound in the range of about 1 to about 50 mg/kg-day or up to 3 g per day and a pharmaceutically effective carrier.
- 25 17. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a compound of Claim 1.
18. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a

-249-

compound of Claim 1 in combination with an HIV reverse transcriptase inhibitor.

19. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a compound of Claim 1 in combination with AZT.
20. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a compound of Claim 1 in combination with ddC.
21. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a compound of Claim 1 in combination with an HIV protease inhibitor.
22. A compound according to Claim 1 and selected from:
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-4-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-2-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-2-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

-250-

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

5 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

10 (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one;

15 6-[2-(4-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

3-(2-Amino-5-isopropyl-benzothiazol-6-yl-sulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-phenethyl-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

20 (S)-3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-yl-sulfanyl)-4-hydroxy-6-isopropyl-6-phenethyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

25 (S)-(6-{4-Hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-isopropyl-7-methyl-benzothiazol-2-yl)-carbamic acid methyl ester;

(S)-3-(2-Amino-6-tert-butyl-benzothiazol-4-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 (S)-3-(3-tert-Butyl-benzo[b]thiophen-2-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(3-tert-Butyl-benzo[b]thiophen-2-yl-sulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-251-

4-Hydroxy-3-(5-hydroxymethyl-2-isopropyl-thiophen-3-ylsulfanyl)-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

5 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

10 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

N-(4-{2-[5-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl]-ethyl}-thiazol-2-yl)-acetamide;

15 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

20 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

25 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

30 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-isopropyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

-252-

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

5 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

10 3-(2-tert-Butyl-4-hydroxy-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

15 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(1H-pyrazol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

20 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-[2-(1H-pyrazol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyrimidin-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

25 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyrimidin-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

6-[2-(2-Amino-pyrimidin-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

30 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(2-amino-pyrimidin-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiazol-4-yl-ethyl)-5,6-dihydro-pyran-2-one;

-253-

(S)-4-Hydroxy-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-6-phenethyl-5,6-dihydro-pyran-2-one;

(S)-4-Hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

5 3-(2-Amino-7-isopropyl-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

10 (S)-3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

(R)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(2-Amino-5-isopropyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

15 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

25 3-[2-tert-Butyl-4-(2-hydroxy-ethyl)-5-methyl-phenylsulfanyl]-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

5-Trifluoromethyl-pyridine-2-sulfonic acid {5-tert-butyl-4-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-2-methyl-phenyl}-amide;

30 5-Trifluoromethyl-pyridine-2-sulfonic acid {5-tert-butyl-4-[4-hydroxy-6-isopropyl-2-oxo-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-2-methyl-phenyl}-amide;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-254-

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

5 (S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

10 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

15 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

4-Hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

20 4-Hydroxy-3-(6-tert-butyl-indol-5-ylsulfanyl)-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

(S)-N-[6-tert-Butyl-5-(4-hydroxy-6-isopropyl-2-oxo-6-[2-(4-hydroxy-phenyl)ethyl]-5,6-dihydro-2H-pyran-3-ylsulfanyl)-indolin-1-yl]-(4-cyanophenyl)sulfonamide;

25 (S)-6-tert-Butyl-5-{4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-indole-1-carboxylic acid tert-butyl ester;

(S)-3-(6-tert-Butyl-indol-5-yl-sulfanyl)-4-hydroxy 6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 (S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-4-hydroxy-6-isopropyl-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

-255-

(S)-4-Hydroxy-6-isopropyl-3-(6-isopropyl-3H-benzimidazol-5-ylsulfanyl)-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

(S)-5-(4-Hydroxy-6-isopropyl-2-oxo-6-phenethyl-5,6-dihydro-2H-pyran-3-ylsulfanyl)-6-isopropyl-1,3-dihydro-benzimidazol-2-one;

5 (S)-3-(6-tert-Butyl-3H-benzimidazol-5-yl-sulfanyl)-4-hydroxy-6-isopropyl-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

(S)-5-tert-Butyl-6-(4-hydroxy-6-isopropyl-2-oxo-6-(2-phenethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl)-1,3-dihydro-benzimidazol-2-one;

10 (S)-N-[5-tert-Butyl-6-(4-hydroxy-6-isopropyl-2-oxo-6-phenethyl-5,6-dihydro-2H-pyran-3-ylsulfanyl)-benzothiazol-2-yl]-methanesulfonamide;

6-tert-Butyl-5-(4-hydroxy-6-isopropyl-2-oxo-6-phenethyl-5,6-dihydro-2H-pyran-3-ylsulfanyl)-1H-indole-2-carboxylic acid ethyl ester;

15 6-tert-Butyl-5-(4-hydroxy-6-isopropyl-2-oxo-6-phenethyl-5,6-dihydro-2H-pyran-3-ylsulfanyl)-1H-indole-2-carboxylic acid;

3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-[(2-(3-thiophenyl)-ethyl)-5,6-dihydro-pyran-2-one];

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3-fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3,5-difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

25 (S)-N-[5-tert-Butyl-6-(4-hydroxy-6-isopropyl-2-oxo-6-(2-phenethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl)-benzothiazol-2-yl]-acetamide;

(S)-N-{5-tert-Butyl-6-(4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl)-benzothiazol-2-yl}-4-cyanobenzenesulfonamide;

30 (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

-256-

6-[2-(2-Amino-thiazol-4-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

- 5 (S)-3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- (S)-3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-isopropyl-6-[(2-(3-thiophenyl)-ethyl)-5,6-dihydro-pyran-2-one]; and
- 4-(6-tert-Butyl-5-{4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-2,3-dihydro-indole-1-sulfonyl)-benzonitrile.
- 10

23. A compound according to Claim 1 and selected from:

- 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(3-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 15 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 20 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one
- 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 25 3-(2-Cyclopentyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 3-(4-Amino-2-cyclopentyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 30

-257-

3-(4-Amino-2-cyclopentyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

5 3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one; and

3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one.

24. A compound according to Claim 1 and selected from:

10 3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

15 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-furan-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(3-hydroxymethyl-furan-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-furan-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(2-methyl-furan-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

25 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(3-methyl-furan-2-yl)-ethyl]-5,6-dihydro-pyran-2-one; and

30 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-furan-3-yl)-ethyl]-5,6-dihydro-pyran-2-one.

25. A compound according to Claim 1 and selected from:

-258-

3-(2-Cyclopentyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one and

3-(6-tert-Butyl-1-hydroxy-indan-5-ylsulfanyl)-6-(2-furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one.

5 26. A compound according to Claim 1 and selected from:

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(4-hydroxymethyl-thiazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

10 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(5-hydroxymethyl-thiazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

15 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

20 6-[2-(2-Amino-thiazol-4-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(2-Amino-thiazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

25 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-4-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-isothiazol-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

30 and

-259-

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-[2-(4-methyl-oxazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one.

27. A compound according to Claim 1 and selected from:

5 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

10 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

15 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(1H-benzoimidazol-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(1H-Benzoimidazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

20 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(3-amino-1H-indazol-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
and

25 6-[2-(3-Amino-1H-indazol-5-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one.

28. A compound according to Claim 1 and selected from:

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-260-

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(1H-indazol-4-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-[2-(3-amino-1H-indazol-4-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

6-[2-(3-Amino-1H-indazol-4-yl)-ethyl]-3-(2-tert-butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one.

29. A compound according to Claim 1 and selected from:

3-(2-tert-Butyl-6-hydroxymethyl-5-methyl-pyridin-3-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-5-methyl-pyridin-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(3-tert-Butyl-2-hydroxymethyl-6-methyl-pyridin-4-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

3-(5-tert-Butyl-2-methyl-pyridin-4-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one.

30. A compound according to Claim 1 and selected from:

3-(4-tert-Butyl-5-hydroxymethyl-thiophen-3-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(4-tert-Butyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(4-Amino-phenyl)-ethyl]-3-(2-tert-butyl-5-methyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-5-methyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-261-

6-[2-(4-Amino-phenyl)-ethyl]-3-(4-tert-butyl-5-hydroxymethyl-thiophen-3-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(4-Amino-phenyl)-ethyl]-3-(3-tert-butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(3-tert-Butyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(3-tert-Butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

3-(3-tert-Butyl-4-hydroxymethyl-5-methyl-thiophen-2-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one.

31. A compound according to Claim 1 and selected from:

3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one.

32. A compound according to Claim 1 and selected from:

N-(7-tert-Butyl-6-{6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-acetamide;

-262-

N-(7-tert-Butyl-6-{4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-acetamide;

5 N-(7-tert-Butyl-6-{6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-methanesulfonamide;

N-(7-tert-Butyl-6-{4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-2-oxo-5,6-dihydro-2H-pyran-3-ylsulfanyl}-4-methyl-benzothiazol-2-yl)-methanesulfonamide;

10 3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

15 3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-6-[2-(4-fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

20 3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one.

33. A compound according to Claim 1 and selected from:

3-(2-tert-Butyl-6-hydroxymethyl-5-methyl-pyridin-3-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

25 3-(3-tert-Butyl-2-hydroxymethyl-6-methyl-pyridin-4-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(6-tert-Butyl-2,3-dihydro-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 3-(6-tert-Butyl-1H-indol-5-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-263-

3-(7-tert-Butyl-1,2,3,4-tetrahydro-quinolin-6-ylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

5 N-{7-tert-Butyl-6-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-4-methyl-benzothiazol-2-yl}-acetamide;

N-{7-tert-Butyl-6-[4-hydroxy-6-isopropyl-2-oxo-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-2H-pyran-3-ylsulfanyl]-4-methyl-benzothiazol-2-yl}-methanesulfonamide;

10 3-(7-tert-Butyl-2-dimethylamino-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(7-tert-Butyl-2-hydroxy-4-methyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

15 (S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-cyclohexyl-4-hydroxy-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

(S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-cyclopentyl-4-hydroxy-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

20 (S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-n-pentyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-cyclohexyl-4-hydroxy-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

25 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-6-cyclopentyl-4-hydroxy-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

30 3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-n-pentyl-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-cyclohexyl-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

-264-

3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-6-cyclopentyl-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

5 3-(4-Amino-2-tert-butyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-n-pentyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-4-hydroxy-6-n-pentyl-6-phenethyl-5,6-dihydro-pyran-2-one;

10 (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-cyclohexyl-4-hydroxy-6-phenethyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-cyclopentyl-4-hydroxy-6-phenethyl-5,6-dihydro-pyran-2-one;

(S)-4-Hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-6-propyl-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

15 (S)-6-Cyclohexyl-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

(S)-6-Cyclopentyl-4-hydroxy-6-[2-(4-hydroxy-phenyl)-ethyl]-3-(5-isopropyl-benzothiazol-6-ylsulfanyl)-5,6-dihydro-pyran-2-one;

20 (S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-4-hydroxy-6-propyl-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

(S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-6-cyclohexyl-4-hydroxy-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

(S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-6-cyclopentyl-4-hydroxy-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

25 (S)-3-(6-tert-Butyl-1H-indol-5-yl-sulfanyl)-4-hydroxy-6-pentyl-6-(2-phenethyl)-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-6-propyl-5,6-dihydro-pyran-2-one;

30 (S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-6-pentyl-5,6-dihydro-pyran-2-one;

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-cyclohexyl-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-5,6-dihydro-pyran-2-one; and

-265-

(S)-3-(2-Amino-5-tert-butyl-benzothiazol-6-ylsulfanyl)-6-cyclopentyl-6-[2-(3-fluoro-phenyl)-ethyl]-4-hydroxy-5,6-dihydro-pyran-2-one.

34. A compound according to Claim 1 named (S)-3-(2-tert-Butyl-4-hydroxymethyl-5-methyl-phenylsulfanyl)-4-hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one.
35. A compound selected from:
- 5-Hydroxy-6-methyl-5-[2-(4-methyl-thiazol-5-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-(2-thiazol-2-yl-ethyl)-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-(2-thiazol-4-yl-ethyl)-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-(2-thiazol-5-yl-ethyl)-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-5-[2-(5-methyl-thiazol-2-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-5-[2-(4-methyl-thiazol-2-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-5-[2-(2-methyl-thiazol-4-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-5-[2-(5-methyl-thiazol-4-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-5-[2-(2-isopropyl-thiazol-4-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-5-[2-(4-isopropyl-thiazol-5-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-5-[2-(5-isopropyl-thiazol-4-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-5-[2-(4-isopropyl-thiazol-2-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

-266-

- 5-[2-(2-Acetylamino-thiazol-4-yl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-(2-pyridin-2-yl-ethyl)-heptanoic acid methyl ester;
- 5 5-Hydroxy-6-methyl-3-oxo-5-(2-pyridin-4-yl-ethyl)-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-(2-pyridin-3-yl-ethyl)-heptanoic acid methyl ester;
- 5-(2-Furan-2-yl-ethyl)-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 10 5-(2-Furan-3-yl-ethyl)-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-[2-(tetrahydro-furan-2-yl)-ethyl]-heptanoic acid methyl ester;
- 15 5-Hydroxy-6-methyl-3-oxo-5-(2-thiophen-3-yl-ethyl)-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-(2-thiophen-2-yl-ethyl)-heptanoic acid methyl ester;
- 5-{2-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 20 5-{2-[5-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-{2-[5-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-2-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 25 5-Hydroxy-5-[2-(5-hydroxymethyl-thiophen-3-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-5-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-5-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 30 5-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

-267-

- 5-[2-(2-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-[2-(4-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5 5-Hydroxy-6-methyl-3-oxo-5-[2-(1-trityl-1H-pyrazol-3-yl)-ethyl]-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-[2-(1H-pyrrol-3-yl)-ethyl]-heptanoic acid methyl ester;
- 5-Hydroxy-5-[2-(1H-imidazol-4-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 10 5-Hydroxy-6-methyl-3-oxo-5-(2-pyrimidin-5-yl-ethyl)-heptanoic acid methyl ester;
- 5-[2-(2-Amino-pyrimidin-5-yl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 15 5-[2-(4-Cyano-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-5-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-{2-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-phenyl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 20 5-Hydroxy-5-[2-(1H-indol-5-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-3-oxo-5-{2-[2-(2,2,2-trifluoro-acetylamino)-thiazol-4-yl]-ethyl}-heptanoic acid methyl ester;
- 25 5-Hydroxy-6-methyl-5-[2-(4-methyl-thiophen-3-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;
- 5-Hydroxy-6-methyl-5-[2-(3-methyl-thiophen-2-yl)-ethyl]-3-oxo-heptanoic acid methyl ester;
- 5-{2-[3-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-2-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;
- 30 5-Hydroxy-5-[2-(3-hydroxymethyl-thiophen-2-yl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester;

-268-

5-[2-(2,6-Difluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

5-[2-(3,5-Difluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

5 5-[2-(2,4-Difluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

5-[2-(3,4,5-Trifluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

10 5-[2-(5-Fluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

5-[2-(2-Ethyl-5-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester;

5-{2-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-4-fluoro-phenyl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid methyl ester; and

15 5-Hydroxy-5-[2-(4-fluoro-2-hydroxymethyl-phenyl)-ethyl]-6-methyl-3-oxo-heptanoic acid methyl ester.

36. A compound selected from:

4-Hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;

20 4-Hydroxy-6-isopropyl-6-(2-thiazol-2-yl-ethyl)-5,6-dihydro-pyran-2-one;

4-Hydroxy-6-isopropyl-6-(2-thiazol-4-yl-ethyl)-5,6-dihydro-pyran-2-one;

25 4-Hydroxy-6-isopropyl-6-(2-thiazol-5-yl-ethyl)-5,6-dihydro-pyran-2-one;

4-Hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

4-Hydroxy-6-isopropyl-6-[2-(4-methyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;

30 4-Hydroxy-6-isopropyl-6-[2-(2-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;

-269-

- 4-Hydroxy-6-isopropyl-6-[2-(5-methyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-[2-(2-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 5 4-Hydroxy-6-isopropyl-6-[2-(4-isopropyl-thiazol-5-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-[2-(5-isopropyl-thiazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-[2-(4-isopropyl-thiazol-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 10 N-{4-[2-(4-Hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl)-ethyl]-thiazol-2-yl}-acetamide;
- 4-Hydroxy-6-isopropyl-6-(2-pyridin-2-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 15 4-Hydroxy-6-isopropyl-6-(2-pyridin-4-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-(2-pyridin-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 6-(2-Furan-2-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 20 6-(2-Furan-3-yl-ethyl)-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-[2-(tetrahydro-furan-2-yl)-5,6-dihydro-pyran-2-one];
- 25 4-Hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-(2-thiophen-2-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-[2-(5-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 30 4-Hydroxy-6-[2-(5-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-270-

- 4-Hydroxy-6-[2-(2-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 6-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 5 6-[2-(2-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 6-[2-(4-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-[2-(1H-pyrazol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 10 4-Hydroxy-6-isopropyl-6-[2-(1H-pyrrol-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-[2-(1-trityl-1H-imidazol-4-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 15 4-Hydroxy-6-isopropyl-6-(2-pyrimidin-5-yl-ethyl)-5,6-dihydro-pyran-2-one;
- 6-[2-(2-Amino-pyrimidin-5-yl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;
- 4-[2-(4-Hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl)-ethyl]-benzonitrile;
- 20 4-Hydroxy-6-isopropyl-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-[2-(1H-indol-5-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;
- 25 2,2,2-Trifluoro-N-{4-[2-(4-hydroxy-2-isopropyl-6-oxo-3,6-dihydro-2H-pyran-2-yl)-ethyl]-thiazol-2-yl}-acetamide;
- 4-Hydroxy-6-isopropyl-6-[2-(4-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 4-Hydroxy-6-isopropyl-6-[2-(3-methyl-thiophen-2-yl)-ethyl]-5,6-dihydro-pyran-2-one;
- 30 4-Hydroxy-6-[2-(3-hydroxymethyl-thiophen-2-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

-271-

6-[2-(2,6-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(3,5-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

5 6-[2-(2,4-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(3,4,5-Trifluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

10 6-[2-(5-Fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(2-Ethyl-5-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-{2-[2-(tert-Butyl-dimethyl-silanyloxymethyl)-4-fluoro-phenyl]-ethyl}-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

15 6-[2-(4-Fluoro-2-hydroxymethyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one.

37. A compound selected from:

(S)-5-Hydroxy-6-methyl-3-oxo-5-(2-thiophen-3-yl-ethyl)-heptanoic acid ethyl ester;

20 (R)-5-Hydroxy-6-methyl-3-oxo-5-(2-thiophen-3-yl-ethyl)-heptanoic acid ethyl ester;

5-[2-(3-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester, (S) enantiomer;

25 (R)-5-[2-(3-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

5-[2-(3,4-Difluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester (R and S isomers);

5-Hydroxy-6-methyl-3-oxo-5-[2-(2-trifluoromethyl-phenyl)-ethyl]-heptanoic acid ethyl ester;

30 5-Hydroxy-6-methyl-3-oxo-5-[2-(3-trifluoromethyl-phenyl)-ethyl]-heptanoic acid ethyl ester;

-272-

5-Hydroxy-6-methyl-3-oxo-5-[2-(4-trifluoromethyl-phenyl)-ethyl]-heptanoic acid ethyl ester;

5-[2-(3-Fluoro-5-trifluoromethyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

5 [2-(4-Fluoro-3-trifluoromethyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

(S)-5-Hydroxy-5-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-methyl-3-oxo-heptanoic acid ethyl ester;

10 (R)-5-Hydroxy-5-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-methyl-3-oxo-heptanoic acid ethyl ester;

5-Hydroxy-6-methyl-3-oxo-5-(2-phenyl-ethyl)-heptanoic acid ethyl ester (R and S isomers);

5-[2-(4-Fluoro-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester (S and R isomers);

15 5-{2-[4-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-ethyl}-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;


(S)-5-Hydroxy-6-methyl-5-[2-(2-methyl-thiophen-3-yl)-ethyl]-3-oxo-heptanoic acid ethyl ester;

20 (S)-5-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester;

(S)-5-[2-(3,5-Difluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester; and

(S)-5-[2-(4-Fluoro-2-methyl-phenyl)-ethyl]-5-hydroxy-6-methyl-3-oxo-heptanoic acid ethyl ester.

25 38. A compound of Formula 1A



1A

-273-

wherein

R₄, R₅, and R₆ are independently H, a straight or branched alkyl of

1-6 carbons, a cycloalkyl of 3-6 carbons, (CR'₂)_nOR,

(CR'₂)_nN(R)₂, F, Cl, Br, CN, CF₃, =O, (CR'₂)_pNR'COR,

5 (CR'₂)_pSO_pR, (CR'₂)_pR, (CR'₂)_pOCOR, O(CR'₂)_mOR,

NR(CR'₂)_mOR, (CR'₂)_pNR'CON(R)₂, (CR'₂)_pOCON(R)₂,

(CR'₂)_pNR'CO₂R, (CR'₂)_pCOR, (CR'₂)_pCO₂R,

(CR'₂)_pCON(R)₂, (CR'₂)_pNR'SO₂R, (CR'₂)_pSO₂N(R)₂,

(CR'₂)_pNR'SO₂OR, (CR'₂)_pOSO₂N(R)₂, (CR'₂)_pNR'SO₂N(R)₂,

10 (CR'₂)_pC(NR')N(R)₂, (CR'₂)_pNR'C(NR')N(R)₂, (CR'₂)_pHet;

any two of R₄-R₆ may together form a ring of 5-6 total atoms which may
contain 0-3 heteroatoms;

n is an integer of from 0 to 3;

p is an integer from 0 to 2;

15 R₇ is a straight or branched alkyl of 1-6 carbons or a carbocycle of
3-6 carbons;

R is independently H, a straight or branched alkyl of 1-4 carbons,

(CH₂)_nPh, or a (CH₂)_nheterocycle of 5-6 atoms containing

1-2 heteroatoms and wherein the (R)₂ in N(R)₂ may form a

20 heterocycle containing the nitrogen, all optionally substituted by F,
Cl, Br, OR', CN, CO₂R', N(R')₂, NR'COR', CF₃, or =O;

R' is independently H, a straight or branched alkyl of 1-4 carbons, or
phenyl;

R'' is independently H, a straight or branched alkyl of 1-4 carbons, F, Cl,

25 Br, OR', or N(R')₂;

Ar₂ is phenyl or Het wherein Het is a heterocycle of from 5-6 atoms
having from 1-4 heteroatoms.

39. A compound selected from:

2-Amino-5-tert-butyl-benzothiazole-6-thiol;

-274-

- 2-Amino-5-isopropyl-benzothiazole-6-thiol;
 5-Isopropyl-benzothiazole-6-thiol;
 2-Amino-7-isopropyl-4-methyl-benzothiazole-6-thiol;
 2-Amino-6-tert-butyl-benzothiazole-4-thiol;
 5 2-(Methyloxycarbonyl)amino-4-isopropyl-7-methyl-benzothiazole-6-thiol;
 3-tert-Butyl-2-mercapto-benzo[b]thiophene;
 5-(tert-Butyl-dimethyl-silanyloxymethyl)-2-isopropyl-3-mercapto-thiophene;
 10 (5-tert-Butyl-4-mercapto-5-methyl-phenyl)methanol;
 (3-tert-Butyl-4-mercapto-phenyl)methanol; and
 2-(5-tert-Butyl-4-mercapto-2-methyl-phenyl)-ethanol.
40. A compound selected from:
- 15 Toluene-4-thiosulfonic acid S-(2-amino-5-tert-butyl-benzothiazol-6-yl)ester;
 Toluene-4-thiosulfonic acid S-(2-amino-5-isopropyl-benzothiazol-6-yl)ester;
 Toluene-4-thiosulfonic acid S-(5-isopropyl-benzothiazol-6-yl)ester;
 Toluene-4-thiosulfonic acid S-(2-amino-7-isopropyl-4-methyl-
 20 benzothiazol-6-yl)ester;
 Toluene-4-thiosulfonic acid S-(4-isopropyl-2-methoxycarbonylamino-7-methyl-benzothiazol-6-yl)ester;
 Toluene-4-thiosulfonic acid S-(2-amino-6-tert-butyl-benzothiazol-4-yl)ester;
 25 Toluene-4-thiosulfonic acid S-(3-tert-butyl-benzo[b]thiophen-2-yl)ester;
 Toluene-4-thiosulfonic acid S-[5-(tert-butyl-silanyloxymethyl)-2-isopropyl-thiophen-3-yl]ester;
 Toluene-4-thiosulfonic acid S-(2-tert-butyl-4-hydroxymethyl-
 30 phenyl)ester;
 Toluene-4-thiosulfonic acid S-[2-tert-butyl-4-(2-hydroxy-ethyl)-5-methyl-phenyl]ester;

-275-

Benzenesulfonylthioic acid, 4-methyl-, S-[2-(1,1-dimethylethyl)-5-methyl-4-[[[(5-trifluoromethyl)-2-pyridinyl]sulfonyl]amino]phenyl]ester;

6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-indole-1-carboxylic acid tert-butyl ester;

5 6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-2,3-dihydro-indole-1-carboxylic acid tert-butyl ester;

Toluene-4-thiosulfonic acid S-[6-tert-butyl-1-(4-cyanobenzenesulfonyl)-2,3-dihydro-1H-indol-5-yl] ester;

10 Toluene-4-thiosulfonic acid S-(6-isopropyl-3H-benzoimidazol-5-yl) ester;

Toluene-4-thiosulfonic acid S-(6-tert-butyl-3H-benzoimidazol-5-yl) ester;

Toluene-4-thiosulfonic acid S-(6-tert-butyl-2-oxo-2,3-dihydro-1H-benzimidazol-5-yl) ester;

15 6-tert-Butyl-5-(toluene-4-sulfonylsulfanyl)-1H-indole-2-carboxylic acid ethyl ester;

Toluene-4-thiosulfonic acid S-(7-tert-butyl-2,3-dioxo-1,2,3,4-tetrahydro-quinoxalin-6-yl) ester;

20 Toluene-4-thiosulfonic acid S-[5-tert-butyl-2-(4-cyanobenzenesulfonyl-amino)-benzothiazol-6-yl]] ester;

Toluene-4-thiosulfonic acid S-(6-tert-butyl-2-methanesulfonylamino-benzothiazol-6-yl) ester;

Toluene-4-thiosulfonic acid S-(2-acetylamino-5-tert-butyl-benzothiazol-6-yl) ester; and

25 Toluene-4-thiosulfonic acid S-(6-isopropyl-2-oxo-2,3-dihydro-1H-benzimidazol-5-yl) ester.

41. A compound selected from:

3-Hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid tert-butyl ester;

30 5-(3-Fluoro-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;

-276-

- 5-(3,4-Difluoro-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;
- 3-Hydroxy-3-isopropyl-5-(2-trifluoromethyl-phenyl)-pent-4-enoic acid benzyl ester;
- 5 3-Hydroxy-3-isopropyl-5-(3-trifluoromethyl-phenyl)-pent-4-enoic acid benzyl ester;
- 3-Hydroxy-3-isopropyl-5-(4-trifluoromethyl-phenyl)-pent-4-enoic acid benzyl ester;
- 10 5-(3-Fluoro-5-trifluoromethyl-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;
- 5-(4-Fluoro-3-trifluoromethyl-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;
- 3-Hydroxy-3-[2-(2-hydroxymethyl-phenyl)-ethyl]-4-methyl-pentanoic acid tert-butyl ester;
- 15 3-Hydroxy-4-methyl-3-(2-phenyl-ethyl)-pentanoic acid tert-butyl ester;
- 3-Hydroxy-3-isopropyl-5-phenyl-3-pent-4-enoic acid benzyl ester;
- 5-(4-Fluoro-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;
- 20 3-{2-[4-(tert-Butyl-dimethyl-silanyloxymethyl)-thiophen-3-yl]-ethyl}-3-hydroxy-4-methyl-pentanoic acid ethyl ester;
- 3-Hydroxy-4-methyl-3-[2-(2-methyl-thiophen-3-yl)-ethyl]-pentanoic acid tert-butyl ester;
- 5-(3-Fluoro-2-methyl-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester;
- 25 5-(3,5-Difluoro-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid benzyl ester; and
- 5-(4-Fluoro-2-methyl-phenyl)-3-hydroxy-3-isopropyl-pent-4-enoic acid tert-butyl ester.
- 30 42. A compound selected from:
- (S)-3-Hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid;
- (R)-3-Hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid;

-277-

- (+/-)-3-Hydroxy-4-methyl-3-(2-thiophen-3-yl-ethyl)-pentanoic acid;
- 3-[2-(3-Fluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;
- 3-[2-(3,4-Difluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;
- 3-Hydroxy-4-methyl-3-[2-(2-trifluoromethyl-phenyl)-ethyl]-pentanoic acid;
- 3-Hydroxy-4-methyl-3-[2-(3-trifluoromethyl-phenyl)-ethyl]-pentanoic acid;
- 3-Hydroxy-4-methyl-3-[2-(4-trifluoromethyl-phenyl)-ethyl]-pentanoic acid;
- 3-[2-(3-Fluoro-5-trifluoromethyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;
- 3-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;
- (S)-3-Hydroxy-3-[2-(2-hydroxymethyl-phenyl)-ethyl]-4-methyl-pentanoic acid;
- (R)-3-Hydroxy-3-[2-(2-hydroxymethyl-phenyl)-ethyl]-4-methyl-pentanoic acid;
- 3-Hydroxy-4-methyl-3-(2-phenyl-ethyl)-pentanoic acid;
- 3-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;
- 3-{2-[4-(tert-Butyl-dimethyl-silanyloxy-methyl)-thiophen-3-yl]-ethyl}-3-hydroxy-4-methyl-pentanoic acid;
- 3-Hydroxy-4-methyl-3-[2-(2-methyl-thiophen-3-yl)-ethyl]-pentanoic acid;
- 3-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid;
- 3-[2-(3,5-Difluoro-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid; and
- 3-[2-(4-Fluoro-2-methyl-phenyl)-ethyl]-3-hydroxy-4-methyl-pentanoic acid.

43. A compound selected from:

-278-

(S)-4-Hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

(R)-4-Hydroxy-6-isopropyl-6-(2-thiophen-3-yl-ethyl)-5,6-dihydro-pyran-2-one;

5 6-[2-(3-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (R and S isomers);

6-[2-(3,4-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (R and S isomers);

10 4-Hydroxy-6-isopropyl-6-[2-(2-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

4-Hydroxy-6-isopropyl-6-[2-(3-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

4-Hydroxy-6-isopropyl-6-[2-(4-trifluoromethyl-phenyl)-ethyl]-5,6-dihydro-pyran-2-one;

15 6-[2-(3-Fluoro-5-trifluoromethyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

6-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

20 (S)-4-Hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

(R)-4-Hydroxy-6-[2-(2-hydroxymethyl-phenyl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

4-Hydroxy-6-isopropyl-6-(2-phenyl-ethyl)-5,6-dihydro-pyran-2-one (R and S isomers);

25 6-[2-(4-Fluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one (R and S isomers);

4-Hydroxy-6-[2-(4-hydroxymethyl-thiophen-3-yl)-ethyl]-6-isopropyl-5,6-dihydro-pyran-2-one;

30 (S)-4-Hydroxy-6-isopropyl-6-[2-(2-methyl-thiophen-3-yl)-ethyl]-5,6-dihydro-pyran-2-one;

(S)-6-[2-(3-Fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one;

-279-

(S)-6-[2-(3,5-Difluoro-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one; and

(S)-6-[2-(4-Fluoro-2-methyl-phenyl)-ethyl]-4-hydroxy-6-isopropyl-5,6-dihydro-pyran-2-one.

- 5 44. A compound and the R and S isomer thereof selected from:
- 3-[2-(3-Fluoro-phenyl)-ethyl]-3-hydroxy-1-imidazol-1-yl-4-methyl-pentan-1-one;
- 3-Hydroxy-1-imidazol-1-yl-4-methyl-3-phenethyl-pentan-1-one;
- 3-[2-(3,4-Difluoro-phenyl)-ethyl]-3-hydroxy-1-imidazol-1-yl-4-methyl-pentan-1-one;
- 10 3-Hydroxy-1-imidazol-1-yl-4-methyl-3-[2-(2-trifluoromethyl-phenyl)-ethyl]-pentan-1-one;
- 3-[2-(4-Fluoro-3-trifluoromethyl-phenyl)-ethyl]-3-hydroxy-1-imidazol-1-yl-4-methyl-pentan-1-one;
- 15 3-Hydroxy-1-imidazol-1-yl-4-methyl-3-[2-(3-trifluoromethyl-phenyl)-ethyl]-pentan-1-one;
- 3-[2-(4-Fluoro-2-trifluoromethyl-phenyl)-ethyl]-3-hydroxy-1-imidazol-1-yl-4-methyl-pentan-1-one;
- 3-Hydroxy-1-imidazol-1-yl-4-methyl-3-[2-(4-trifluoromethyl-phenyl)-ethyl]-pentan-1-one;
- 20 3-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-1-imidazol-1-yl-4-methyl-pentan-1-one;
- 3-Hydroxy-1-imidazol-1-yl-4-methyl-3-[2-thiophen-3-yl-ethyl]-pentan-1-one; and
- 25 3-Hydroxy-3-[2-(2-hydroxymethyl-phenyl)-ethyl]-1-imidazol-1-yl-4-methyl-pentan-1-one.
45. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a compound according to Claim 1 and another antiretroviral agent.

-280-

46. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a compound of Claim 1 in combination with AZT and another antiretroviral agent.
- 5 47. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a compound of Claim 1 in combination with ddC and another antiretroviral agent.
- 10 48. A method of treatment of infection or disease caused by a retrovirus, which comprises administering to a subject in need of such treatment a compound of Claim 1 in combination with an HIV protease inhibitor and another antiretroviral agent.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 405/06, A61K 31/35, C07D 407/06, 409/06, 417/12, 409/12, 409/14, 417/06, 405/14, 405/12, 417/14, 413/06	A3	(11) International Publication Number: WO 00/15634 (43) International Publication Date: 23 March 2000 (23.03.00)
(21) International Application Number: PCT/US99/18986 (22) International Filing Date: 18 August 1999 (18.08.99) (30) Priority Data: 60/099,946 11 September 1998 (11.09.98) US (71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BOYER, Frederick, Earl, Jr. [US/US]; 1624 Ironwood Drive, Canton Township, MI 48188 (US). DOMAGALA, John, Michael [US/US]; 47693 Red Run, Canton, MI 48187 (US). ELLSWORTH, Edmund, Lee [US/US]; 6899 Kestrel Court, Brighton, MI 48116 (US). GAJDA, Christopher, Andrew [CA/US]; 2684 Bellwood, Ann Arbor, MI 48104 (US). HAGEN, Susan, Elizabeth [US/US]; 1035 Paddington Road, Canton Township, MI 48187 (US). LOVDAHL, Michael, James [US/US]; 2948 Renfrew Street, Ann Arbor, MI 48105 (US). LUNNEY, Elizabeth, Ann [US/US]; 619 Ridgewood Court, Ann Arbor, MI 48103 (US). MARKOSKI, Larry, James [US/US]; 303 W. Green 202D, Champaign, IL 61820 (US).	JOSYULA, Vara, Prasad, Venkata, Nagendra [IN/US]; 3129 Fawnmeadow Court, Ann Arbor, MI 48105 (US). TAIT, Bradley, Dean [US/US]; 8431 Congress Drive, Canton, MI 48187 (US). (74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al. (81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 16 November 2000 (16.11.00)	
(54) Title: HIV PROTEASE INHIBITORS (57) Abstract The present invention relates to novel dihydropyrones with tethered heterocycles having improved pharmacologic properties which potently inhibit the HIV aspartyl protease blocking HIV infectivity. The dihydropyrones are useful in the development of therapies for the treatment of viral infections and diseases, including AIDS. The present invention is also directed to methods of synthesis of the dihydropyrones and intermediates useful in the preparation of the final compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

In national Application No

PCT/US 99/18986

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D405/06 A61K31/35 C07D407/06 C07D409/06 C07D417/12
C07D409/12 C07D409/14 C07D417/06 C07D405/14 C07D405/12
C07D417/14 C07D413/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, Y	WO 95 14012 A (PARKE DAVIS & CO) 26 May 1995 (1995-05-26) see in particular examples on pages 15,16,18,19,23 and example 133 ---	1-48
X, Y	WO 95 14011 A (PARKE DAVIS & CO) 26 May 1995 (1995-05-26) see in particular examples on pages 15,16,18,19,24 and example 133 ---	1-48
X, Y	WO 98 19997 A (BOYER FREDERICK EARL JR ;DOMAGALA JOHN MICHAEL (US); HAGEN SUSAN E) 14 May 1998 (1998-05-14) see in particular examples 61 and 84 --- -/--	1-48

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

26 April 2000

Date of mailing of the international search report

22.05.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Steendijk, M

INTERNATIONAL SEARCH REPORT

In. ational Application No

PCT/US 99/18986

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 84, no. 21, 24 May 1976 (1976-05-24) Columbus, Ohio, US; abstract no. 150164, GUETTE ET AL.: "Preparation of amino hydroxyesters by reaction of ..." XP002126823 & Bull.Soc.Chim.Fr. 1975, (11-12, pt.2), 2759-2762 abstract	38
X	--- HAGEN ET AL.: "Synthesis of 5,6-dihydro-4-hydroxy-2-pyrones ...w" J.MED.CHEM., vol. 40, no. 23, 1997, pages 3707-3711, XP002136420 see schemes 1 and 2	38-40
A	--- US 3 786 063 A (ARNOLD R) 15 January 1974 (1974-01-15) see formula column 1	39,40
P,X	--- ELLSWORTH E L ET AL: "4-hydroxy-5,6-dihydro-2H-pyran-2-ones. 3. bicyclic and heteroaromatic systems as 3-position scaffolds to the S1' and S2' of the HIV-1 protease enzyme" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS,GB,OXFORD, vol. 9, no. 14, 19 July 1999 (1999-07-19), pages 2019-2024, XP004171629 ISSN: 0960-894X the whole document	1,2,6,8, 10,11, 15-22, 29-33, 38-40, 45-48
P,X	--- VARA PRASAD J V N ET AL: "Nonpeptidic HIV protease inhibitors: 6-alkyl-5,6-dihydropyran-2-ones possessing achiral 3-(4-amino/carboxamide-2-t-butyl, 5-methylphenyl thio) moiety: antiviral activities and pharmacokinetic properties" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS,GB,OXFORD, vol. 9, no. 11, 7 June 1999 (1999-06-07), pages 1481-1486, XP004169615 ISSN: 0960-894X table 1	38-40

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/18986

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-2(part),3-5,7,8(part),9,12-14,16-22(part),23-28,30(part),33(part)34,35-38(part),41-48(part)

Endproducts and related intermediates in which Ar2 occurs as a heterocycle as well as their application

2. Claims: 1-2(part),6,8(part),10,11,15,16-22(part),29,30(part),31,32,33(part),39-40(part),45-48(part)

Endproducts and related intermediates in which Ar1 occurs as a heterocycle (and Ar2 does not represent a heterocycle),as well as their application

3. Claims: 35-38(part),41-44(part)

Intermediates in which Ar2 occurs and does not represent a heterocycle

4. Claims: 39-40(part)

Intermediates in which Ar1 occurs and does not represent a heterocycle

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/US 99/18986

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9514012 A	26-05-1995	US 5840751 A	24-11-1998
		AU 695088 B	06-08-1998
		AU 4097497 A	08-01-1998
		AU 680064 B	17-07-1997
		AU 8090094 A	06-06-1995
		AU 689158 B	26-03-1998
		AU 8125694 A	06-06-1995
		BG 100563 A	31-12-1996
		CA 2176041 A	26-05-1995
		CA 2176043 A	26-05-1995
		CZ 9601368 A	17-12-1997
		EP 0729463 A	04-09-1996
		EP 0729464 A	04-09-1996
		FI 962021 A	12-07-1996
		FI 962022 A	12-07-1996
		HR 940935 A	30-04-1997
		HR 940936 A	31-08-1997
		HU 77668 A	28-07-1998
		HU 75225 A	28-04-1997
		IL 111673 A	11-04-1999
		JP 9505292 T	27-05-1997
		JP 9505294 T	27-05-1997
		NO 962017 A	08-07-1996
		NO 962018 A	04-07-1996
		NZ 275318 A	29-06-1999
		PL 314483 A	16-09-1996
		SK 64696 A	05-03-1997
		US 5936128 A	10-08-1999
		WO 9514011 A	26-05-1995
		US 5789440 A	04-08-1998
		ZA 9409148 A	21-07-1995
		ZA 9409151 A	29-07-1995
WO 9514011 A	26-05-1995	US 5789440 A	04-08-1998
		AU 695088 B	06-08-1998
		AU 4097497 A	08-01-1998
		AU 680064 B	17-07-1997
		AU 8090094 A	06-06-1995
		AU 689158 B	26-03-1998
		AU 8125694 A	06-06-1995
		BG 100563 A	31-12-1996
		CA 2176041 A	26-05-1995
		CA 2176043 A	26-05-1995
		CZ 9601368 A	17-12-1997
		EP 0729463 A	04-09-1996
		EP 0729464 A	04-09-1996
		FI 962021 A	12-07-1996
		FI 962022 A	12-07-1996
		HR 940935 A	30-04-1997
		HR 940936 A	31-08-1997
		HU 77668 A	28-07-1998
		HU 75225 A	28-04-1997
		IL 111673 A	11-04-1999
		JP 9505292 T	27-05-1997
		JP 9505294 T	27-05-1997
		NO 962017 A	08-07-1996
		NO 962018 A	04-07-1996
		NZ 275318 A	29-06-1999

INTERNATIONAL SEARCH REPORT

Information on patent family members

In tional Application No

PCT/US 99/18986

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9514011 A		PL 314483 A	16-09-1996
		SK 64696 A	05-03-1997
		US 5936128 A	10-08-1999
		WO 9514012 A	26-05-1995
		US 5840751 A	24-11-1998
		ZA 9409148 A	21-07-1995
		ZA 9409151 A	29-07-1995
WO 9819997 A	14-05-1998	US 5834506 A	10-11-1998
		AU 5100198 A	29-05-1998
		EP 0935597 A	18-08-1999
		HR 970571 A	31-08-1998
		NO 992103 A	30-04-1999
		PL 332904 A	25-10-1999
US 3786063 A	15-01-1974	NONE	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.